



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 01:19 AM UTC

PDB ID : 3H2Y / pdb_00003h2y
Title : Crystal structure of YqeH GTPase from Bacillus anthracis with dGDP bound
Authors : Brunzelle, J.S.; Anderson, S.M.; Xu, X.; Savchenko, A.; Anderson, W.F.;
Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2009-04-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	24% 68% 15% • 16%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 2587 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GTPase family protein.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	309	2434	1561	414	451	3	5	17	2	0

- Molecule 2 is 2'-DEOXYGUANOSINE-5'-DIPHOSPHATE (CCD ID: DGI) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	27	10	5	10	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	126	Total	O	0	0
			126	126		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GTPase family protein



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	83.13Å 58.99Å 77.72Å 90.00° 99.38° 90.00°	Depositor
Resolution (Å)	30.00 – 1.80 30.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.4 (30.00-1.80) 98.4 (30.00-1.80)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.67 (at 1.80Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.215 , 0.242 0.213 , 0.242	Depositor DCC
R_{free} test set	1714 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.546	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 69.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	2587	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.47% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DGI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	2/2485 (0.1%)	0.85	7/3359 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	331	LYS	CG-CD	-11.67	1.17	1.52
1	A	328	ARG	CB-CG	-8.26	1.27	1.52

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	328	ARG	CA-CB-CG	9.29	132.68	114.10
1	A	331	LYS	CB-CG-CD	8.45	130.74	111.30
1	A	328	ARG	CB-CG-CD	6.79	126.92	111.30
1	A	205	ILE	N-CA-C	6.23	117.68	109.21
1	A	365	LYS	CA-CB-CG	5.83	125.75	114.10
1	A	335	VAL	N-CA-C	5.52	115.74	107.51
1	A	197	PHE	N-CA-C	-5.49	102.86	109.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2434	0	2400	34	1
2	A	27	0	12	0	0
3	A	126	0	0	4	1
All	All	2587	0	2412	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:227:MSE:HE2	1:A:359:VAL:HG11	1.72	0.70
1:A:265:ARG:HB3	1:A:354:HIS:HB2	1.76	0.65
1:A:324:GLU:HG2	1:A:352:VAL:HG22	1.83	0.60
1:A:181:ILE:HD11	1:A:205:ILE:HD13	1.83	0.58
1:A:62:ILE:HG21	1:A:197:PHE:CE1	2.44	0.53
1:A:80:PHE:CE2	1:A:108:LEU:HB2	2.44	0.53
1:A:235:SER:O	1:A:239:ILE:HD12	2.09	0.53
1:A:355:VAL:HG22	1:A:356:PRO:HD2	1.93	0.50
1:A:355:VAL:HG22	1:A:359:VAL:HB	1.95	0.49
1:A:168[B]:CYS:SG	1:A:223:ASN:ND2	2.85	0.49
1:A:207:ILE:HD12	1:A:215:LEU:HD22	1.95	0.48
1:A:254:GLU:HG2	1:A:255:GLU:HG2	1.96	0.47
1:A:307:SER:HA	1:A:308:PRO:C	2.39	0.47
1:A:67:GLY:HA2	1:A:97:ASN:HD21	1.79	0.46
1:A:368:ILE:HD11	3:A:458:HOH:O	2.14	0.46
1:A:206:ASP:HB3	3:A:465:HOH:O	2.14	0.46
1:A:331:LYS:HB3	1:A:366:SER:HB2	1.97	0.46
1:A:289:THR:HG21	1:A:297:LEU:HD22	1.96	0.46
1:A:368:ILE:HD12	1:A:368:ILE:N	2.31	0.45
1:A:87:LEU:CD2	1:A:168[B]:CYS:SG	3.05	0.45
1:A:187:GLU:HG3	3:A:493:HOH:O	2.16	0.45
1:A:263:LEU:HD22	1:A:317:MSE:HE1	2.00	0.44
1:A:313:GLU:O	1:A:317:MSE:HB2	2.18	0.44
1:A:317:MSE:HE2	1:A:356:PRO:HG2	2.00	0.44
1:A:254:GLU:HG3	1:A:270:SER:HA	2.00	0.43
1:A:75:LYS:NZ	1:A:87:LEU:H	2.17	0.43
1:A:197:PHE:O	1:A:198:PRO:C	2.62	0.42
1:A:246:LYS:HA	1:A:247:PRO:HD3	1.90	0.42
1:A:113:VAL:HG13	1:A:367:LEU:HB2	2.01	0.42
1:A:248:MSE:HB2	1:A:278:CYS:SG	2.60	0.42
1:A:145:GLN:HB2	3:A:459:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:192:ILE:HD13	1:A:205:ILE:HD11	2.01	0.41
1:A:303:GLY:HA2	1:A:309:PRO:O	2.19	0.41
1:A:277:THR:HG22	1:A:279:HIS:CE1	2.57	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:92:ARG:NH2	3:A:493:HOH:O[4_455]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	307/368 (83%)	298 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/318 (83%)	262 (99%)	3 (1%)	65	60

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	LEU
1	A	320	LEU
1	A	328	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	ASN
1	A	170	ASN
1	A	223	ASN
1	A	354	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	DGI	A	369	-	28,29,29	1.13	2 (7%)	43,45,45	1.73	7 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DGI	A	369	-	-	0/16/28/28	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	369	DGI	C5-C4	3.01	1.47	1.38
2	A	369	DGI	C4-N9	-2.12	1.32	1.38

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	369	DGI	C5-C4-N3	-5.47	119.68	128.39
2	A	369	DGI	C2-N3-C4	4.89	120.72	112.30
2	A	369	DGI	N9-C4-N3	4.09	134.14	125.95
2	A	369	DGI	C6-C5-N7	3.62	136.87	130.29
2	A	369	DGI	C4-C5-N7	-2.51	106.69	110.67
2	A	369	DGI	O2A-PA-O3A	2.29	113.45	107.27
2	A	369	DGI	O6-C6-C5	-2.08	121.05	126.53


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	304/368 (82%)	1.72	89 (29%) 1 1	27, 54, 69, 73	6 (1%)

All (89) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	327	ILE	6.3
1	A	221	ILE	5.5
1	A	338	GLY	5.0
1	A	59[A]	PHE	4.8
1	A	355	VAL	4.7
1	A	331	LYS	4.5
1	A	361	VAL	4.4
1	A	320	LEU	4.3
1	A	359	VAL	4.2
1	A	337	SER	4.1
1	A	360	SER	4.1
1	A	368	ILE	4.1
1	A	339	LEU	4.1
1	A	348	GLY	4.0
1	A	340	GLY	3.8
1	A	205	ILE	3.8
1	A	80	PHE	3.8
1	A	356	PRO	3.7
1	A	197	PHE	3.6
1	A	63	LEU	3.5
1	A	325	PHE	3.3
1	A	291	LEU	3.3
1	A	96	ASN	3.3
1	A	297	LEU	3.2
1	A	226	GLN	3.2
1	A	245	ILE	3.2
1	A	238	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	353	ALA	3.2
1	A	224	HIS	3.1
1	A	79	ILE	3.1
1	A	272	GLY	3.1
1	A	76	ILE	3.0
1	A	335	VAL	3.0
1	A	206	ASP	3.0
1	A	342	VAL	3.0
1	A	336	PHE	3.0
1	A	222	ILE	3.0
1	A	94	VAL	3.0
1	A	358	GLY	2.9
1	A	341	TRP	2.9
1	A	236	LEU	2.9
1	A	330	PRO	2.8
1	A	367	LEU	2.8
1	A	93	PHE	2.8
1	A	276	PHE	2.8
1	A	223	ASN	2.7
1	A	172	GLY	2.7
1	A	198	PRO	2.7
1	A	278	CYS	2.6
1	A	215	LEU	2.6
1	A	109	ILE	2.6
1	A	77	VAL	2.6
1	A	323	TYR	2.6
1	A	328	ARG	2.5
1	A	263	LEU	2.5
1	A	207	ILE	2.5
1	A	235	SER	2.5
1	A	74	VAL	2.5
1	A	62	ILE	2.5
1	A	321	VAL	2.5
1	A	346	GLU	2.5
1	A	204	LEU	2.5
1	A	250	PHE	2.5
1	A	171	VAL	2.5
1	A	196	HIS	2.4
1	A	81	ASP	2.4
1	A	145	GLN	2.4
1	A	191	VAL	2.4
1	A	211	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	284	LEU	2.4
1	A	82	PHE	2.4
1	A	176	PHE	2.3
1	A	249	VAL	2.3
1	A	362	SER	2.3
1	A	306	LEU	2.3
1	A	239	ILE	2.3
1	A	203	ASP	2.3
1	A	357	LYS	2.3
1	A	108	LEU	2.2
1	A	326	ASN	2.1
1	A	345	ASN	2.1
1	A	344	VAL	2.1
1	A	104	ASN	2.1
1	A	188	THR	2.1
1	A	262	GLY	2.1
1	A	228	ALA	2.1
1	A	231	VAL	2.0
1	A	225	HIS	2.0
1	A	230	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	DGI	A	369	27/27	0.95	0.10	34,39,42,46	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.