



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 7, 2026 – 04:49 AM UTC

PDB ID : 2H35 / pdb_00002h35
Title : Solution structure of Human normal adult hemoglobin
Authors : Fan, J.S.; Yang, D.
Deposited on : 2006-05-22

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<https://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	141	26% 60% 10% .
1	C	141	26% 60% 10% .
2	B	146	47% 50% ..
2	D	146	44% 53% ..

2 Ensemble composition and analysis ⓘ

This entry contains 20 models. Model 5 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:41, A:45-A:138, B:2-B:146, C:1-C:41, C:45- C:138, D:2-D:146 (560)	0.98	5

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 4 single-model clusters were found.

Cluster number	Models
1	5, 6, 7, 10, 11, 12, 18
2	4, 8, 16, 17
3	9, 13, 14
4	3, 15
Single-model clusters	1; 2; 19; 20

3 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 9096 atoms, of which 4540 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called Hemoglobin alpha subunit.

Mol	Chain	Residues	Atoms						Trace
1	A	141	Total	C	H	N	O	S	0
			2152	685	1083	187	194	3	
1	C	141	Total	C	H	N	O	S	0
			2152	685	1083	187	194	3	

- Molecule 2 is a protein called Hemoglobin beta subunit.

Mol	Chain	Residues	Atoms						Trace
2	B	146	Total	C	H	N	O	S	0
			2250	724	1127	195	201	3	
2	D	146	Total	C	H	N	O	S	0
			2250	724	1127	195	201	3	

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).



Mol	Chain	Residues	Atoms					
3	A	1	Total	C	Fe	H	N	O
			73	34	1	30	4	4
3	B	1	Total	C	Fe	H	N	O
			73	34	1	30	4	4

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					
			Total	C	Fe	H	N	O
3	C	1	73	34	1	30	4	4
3	D	1	73	34	1	30	4	4

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Hemoglobin alpha subunit




- Molecule 1: Hemoglobin alpha subunit



- Molecule 2: Hemoglobin beta subunit




- Molecule 2: Hemoglobin beta subunit




4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 5. Colouring as in section 4.1 above.


- Molecule 1: Hemoglobin alpha subunit




- Molecule 1: Hemoglobin alpha subunit



- Molecule 2: Hemoglobin beta subunit



- Molecule 2: Hemoglobin beta subunit



5 Refinement protocol and experimental data overview

The models were refined using the following method: *torsion angle dynamics*.

Of the 20 calculated structures, 20 were deposited, based on the following criterion: *structures with acceptable covalent geometry, structures with the least restraint violations, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
xplor-nih	structure solution	2.14
xplor-nih	refinement	2.14

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.46±0.01	0±0/1031 (0.0± 0.0%)	1.21±0.01	0±0/1405 (0.0± 0.0%)
1	C	1.46±0.01	0±0/1031 (0.0± 0.0%)	1.21±0.01	0±0/1405 (0.0± 0.0%)
2	B	1.45±0.01	0±0/1146 (0.0± 0.0%)	1.22±0.01	0±0/1556 (0.0± 0.0%)
2	D	1.45±0.01	0±0/1146 (0.0± 0.0%)	1.22±0.01	0±0/1556 (0.0± 0.0%)
All	All	1.45	2/87080 (0.0%)	1.21	12/118440 (0.0%)

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	C	3	SER	N-CA	5.56	1.49	1.46	4	1
1	A	3	SER	N-CA	5.44	1.49	1.46	4	1

5 of 8 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	138	SER	CB-CA-C	-5.31	109.47	115.79	19	1
1	C	20	HIS	N-CA-C	-5.29	106.42	112.92	16	3
1	C	138	SER	CB-CA-C	-5.26	109.53	115.79	19	1
1	A	20	HIS	N-CA-C	-5.26	106.45	112.92	16	3
2	B	47	ASP	CB-CA-C	-5.24	109.55	115.79	20	1

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1006	1023	1013	133±19
1	C	1006	1023	1013	133±22
2	B	1116	1116	1107	117±25
2	D	1116	1116	1107	122±30
3	A	43	30	30	10±5
3	B	43	30	30	24±11
3	C	43	30	30	12±11
3	D	43	30	30	24±14
All	All	88320	87960	87181	9349

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 53.

5 of 3504 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:40:ARG:CD	1:C:92:ARG:HD3	1.59	1.25	18	2
2:B:146:HIS:CD2	2:D:139:ASN:HD21	1.58	1.16	6	2
2:B:139:ASN:HD21	2:D:146:HIS:CD2	1.57	1.16	6	2
2:B:106:LEU:HD22	3:B:147:HEM:CBB	1.57	1.16	19	1
2:B:146:HIS:CD2	2:D:2:HIS:NE2	1.57	1.73	15	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	134/141 (95%)	100±3 (74±2%)	22±3 (16±2%)	13±2 (9±2%)	1	10
1	C	134/141 (95%)	100±3 (74±2%)	22±3 (16±2%)	13±2 (9±2%)	1	10

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	144/146 (99%)	111±4 (77±3%)	29±4 (20±3%)	4±2 (3±1%)	5	38
2	D	144/146 (99%)	111±4 (77±3%)	29±4 (20±3%)	4±2 (3±1%)	5	38
All	All	11120/11480 (97%)	8418 (76%)	2024 (18%)	678 (6%)	2	19

5 of 118 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	73	VAL	20
1	C	73	VAL	20
1	A	75	ASP	19
1	A	119	PRO	19
1	C	75	ASP	19

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/113 (95%)	93±3 (87±3%)	14±3 (13±3%)	6	47
1	C	107/113 (95%)	93±3 (87±3%)	14±3 (13±3%)	6	47
2	B	117/118 (99%)	105±2 (90±2%)	12±2 (10±2%)	9	53
2	D	117/118 (99%)	105±2 (90±2%)	12±2 (10±2%)	9	53
All	All	8960/9240 (97%)	7922 (88%)	1038 (12%)	7	50

5 of 229 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	3	SER	20
1	A	127	LYS	20
1	C	3	SER	20
1	C	127	LYS	20
2	B	3	LEU	19

6.3.3 RNA ⓘ

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
3	HEM	A	142	-	50,50,50	1.26±0.00	4±0 (8±0%)
3	HEM	D	147	-	50,50,50	1.25±0.00	4±0 (8±0%)
3	HEM	B	147	-	50,50,50	1.25±0.00	4±0 (8±0%)
3	HEM	C	142	-	50,50,50	1.26±0.01	4±0 (8±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
3	HEM	A	142	-	67,82,82	1.02±0.00	3±0 (4±0%)
3	HEM	D	147	-	67,82,82	1.02±0.00	3±0 (4±0%)

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
3	HEM	B	147	-	67,82,82	1.02±0.00	3±0 (4±0%)
3	HEM	C	142	-	67,82,82	1.02±0.01	3±0 (4±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	B	147	-	-	0±0,14,54,54	-
3	HEM	A	142	-	-	0±0,14,54,54	-
3	HEM	C	142	-	-	0±0,14,54,54	-
3	HEM	D	147	-	-	0±0,14,54,54	-

5 of 16 unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
3	C	142	HEM	CBB-CAB	4.93	1.54	1.30	16	20
3	A	142	HEM	CBB-CAB	4.92	1.54	1.30	1	20
3	B	147	HEM	CBB-CAB	4.90	1.53	1.30	10	20
3	D	147	HEM	CBB-CAB	4.89	1.53	1.30	1	20
3	C	142	HEM	CBC-CAC	4.84	1.53	1.30	18	20

5 of 12 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
3	D	147	HEM	C3B-C4B-NB	3.05	111.66	109.47	7	20
3	B	147	HEM	C3B-C4B-NB	3.00	111.62	109.47	1	20
3	C	142	HEM	C3B-C4B-NB	3.00	111.62	109.47	12	20
3	A	142	HEM	C3B-C4B-NB	2.98	111.61	109.47	7	20
3	D	147	HEM	CBB-CAB-C3B	2.94	112.84	127.53	12	20


There are no chirality outliers.


There are no torsion outliers.


There are no ring outliers.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided