



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 03:00 AM UTC

PDB ID : 4H3C / pdb_00004h3c
Title : Structure of E. coli undecaprenyl diphosphate synthase in complex with BPH-987
Authors : Zhu, W.; Oldfield, E.
Deposited on : 2012-09-13
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1452 (1.94-1.94)
Clashscore	190562	1494 (1.94-1.94)
Ramachandran outliers	187476	1479 (1.94-1.94)
Sidechain outliers	187428	1479 (1.94-1.94)
RSRZ outliers	180081	1453 (1.94-1.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	253	10% 66% 14% • 17%
1	B	253	8% 67% 14% 19%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 3407 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Undecaprenyl pyrophosphate synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	0	0
			1629	1024	301	298	6			
1	B	206	Total	C	N	O	S	0	0	0
			1628	1023	305	295	5			

- Molecule 2 is (2Z)-4-[3-(decyloxy)phenyl]-2-hydroxy-4-oxobut-2-enoic acid (CCD ID: 0YZ) (formula: C₂₀H₂₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			25	20	5		
2	A	1	Total	C	O	0	0
			25	20	5		
2	B	1	Total	C	O	0	0
			25	20	5		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	43	Total 43	O 43	0	0
3	B	32	Total 32	O 32	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Undecaprenyl pyrophosphate synthase



• Molecule 1: Undecaprenyl pyrophosphate synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.33Å 69.10Å 111.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	43.43 – 1.93 43.43 – 1.93	Depositor EDS
% Data completeness (in resolution range)	99.1 (43.43-1.93) 99.0 (43.43-1.93)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.98 (at 1.94Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.206 , 0.250 0.207 , 0.247	Depositor DCC
R_{free} test set	1871 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	27.7	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 34.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3407	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OYZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.52	10/1659 (0.6%)	1.30	13/2242 (0.6%)
1	B	1.54	13/1659 (0.8%)	1.25	6/2240 (0.3%)
All	All	1.53	23/3318 (0.7%)	1.27	19/4482 (0.4%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	20	HIS	ND1-CE1	7.47	1.40	1.32
1	B	45	ALA	CA-C	7.22	1.62	1.52
1	B	199	HIS	CG-ND1	-6.52	1.31	1.38
1	A	172	ILE	N-CA	6.28	1.53	1.46
1	A	20	HIS	CG-CD2	6.25	1.42	1.35
1	A	138	THR	CA-C	6.15	1.60	1.52
1	A	215	TYR	CA-C	6.07	1.59	1.52
1	B	66	THR	CA-C	5.95	1.59	1.52
1	A	199	HIS	CG-ND1	-5.91	1.31	1.38
1	B	103	HIS	CG-ND1	-5.91	1.31	1.38
1	B	207	TRP	NE1-CE2	-5.88	1.30	1.37
1	B	18	CYS	C-O	-5.78	1.16	1.24
1	B	219	VAL	CA-CB	5.65	1.60	1.54
1	B	108	ARG	C-O	-5.57	1.17	1.23
1	B	199	HIS	CD2-NE2	-5.54	1.31	1.37
1	A	236	PHE	CA-C	5.50	1.59	1.52
1	B	232	ALA	N-CA	-5.33	1.39	1.46
1	B	225	ASP	C-O	-5.29	1.17	1.24
1	A	38	ILE	CA-C	-5.24	1.46	1.53
1	B	20	HIS	ND1-CE1	5.16	1.37	1.32
1	B	175	GLU	C-O	-5.12	1.18	1.24
1	A	147	GLY	C-O	5.11	1.29	1.23
1	A	49	SER	CA-C	5.09	1.59	1.52

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	39	ARG	N-CA-C	9.67	123.03	111.33
1	B	234	ASN	N-CA-C	8.14	120.16	111.28
1	B	207	TRP	N-CA-C	-7.94	102.57	111.14
1	B	221	TRP	CA-C-N	-7.00	112.49	119.56
1	B	221	TRP	C-N-CA	-7.00	112.49	119.56
1	A	204	PHE	N-CA-C	-6.22	97.55	110.80
1	A	209	ILE	N-CA-C	6.21	117.44	111.91
1	A	119	ARG	N-CA-C	6.18	117.68	111.07
1	A	90	VAL	N-CA-CB	5.80	117.34	110.55
1	A	196	GLY	N-CA-C	-5.78	107.40	114.92
1	A	197	GLY	N-CA-C	5.77	122.71	115.21
1	A	38	ILE	O-C-N	-5.58	116.79	122.54
1	A	38	ILE	CA-C-O	5.56	128.30	122.13
1	A	94	ASP	N-CA-C	5.48	122.46	110.80
1	B	90	VAL	CA-C-N	-5.39	111.24	121.54
1	B	90	VAL	C-N-CA	-5.39	111.24	121.54
1	A	25	MET	N-CA-C	5.34	117.62	109.62
1	A	208	GLN	CA-C-N	-5.18	115.84	122.26
1	A	208	GLN	C-N-CA	-5.18	115.84	122.26

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1629	0	1572	44	1
1	B	1628	0	1591	13	1
2	A	50	0	53	7	0
2	B	25	0	26	0	0
3	A	43	0	0	0	0
3	B	32	0	0	0	0
All	All	3407	0	3242	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 9.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:93:LEU:CD2	1:A:94:ASP:N	1.73	1.51
1:A:117:ASN:ND2	1:A:119:ARG:N	1.79	1.30
1:A:117:ASN:ND2	1:A:119:ARG:H	1.33	1.20
1:A:93:LEU:C	1:A:93:LEU:HD23	1.74	1.13
1:A:93:LEU:HD23	1:A:94:ASP:N	1.37	1.12
1:A:93:LEU:HD22	1:A:94:ASP:N	1.47	1.09
1:A:93:LEU:HD22	1:A:94:ASP:H	1.03	1.05
1:A:93:LEU:HD23	1:A:94:ASP:CA	1.84	1.05
1:A:93:LEU:CD2	1:A:93:LEU:C	2.29	1.03
1:A:117:ASN:HD21	1:A:119:ARG:CA	1.71	1.03
1:A:93:LEU:CD2	1:A:94:ASP:CA	2.40	0.98
1:A:117:ASN:HD21	1:A:119:ARG:CB	1.77	0.96
1:A:117:ASN:HD22	1:A:117:ASN:C	1.73	0.96
1:A:117:ASN:HD21	1:A:119:ARG:N	1.44	0.96
1:A:117:ASN:ND2	1:A:117:ASN:C	2.29	0.90
1:A:93:LEU:HD23	1:A:94:ASP:HA	1.50	0.90
1:A:117:ASN:HD22	1:A:119:ARG:H	1.17	0.87
1:A:93:LEU:CD2	1:A:94:ASP:HA	2.06	0.85
1:A:117:ASN:HD22	1:A:119:ARG:N	1.66	0.83
1:A:141:ILE:HD12	2:A:301:OYZ:H24	1.61	0.82
1:A:117:ASN:HD22	1:A:118:SER:N	1.80	0.80
1:A:141:ILE:CD1	2:A:301:OYZ:H24	2.12	0.78
1:B:90:VAL:O	1:B:91:TRP:C	2.26	0.72
1:A:93:LEU:HB2	2:A:301:OYZ:H27	1.74	0.69
1:A:117:ASN:ND2	1:A:119:ARG:CB	2.57	0.66
1:A:98:LYS:O	1:A:102:ARG:HG3	2.00	0.60
1:B:57:ALA:HA	1:B:62:ILE:HD12	1.84	0.60
1:B:201:ILE:HD13	1:B:209:ILE:HD11	1.84	0.59
1:B:168:GLN:HB2	1:B:171:GLN:HG3	1.85	0.58
1:B:176:MET:O	1:B:180:HIS:HD2	1.87	0.58
1:A:103:HIS:NE2	2:A:301:OYZ:H3	2.19	0.57
1:B:91:TRP:HA	1:B:91:TRP:CE3	2.41	0.56
1:A:101:HIS:HD2	1:A:134:ASN:HD21	1.54	0.55
1:A:93:LEU:O	1:A:96:GLU:N	2.40	0.53
1:A:163:GLN:HE22	1:B:168:GLN:HE22	1.57	0.53
1:B:44:LYS:HG3	1:B:48:LYS:HZ2	1.75	0.52
1:A:94:ASP:CB	1:A:123:ARG:HH22	2.23	0.50
1:B:238:ASN:O	1:B:239:ARG:C	2.54	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:HIS:CD2	1:A:134:ASN:HD21	2.30	0.50
1:A:117:ASN:ND2	1:A:117:ASN:O	2.42	0.49
1:A:90:VAL:O	1:A:93:LEU:HB3	2.12	0.49
1:A:106:ARG:NH1	1:A:108:ARG:NH2	2.61	0.49
2:A:302:0YZ:OAC	2:A:302:0YZ:OAE	2.30	0.48
1:A:94:ASP:CB	1:A:123:ARG:NH2	2.77	0.48
1:A:103:HIS:NE2	2:A:301:0YZ:CAF	2.77	0.48
1:A:226:GLU:O	1:A:230:GLU:HG3	2.14	0.47
1:B:98:LYS:H	1:B:98:LYS:HG2	1.53	0.46
1:A:47:ALA:HB1	2:A:302:0YZ:H28	1.98	0.46
1:A:142:ALA:HB1	1:A:145:TYR:HB3	1.98	0.45
1:B:142:ALA:HB1	1:B:145:TYR:HB3	1.99	0.45
1:A:117:ASN:O	1:A:118:SER:C	2.60	0.45
1:A:163:GLN:HE22	1:B:168:GLN:NE2	2.15	0.43
1:A:117:ASN:HD22	1:A:118:SER:C	2.25	0.43
1:B:151:ILE:O	1:B:155:VAL:HG23	2.20	0.42
1:A:93:LEU:O	1:A:96:GLU:HB2	2.18	0.42
1:A:198:GLU:HG3	1:A:200:ARG:HG3	2.02	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:179:GLN:NE2	1:B:106:ARG:NH2[3_645]	1.79	0.41

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	206/253 (81%)	200 (97%)	6 (3%)	0	100	100
1	B	202/253 (80%)	195 (96%)	7 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	408/506 (81%)	395 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	161/206 (78%)	155 (96%)	6 (4%)	30	16
1	B	164/206 (80%)	162 (99%)	2 (1%)	63	57
All	All	325/412 (79%)	317 (98%)	8 (2%)	42	29

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90	VAL
1	A	93	LEU
1	A	117	ASN
1	A	126	LYS
1	A	172	ILE
1	A	200	ARG
1	B	118	SER
1	B	126	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	101	HIS
1	A	117	ASN
1	B	43	HIS
1	B	59	ASN
1	B	164	GLN
1	B	166	ASN
1	B	168	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	180	HIS
1	B	199	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	0YZ	B	301	-	25,25,25	2.34	5 (20%)	30,30,30	1.91	7 (23%)
2	0YZ	A	302	-	25,25,25	2.53	5 (20%)	30,30,30	3.50	5 (16%)
2	0YZ	A	301	-	25,25,25	2.35	4 (16%)	30,30,30	2.68	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	0YZ	B	301	-	-	2/23/23/23	0/1/1/1
2	0YZ	A	302	-	-	3/23/23/23	0/1/1/1
2	0YZ	A	301	-	-	3/23/23/23	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	302	0YZ	CAF-CAW	8.51	1.55	1.36
2	B	301	0YZ	CAF-CAW	8.01	1.54	1.36
2	A	301	0YZ	CAY-CAV	-7.66	1.38	1.49
2	A	302	0YZ	CAY-CAV	-7.06	1.38	1.49
2	A	301	0YZ	CAF-CAW	6.33	1.50	1.36
2	B	301	0YZ	CAW-CAU	4.50	1.55	1.48
2	B	301	0YZ	CAY-CAV	-4.22	1.43	1.49
2	B	301	0YZ	CAF-CAV	3.90	1.55	1.43
2	A	302	0YZ	CAW-CAU	3.84	1.54	1.48
2	A	301	0YZ	CAW-CAU	3.47	1.53	1.48
2	A	302	0YZ	CAF-CAV	2.58	1.51	1.43
2	A	301	0YZ	OAD-CAU	-2.42	1.24	1.30
2	A	302	0YZ	OAD-CAU	-2.22	1.24	1.30
2	B	301	0YZ	CAG-CAI	2.08	1.42	1.38

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	302	0YZ	CAW-CAF-CAV	-17.24	110.89	120.64
2	A	301	0YZ	OAE-CAW-CAU	10.03	123.68	114.17
2	B	301	0YZ	OAE-CAW-CAU	6.54	120.37	114.17
2	A	301	0YZ	CAF-CAW-CAU	-5.34	114.71	122.51
2	A	301	0YZ	OAC-CAV-CAY	4.48	126.27	119.20
2	A	302	0YZ	OAE-CAW-CAU	4.44	118.38	114.17
2	B	301	0YZ	OAD-CAU-CAW	4.22	120.48	115.78
2	A	302	0YZ	OAD-CAU-CAW	3.72	119.93	115.78
2	A	301	0YZ	OAD-CAU-CAW	3.65	119.85	115.78
2	A	301	0YZ	CAW-CAF-CAV	-3.63	118.59	120.64
2	A	301	0YZ	CAY-CAV-CAF	-3.27	112.85	119.83
2	A	302	0YZ	CAS-OAT-CAX	3.12	126.05	117.93
2	B	301	0YZ	CAY-CAJ-CAX	2.84	122.49	119.56
2	A	301	0YZ	CAS-OAT-CAX	2.81	125.23	117.93
2	B	301	0YZ	OAC-CAV-CAY	2.64	123.37	119.20
2	B	301	0YZ	CAW-CAF-CAV	-2.31	119.34	120.64
2	A	301	0YZ	CAL-CAM-CAN	-2.25	102.98	114.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	0YZ	CAY-CAV-CAF	-2.08	115.38	119.83
2	B	301	0YZ	CAF-CAW-CAU	-2.05	119.52	122.51
2	A	302	0YZ	OAD-CAU-OAB	-2.02	119.09	123.90

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	302	0YZ	CAN-CAO-CAP-CAQ
2	A	302	0YZ	CAK-CAL-CAM-CAN
2	A	302	0YZ	CAA-CAK-CAL-CAM
2	A	301	0YZ	CAM-CAN-CAO-CAP
2	A	301	0YZ	CAA-CAK-CAL-CAM
2	B	301	0YZ	CAM-CAN-CAO-CAP
2	B	301	0YZ	CAN-CAO-CAP-CAQ
2	A	301	0YZ	CAK-CAL-CAM-CAN

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	302	0YZ	2	0
2	A	301	0YZ	5	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	210/253 (83%)	0.72	25 (11%) 9 10	15, 30, 62, 75	0
1	B	206/253 (81%)	0.59	20 (9%) 13 15	15, 29, 56, 71	0
All	All	416/506 (82%)	0.66	45 (10%) 11 12	15, 29, 60, 75	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	93	LEU	7.3
1	B	89	PHE	7.0
1	B	92	ALA	5.9
1	B	88	LEU	5.7
1	A	90	VAL	5.7
1	B	91	TRP	5.6
1	A	118	SER	5.2
1	B	169	PRO	4.8
1	A	88	LEU	4.3
1	A	94	ASP	4.0
1	A	95	SER	4.0
1	B	93	LEU	4.0
1	A	91	TRP	3.8
1	A	92	ALA	3.7
1	B	172	ILE	3.7
1	A	122	GLU	3.5
1	A	40	ALA	3.4
1	B	90	VAL	3.4
1	A	170	ASP	3.3
1	A	39	ARG	3.2
1	B	159	ALA	3.2
1	B	165	GLY	3.1
1	A	120	LEU	3.0
1	A	89	PHE	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	119	ARG	2.9
1	B	70	PHE	2.8
1	B	36	GLY	2.8
1	B	167	LEU	2.7
1	B	41	PHE	2.6
1	A	86	MET	2.6
1	B	162	VAL	2.5
1	B	94	ASP	2.5
1	B	17	GLY	2.4
1	A	129	ALA	2.4
1	A	125	ARG	2.3
1	A	70	PHE	2.3
1	A	171	GLN	2.3
1	B	95	SER	2.3
1	B	98	LYS	2.3
1	A	172	ILE	2.3
1	A	175	GLU	2.2
1	A	166	ASN	2.2
1	B	38	ILE	2.1
1	A	169	PRO	2.1
1	A	165	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	0YZ	B	301	25/25	0.76	0.19	32,46,83,90	0
2	0YZ	A	302	25/25	0.79	0.16	41,63,93,94	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	0YZ	A	301	25/25	0.82	0.16	40,49,80,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 0YZ B 301:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 0YZ A 302:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 0YZ A 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.