



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 01:08 PM EDT

PDB ID : 6H3D / pdb_00006h3d
Title : Staphylopine dehydrogenase in complex with xNA
Authors : Hajjar, C.; Arnoux, P.
Deposited on : 2018-07-18
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	441	

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3738 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called DUF2338 domain-containing protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	428	3484	2236	569	656	23	0	1	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	434	LEU	-	expression tag	UNP A0A1K7Y513
A	435	GLU	-	expression tag	UNP A0A1K7Y513
A	436	HIS	-	expression tag	UNP A0A1K7Y513
A	437	HIS	-	expression tag	UNP A0A1K7Y513
A	438	HIS	-	expression tag	UNP A0A1K7Y513
A	439	HIS	-	expression tag	UNP A0A1K7Y513
A	440	HIS	-	expression tag	UNP A0A1K7Y513
A	441	HIS	-	expression tag	UNP A0A1K7Y513

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	1
			96	42	14	34	6		

- Molecule 3 is D-HISTIDINE (CCD ID: DHI) (formula: $C_6H_{10}N_3O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			11	6	3	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			13	8	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	128	Total	O	0	0
			128	128		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DUF2338 domain-containing protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	185.09Å 48.75Å 59.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	92.54 – 2.05 92.54 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.2 (92.54-2.05) 99.2 (92.54-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.192 , 0.229 0.202 , 0.234	Depositor DCC
R_{free} test set	1733 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	49.0	Xtriage
Anisotropy	0.067	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 33.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3738	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PG4, NDP, DHI

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.35	13/3562 (0.4%)	1.29	19/4814 (0.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	355	ILE	N-CA	-9.17	1.38	1.46
1	A	328	PRO	C-O	-7.62	1.15	1.23
1	A	39	LYS	CA-C	7.44	1.62	1.52
1	A	380	THR	N-CA	6.36	1.55	1.46
1	A	72	ASN	C-O	-6.22	1.16	1.24
1	A	395	GLN	N-CA	5.66	1.53	1.46
1	A	197	LEU	C-O	-5.54	1.17	1.24
1	A	369	ILE	C-O	-5.44	1.17	1.24
1	A	56	LYS	N-CA	5.28	1.51	1.45
1	A	360	SER	N-CA	5.26	1.52	1.46
1	A	163	LEU	CA-C	-5.12	1.46	1.52
1	A	241	LYS	C-O	5.09	1.30	1.23
1	A	229	ILE	C-O	-5.09	1.18	1.24

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	379	LYS	N-CA-C	10.15	122.10	111.14
1	A	229	ILE	CB-CA-C	-8.20	101.28	112.02
1	A	52	GLN	CA-CB-CG	7.30	128.69	114.10
1	A	413	ASN	N-CA-C	-6.70	98.70	109.96
1	A	31	VAL	CB-CA-C	-6.59	101.53	110.42
1	A	124	SER	N-CA-C	-6.06	104.24	111.69
1	A	380	THR	N-CA-C	5.77	122.57	109.81
1	A	413	ASN	CB-CA-C	5.68	118.90	109.53
1	A	299	GLU	N-CA-C	5.58	117.36	111.28
1	A	422	VAL	CB-CA-C	-5.56	104.74	112.02
1	A	69	PHE	N-CA-C	5.54	117.27	108.79
1	A	306	ASP	N-CA-C	5.53	117.75	111.11
1	A	133	SER	N-CA-C	5.52	118.06	111.71
1	A	340	VAL	CA-C-N	-5.42	114.67	120.03
1	A	340	VAL	C-N-CA	-5.42	114.67	120.03
1	A	263	MET	N-CA-C	5.26	117.70	111.33
1	A	43	LEU	CA-CB-CG	-5.14	98.32	116.30
1	A	43	LEU	N-CA-C	-5.07	105.83	111.36
1	A	28	ILE	CB-CA-C	-5.01	103.64	110.96

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	412	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3484	0	3481	46	0
2	A	96	0	52	3	0
3	A	11	0	9	0	0
4	A	6	0	8	0	0
5	A	13	0	18	0	0
6	A	128	0	0	14	0
All	All	3738	0	3568	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:256:MET:SD	6:A:823:HOH:O	2.14	1.05
1:A:92:CYS:SG	6:A:778:HOH:O	2.20	0.98
1:A:9:THR:HB	1:A:43:LEU:HD22	1.59	0.83
1:A:60:GLU:OE1	1:A:63:GLN:NE2	2.17	0.77
1:A:404:GLN:HG2	6:A:804:HOH:O	1.89	0.72
1:A:7:ILE:HG13	1:A:90:MET:HE3	1.72	0.70
1:A:64:HIS:HB3	6:A:800:HOH:O	1.94	0.67
1:A:249:THR:HG22	1:A:252:LEU:H	1.60	0.66
1:A:256:MET:HE1	1:A:281:MET:HE1	1.80	0.62
1:A:349:GLU:H	1:A:349:GLU:CD	2.07	0.61
1:A:217:PRO:HB3	1:A:221:MET:CE	2.32	0.59
1:A:70:GLU:HG2	6:A:754:HOH:O	2.05	0.56
1:A:225:SER:O	1:A:229:ILE:HG12	2.06	0.55
1:A:412:THR:HA	6:A:734:HOH:O	2.07	0.55
1:A:256:MET:HE3	1:A:277:LEU:HD21	1.90	0.54
1:A:7:ILE:HG21	1:A:100:THR:HG21	1.90	0.52
1:A:217:PRO:HB3	1:A:221:MET:HE2	1.90	0.52
1:A:77:ASP:HB2	6:A:702:HOH:O	2.10	0.51
1:A:366:THR:HG23	1:A:383:ILE:HG23	1.91	0.51
1:A:173:MET:HE2	1:A:186:ILE:HG21	1.93	0.50
1:A:185:ARG:NH1	6:A:708:HOH:O	2.46	0.48
1:A:173:MET:HB2	1:A:197:LEU:HD11	1.96	0.48
1:A:221:MET:HE3	1:A:386:PHE:HB3	1.95	0.48
1:A:52:GLN:HE21	1:A:68:LYS:HB3	1.79	0.47
1:A:211:SER:CB	1:A:285:ASN:HD21	2.27	0.47
1:A:321:LEU:HA	1:A:338:SER:OG	2.14	0.47
1:A:286:TYR:OH	1:A:323:ASP:OD1	2.26	0.47
1:A:173:MET:HE2	1:A:186:ILE:CG2	2.45	0.47
1:A:19:ILE:HD11	1:A:65:LEU:HD23	1.97	0.46
1:A:191:GLU:HG3	6:A:795:HOH:O	2.16	0.45
1:A:90:MET:HE2	6:A:778:HOH:O	2.16	0.45
1:A:150:ASP:HA	2:A:600[B]:NDP:N7N	2.33	0.44
1:A:256:MET:HE2	1:A:313:LEU:HD22	2.00	0.44
1:A:256:MET:CE	1:A:313:LEU:HD22	2.48	0.44
1:A:259:MET:HE3	1:A:383:ILE:HD11	2.01	0.43
1:A:43:LEU:O	1:A:43:LEU:HG	2.14	0.43
1:A:249:THR:HG21	6:A:820:HOH:O	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:7:ILE:HD13	1:A:31:VAL:HG21	2.01	0.42
1:A:256:MET:HE2	1:A:313:LEU:HD13	2.01	0.42
1:A:343:LYS:HG2	6:A:793:HOH:O	2.19	0.42
1:A:329:ASP:OD1	1:A:329:ASP:C	2.63	0.42
1:A:365:ARG:NE	6:A:719:HOH:O	2.53	0.41
2:A:600[B]:NDP:H6N	6:A:750:HOH:O	2.20	0.41
1:A:150:ASP:HA	2:A:600[A]:NDP:N7N	2.35	0.41
1:A:6:MET:HE2	1:A:6:MET:HB2	1.98	0.41
1:A:258:LEU:HD23	1:A:258:LEU:HA	1.95	0.40
1:A:321:LEU:O	1:A:338:SER:HB2	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	427/441 (97%)	409 (96%)	17 (4%)	1 (0%)	43	37

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	64	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	392/404 (97%)	369 (94%)	23 (6%)	18	10

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	ILE
1	A	36	THR
1	A	43	LEU
1	A	64	HIS
1	A	70	GLU
1	A	82	LYS
1	A	90	MET
1	A	125	GLN
1	A	136	SER
1	A	152	ARG
1	A	156	LYS
1	A	157	GLU
1	A	173	MET
1	A	194	LYS
1	A	213	LEU
1	A	242	LEU
1	A	249	THR
1	A	259	MET
1	A	299	GLU
1	A	338	SER
1	A	349	GLU
1	A	357	ARG
1	A	385	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	15	GLN
1	A	52	GLN
1	A	58	GLN
1	A	72	ASN
1	A	125	GLN
1	A	177	HIS
1	A	179	ASN
1	A	184	GLN
1	A	192	GLN
1	A	205	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	285	ASN
1	A	350	GLN
1	A	395	GLN
1	A	404	GLN
1	A	405	GLN
1	A	429	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NDP	A	600[B]	-	51,52,52	1.42	6 (11%)	71,80,80	1.78	17 (23%)
2	NDP	A	600[A]	-	51,52,52	1.50	7 (13%)	71,80,80	1.87	16 (22%)
3	DHI	A	601	-	10,11,11	1.10	1 (10%)	10,14,14	1.55	2 (20%)
4	GOL	A	602	-	5,5,5	0.55	0	5,5,5	1.43	1 (20%)
5	PG4	A	603	-	12,12,12	0.56	0	11,11,11	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	A	600[B]	-	-	13/34/77/77	0/5/5/5
2	NDP	A	600[A]	-	-	13/34/77/77	0/5/5/5
3	DHI	A	601	-	-	4/8/8/8	0/1/1/1
4	GOL	A	602	-	-	2/4/4/4	-
5	PG4	A	603	-	-	6/10/10/10	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	600[A]	NDP	C5A-C4A	4.94	1.47	1.39
2	A	600[B]	NDP	C5A-C4A	4.83	1.47	1.39
2	A	600[B]	NDP	PN-O3	4.34	1.64	1.59
2	A	600[A]	NDP	PN-O3	4.28	1.64	1.59
2	A	600[A]	NDP	C5A-C6A	3.21	1.49	1.41
2	A	600[A]	NDP	C8A-N7A	3.18	1.37	1.31
2	A	600[B]	NDP	C8A-N7A	2.74	1.36	1.31
2	A	600[A]	NDP	PA-O3	2.37	1.62	1.59
2	A	600[B]	NDP	PA-O3	2.32	1.62	1.59
2	A	600[B]	NDP	C5A-C6A	2.31	1.47	1.41
3	A	601	DHI	OXT-C	-2.27	1.23	1.30
2	A	600[A]	NDP	C6N-C5N	2.22	1.39	1.33
2	A	600[A]	NDP	P2B-O2B	2.10	1.63	1.59
2	A	600[B]	NDP	C6N-C5N	2.08	1.39	1.33

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600[A]	NDP	C5A-C4A-N3A	-5.64	118.95	126.72
2	A	600[B]	NDP	C5A-C4A-N3A	-5.63	118.96	126.72
2	A	600[B]	NDP	N3A-C4A-N9A	4.84	135.40	127.17
2	A	600[A]	NDP	O4B-C1B-N9A	4.76	117.23	108.09
2	A	600[A]	NDP	C2A-N3A-C4A	4.30	122.33	111.83
2	A	600[A]	NDP	N3A-C2A-N1A	-4.22	122.20	128.58
2	A	600[A]	NDP	N3A-C4A-N9A	4.07	134.10	127.17
2	A	600[B]	NDP	O4B-C1B-N9A	4.05	115.87	108.09
2	A	600[B]	NDP	N3A-C2A-N1A	-3.97	122.57	128.58
2	A	600[B]	NDP	C2A-N3A-C4A	3.92	121.40	111.83
2	A	600[A]	NDP	C4A-C5A-N7A	-3.88	106.14	110.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	600[A]	NDP	O4D-C1D-N1N	3.83	115.39	108.08
2	A	600[B]	NDP	O4D-C1D-N1N	3.82	115.37	108.08
2	A	600[A]	NDP	C6A-C5A-N7A	3.31	138.48	132.09
2	A	600[A]	NDP	C5A-N7A-C8A	3.06	108.25	103.45
2	A	600[B]	NDP	C6N-N1N-C2N	2.98	122.50	119.32
4	A	602	GOL	O1-C1-C2	-2.96	97.04	110.38
3	A	601	DHI	CA-CB-CG	2.80	120.67	113.77
2	A	600[B]	NDP	C4A-N9A-C8A	2.76	108.63	105.74
2	A	600[A]	NDP	N9A-C8A-N7A	-2.66	110.16	113.94
2	A	600[A]	NDP	C4A-N9A-C8A	2.62	108.49	105.74
2	A	600[A]	NDP	C6N-N1N-C2N	2.50	121.99	119.32
3	A	601	DHI	OXT-C-O	-2.50	118.41	124.08
2	A	600[B]	NDP	C4A-C5A-N7A	-2.46	107.77	110.58
2	A	600[B]	NDP	C3N-C2N-N1N	-2.41	119.66	123.20
2	A	600[B]	NDP	C5A-N7A-C8A	2.26	107.00	103.45
2	A	600[A]	NDP	C4A-N9A-C1B	-2.23	121.42	126.63
2	A	600[A]	NDP	C5B-C4B-C3B	-2.21	107.25	115.21
2	A	600[B]	NDP	C2A-N1A-C6A	2.21	122.36	118.73
2	A	600[B]	NDP	C5B-C4B-C3B	-2.21	107.27	115.21
2	A	600[B]	NDP	C1D-N1N-C6N	-2.13	116.26	120.77
2	A	600[B]	NDP	N9A-C8A-N7A	-2.10	110.95	113.94
2	A	600[B]	NDP	C2B-C1B-N9A	-2.03	110.41	113.75
2	A	600[A]	NDP	C2A-N1A-C6A	2.01	122.03	118.73
2	A	600[B]	NDP	O7N-C7N-C3N	2.00	124.67	120.90
2	A	600[A]	NDP	C3N-C2N-N1N	-2.00	120.27	123.20

There are no chirality outliers.

All (38) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	600[A]	NDP	C5B-O5B-PA-O2A
2	A	600[A]	NDP	C5B-O5B-PA-O3
2	A	600[A]	NDP	C5D-O5D-PN-O3
2	A	600[A]	NDP	C5D-O5D-PN-O1N
2	A	600[B]	NDP	C5B-O5B-PA-O2A
2	A	600[B]	NDP	C5B-O5B-PA-O3
2	A	600[B]	NDP	C5D-O5D-PN-O3
2	A	600[B]	NDP	C5D-O5D-PN-O1N
4	A	602	GOL	O1-C1-C2-C3
4	A	602	GOL	O1-C1-C2-O2
5	A	603	PG4	O1-C1-C2-O2
5	A	603	PG4	O3-C5-C6-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	600[A]	NDP	C1B-C2B-O2B-P2B
2	A	600[A]	NDP	C3B-C2B-O2B-P2B
5	A	603	PG4	O4-C7-C8-O5
2	A	600[A]	NDP	PN-O3-PA-O1A
2	A	600[B]	NDP	PN-O3-PA-O1A
5	A	603	PG4	C3-C4-O3-C5
3	A	601	DHI	O-C-CA-CB
2	A	600[A]	NDP	C5D-O5D-PN-O2N
2	A	600[B]	NDP	C5D-O5D-PN-O2N
5	A	603	PG4	C4-C3-O2-C2
3	A	601	DHI	OXT-C-CA-CB
2	A	600[B]	NDP	C3B-C2B-O2B-P2B
2	A	600[B]	NDP	C4B-C5B-O5B-PA
2	A	600[A]	NDP	C2B-O2B-P2B-O2X
5	A	603	PG4	O2-C3-C4-O3
2	A	600[B]	NDP	O4D-C1D-N1N-C6N
2	A	600[A]	NDP	O4D-C1D-N1N-C6N
2	A	600[A]	NDP	C4B-C5B-O5B-PA
2	A	600[B]	NDP	C1B-C2B-O2B-P2B
3	A	601	DHI	CA-CB-CG-ND1
2	A	600[A]	NDP	PN-O3-PA-O5B
2	A	600[B]	NDP	PN-O3-PA-O5B
3	A	601	DHI	CA-CB-CG-CD2
2	A	600[B]	NDP	O4B-C1B-N9A-C8A
2	A	600[A]	NDP	O4B-C4B-C5B-O5B
2	A	600[B]	NDP	O4B-C4B-C5B-O5B


There are no ring outliers.


2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	600[B]	NDP	2	0
2	A	600[A]	NDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the

average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	428/441 (97%)	0.17	4 (0%) 81 83	24, 52, 80, 100	1 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	34	ALA	2.7
1	A	428	ILE	2.2
1	A	64	HIS	2.0
1	A	59	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NDP	A	600[A]	48/48	0.75	0.18	56,88,114,124	48
2	NDP	A	600[B]	48/48	0.75	0.18	58,84,97,108	48
3	DHI	A	601	11/11	0.89	0.15	52,65,101,105	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PG4	A	603	13/13	0.91	0.17	58,71,95,96	0
4	GOL	A	602	6/6	0.94	0.09	45,51,52,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.