



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 10:47 AM UTC

PDB ID : 3H4V / pdb_00003h4v
Title : Selective screening and design to identify inhibitors of leishmania major pteridine reductase 1
Authors : Mcluskey, K.; Gibellini, F.; Hunter, W.N.
Deposited on : 2009-04-21
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4912 (2.40-2.40)
Clashscore	190562	5391 (2.40-2.40)
Ramachandran outliers	187476	5320 (2.40-2.40)
Sidechain outliers	187428	5321 (2.40-2.40)
RSRZ outliers	180081	4916 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	288	 73% 17% 5% 5%
1	B	288	 70% 21% 5% 5%
1	C	288	 73% 15% 10% 2%
1	D	288	 73% 16% 9% 2%
1	E	288	 63% 27% 8% 2%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	288	
1	G	288	
1	H	288	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	DVP	G	301	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 17638 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pteridine reductase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	273	Total	C	N	O	S	0	1	0
			2034	1278	362	383	11			
1	B	273	Total	C	N	O	S	0	1	0
			2028	1276	360	381	11			
1	C	259	Total	C	N	O	S	0	2	0
			1941	1220	348	363	10			
1	D	263	Total	C	N	O	S	0	2	0
			1973	1245	350	367	11			
1	E	266	Total	C	N	O	S	0	4	0
			1999	1259	353	376	11			
1	F	270	Total	C	N	O	S	0	1	0
			2018	1268	359	380	11			
1	G	261	Total	C	N	O	S	0	0	0
			1948	1223	348	366	11			
1	H	263	Total	C	N	O	S	0	0	0
			1959	1228	350	371	10			

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	162	VAL	PHE	SEE REMARK 999	UNP Q01782
B	162	VAL	PHE	SEE REMARK 999	UNP Q01782
C	162	VAL	PHE	SEE REMARK 999	UNP Q01782
D	162	VAL	PHE	SEE REMARK 999	UNP Q01782
E	162	VAL	PHE	SEE REMARK 999	UNP Q01782
F	162	VAL	PHE	SEE REMARK 999	UNP Q01782
G	162	VAL	PHE	SEE REMARK 999	UNP Q01782
H	162	VAL	PHE	SEE REMARK 999	UNP Q01782

- Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	E	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	F	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	G	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
2	H	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is METHYL 1-(4-{[(2,4-DIAMINOPTERIDIN-6-YL)METHYL]AMINO}BENZ OYL)PIPERIDINE-4-CARBOXYLATE (CCD ID: DVP) (formula: C₂₁H₂₄N₈O₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			32	21	8	3		
3	B	1	Total	C	N	O	0	0
			32	21	8	3		
3	C	1	Total	C	N	O	0	0
			32	21	8	3		
3	D	1	Total	C	N	O	0	0
			32	21	8	3		
3	E	1	Total	C	N	O	0	0
			32	21	8	3		
3	F	1	Total	C	N	O	0	0
			32	21	8	3		
3	G	1	Total	C	N	O	0	0
			32	21	8	3		
3	H	1	Total	C	N	O	0	0
			32	21	8	3		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	221	Total	O	0	0
			221	221		
4	B	182	Total	O	0	0
			182	182		
4	C	161	Total	O	0	0
			161	161		
4	D	136	Total	O	0	0
			136	136		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	E	115	Total 115	O 115	0	0
4	F	110	Total 110	O 110	0	0
4	G	95	Total 95	O 95	0	0
4	H	78	Total 78	O 78	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Pteridine reductase 1



• Molecule 1: Pteridine reductase 1




• Molecule 1: Pteridine reductase 1




• Molecule 1: Pteridine reductase 1






• Molecule 1: Pteridine reductase 1

Chain E: 63% 27% 8%




• Molecule 1: Pteridine reductase 1

Chain F: 63% 28% 6%



• Molecule 1: Pteridine reductase 1



Chain G: 30% 48% 11% 9%





● Molecule 1: Pteridine reductase 1

Chain H: 29% 51% 9% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	95.21Å 103.49Å 146.48Å 90.00° 108.24° 90.00°	Depositor
Resolution (Å)	30.00 – 2.40 30.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	95.2 (30.00-2.40) 95.2 (30.00-2.40)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.54 (at 2.39Å)	Xtriage
Refinement program	REFMAC 5.2.0003	Depositor
R, R_{free}	0.276 , 0.356 0.277 , 0.354	Depositor DCC
R_{free} test set	4986 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	17.6	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 31.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.036 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	17638	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 74.56 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4920e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, DVP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.21	2/2077 (0.1%)	1.16	1/2829 (0.0%)
1	B	1.20	1/2071 (0.0%)	1.20	7/2821 (0.2%)
1	C	1.07	1/1982 (0.1%)	1.11	5/2695 (0.2%)
1	D	1.06	2/2017 (0.1%)	1.08	2/2745 (0.1%)
1	E	1.23	3/2054 (0.1%)	1.20	4/2797 (0.1%)
1	F	1.24	3/2061 (0.1%)	1.24	8/2807 (0.3%)
1	G	1.19	5/1982 (0.3%)	1.35	14/2694 (0.5%)
1	H	1.18	3/1993 (0.2%)	1.41	17/2710 (0.6%)
All	All	1.17	20/16237 (0.1%)	1.22	58/22098 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
1	C	0	1
1	D	0	3
1	H	0	2
All	All	0	9

The worst 5 of 20 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	148	ALA	CA-C	7.65	1.59	1.52
1	A	193	ILE	CA-CB	6.80	1.62	1.54
1	D	73	PRO	CA-C	6.56	1.61	1.52
1	D	244	LYS	N-CA	6.53	1.53	1.46
1	E	192	THR	CA-CB	-6.24	1.43	1.53

The worst 5 of 58 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	225	GLY	N-CA-C	-8.26	102.70	110.21
1	B	73	PRO	N-CA-C	7.98	128.91	112.47
1	H	133	ARG	N-CA-C	-7.73	102.18	111.69
1	B	72	ALA	CA-C-N	7.52	129.24	119.84
1	B	72	ALA	C-N-CA	7.52	129.24	119.84

There are no chirality outliers.

5 of 9 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	131	GLY	Peptide
1	A	71	THR	Peptide
1	B	77	ALA	Peptide
1	C	132	ASP	Peptide
1	D	72	ALA	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2034	0	2033	71	0
1	B	2028	0	2031	51	1
1	C	1941	0	1949	39	0
1	D	1973	0	1987	68	0
1	E	1999	0	1998	79	1
1	F	2018	0	2016	70	0
1	G	1948	0	1954	212	0
1	H	1959	0	1959	202	0
2	A	48	0	25	0	0
2	B	48	0	25	1	0
2	C	48	0	25	0	0
2	D	48	0	25	1	0
2	E	48	0	25	2	0
2	F	48	0	25	0	0
2	G	48	0	25	17	0
2	H	48	0	25	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	32	0	24	6	0
3	B	32	0	24	2	0
3	C	32	0	24	3	0
3	D	32	0	24	4	0
3	E	32	0	24	3	0
3	F	32	0	24	5	0
3	G	32	0	24	10	0
3	H	32	0	24	3	0
4	A	221	0	0	30	2
4	B	182	0	0	14	6
4	C	161	0	0	4	1
4	D	136	0	0	10	3
4	E	115	0	0	16	3
4	F	110	0	0	18	1
4	G	95	0	0	31	0
4	H	78	0	0	22	0
All	All	17638	0	16319	745	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 23.

The worst 5 of 745 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:168:LYS:CE	1:D:169:HIS:HD2	1.42	1.31
1:H:120:ARG:HG2	1:H:121:ASN:OD1	1.30	1.25
1:D:168:LYS:HD3	1:D:169:HIS:CD2	1.75	1.22
1:A:20:ARG:NH1	4:A:652:HOH:O	1.74	1.21
1:D:73:PRO:HB3	4:D:1001:HOH:O	1.43	1.19

The worst 5 of 9 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:361:HOH:O	4:E:672:HOH:O[2_646]	1.39	0.81
4:A:374:HOH:O	4:C:451:HOH:O[2_545]	1.55	0.65
1:B:46:ALA:CB	4:E:678:HOH:O[2_646]	1.67	0.53
4:A:630:HOH:O	4:F:289:HOH:O[2_546]	1.77	0.43
4:B:1056:HOH:O	4:D:960:HOH:O[2_646]	1.86	0.34

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/288 (93%)	241 (90%)	21 (8%)	6 (2%)	5	6
1	B	268/288 (93%)	245 (91%)	19 (7%)	4 (2%)	8	12
1	C	253/288 (88%)	231 (91%)	16 (6%)	6 (2%)	4	5
1	D	257/288 (89%)	237 (92%)	18 (7%)	2 (1%)	16	25
1	E	264/288 (92%)	244 (92%)	19 (7%)	1 (0%)	30	43
1	F	265/288 (92%)	244 (92%)	18 (7%)	3 (1%)	11	18
1	G	253/288 (88%)	204 (81%)	39 (15%)	10 (4%)	2	2
1	H	255/288 (88%)	216 (85%)	31 (12%)	8 (3%)	3	3
All	All	2083/2304 (90%)	1862 (89%)	181 (9%)	40 (2%)	6	8

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	72	ALA
1	C	120	ARG
1	G	167	ALA
1	G	231	ASP
1	H	121	ASN

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/223 (95%)	200 (94%)	12 (6%)	18	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	211/223 (95%)	201 (95%)	10 (5%)	23	41
1	C	203/223 (91%)	194 (96%)	9 (4%)	25	43
1	D	207/223 (93%)	200 (97%)	7 (3%)	32	54
1	E	210/223 (94%)	202 (96%)	8 (4%)	29	49
1	F	211/223 (95%)	201 (95%)	10 (5%)	23	41
1	G	203/223 (91%)	176 (87%)	27 (13%)	4	5
1	H	204/223 (92%)	177 (87%)	27 (13%)	4	5
All	All	1661/1784 (93%)	1551 (93%)	110 (7%)	15	26

5 of 110 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	26	LEU
1	G	183	MET
1	H	278	LYS
1	H	215	LEU
1	G	40	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 22 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	216	GLN
1	G	68	ASN
1	G	38	HIS
1	G	216	GLN
1	C	216	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAP	E	300	-	50,52,52	1.82	7 (14%)	71,80,80	1.95	17 (23%)
2	NAP	G	300	-	50,52,52	1.97	7 (14%)	71,80,80	1.76	12 (16%)
3	DVP	E	301	-	35,35,35	1.20	4 (11%)	47,49,49	2.02	10 (21%)
3	DVP	G	301	-	35,35,35	1.46	4 (11%)	47,49,49	2.11	12 (25%)
2	NAP	A	300	-	50,52,52	1.71	3 (6%)	71,80,80	1.74	13 (18%)
3	DVP	H	301	-	35,35,35	1.48	3 (8%)	47,49,49	1.96	13 (27%)
3	DVP	B	301	-	35,35,35	1.32	3 (8%)	47,49,49	2.22	15 (31%)
3	DVP	C	301	-	35,35,35	1.39	4 (11%)	47,49,49	2.16	14 (29%)
3	DVP	F	301	-	35,35,35	1.23	3 (8%)	47,49,49	2.08	8 (17%)
2	NAP	F	300	-	50,52,52	1.75	7 (14%)	71,80,80	2.00	15 (21%)
2	NAP	H	300	-	50,52,52	2.19	7 (14%)	71,80,80	2.08	22 (30%)
2	NAP	D	300	-	50,52,52	1.75	8 (16%)	71,80,80	1.85	14 (19%)
2	NAP	C	300	-	50,52,52	1.60	4 (8%)	71,80,80	1.80	14 (19%)
3	DVP	A	301	-	35,35,35	1.28	3 (8%)	47,49,49	1.98	8 (17%)
3	DVP	D	301	-	35,35,35	1.33	5 (14%)	47,49,49	2.04	8 (17%)
2	NAP	B	300	-	50,52,52	1.72	6 (12%)	71,80,80	1.69	16 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	E	300	-	-	5/35/67/67	0/5/5/5
2	NAP	G	300	-	-	4/35/67/67	0/5/5/5
3	DVP	E	301	-	-	2/19/29/29	0/4/4/4
3	DVP	G	301	-	-	6/19/29/29	0/4/4/4
2	NAP	A	300	-	-	5/35/67/67	0/5/5/5
3	DVP	H	301	-	-	5/19/29/29	0/4/4/4
3	DVP	B	301	-	-	2/19/29/29	0/4/4/4
3	DVP	C	301	-	-	2/19/29/29	0/4/4/4
3	DVP	F	301	-	-	2/19/29/29	0/4/4/4
2	NAP	F	300	-	-	5/35/67/67	0/5/5/5
2	NAP	H	300	-	-	7/35/67/67	0/5/5/5
2	NAP	D	300	-	-	5/35/67/67	0/5/5/5
2	NAP	C	300	-	-	4/35/67/67	0/5/5/5
3	DVP	A	301	-	-	2/19/29/29	0/4/4/4
3	DVP	D	301	-	-	2/19/29/29	0/4/4/4
2	NAP	B	300	-	-	5/35/67/67	0/5/5/5

The worst 5 of 78 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	G	300	NAP	O7N-C7N	9.78	1.42	1.24
2	A	300	NAP	O7N-C7N	9.39	1.41	1.24
2	H	300	NAP	O7N-C7N	9.04	1.41	1.24
2	B	300	NAP	O7N-C7N	8.78	1.40	1.24
2	D	300	NAP	O7N-C7N	8.18	1.39	1.24

The worst 5 of 211 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	300	NAP	N3A-C2A-N1A	-7.34	117.47	128.58
2	F	300	NAP	N3A-C2A-N1A	-6.66	118.50	128.58
3	A	301	DVP	OAX-CAW-CAV	6.44	121.39	111.76
3	D	301	DVP	OAX-CAW-CAV	6.30	121.19	111.76
2	D	300	NAP	N9A-C8A-N7A	-6.19	105.16	113.94

There are no chirality outliers.

5 of 63 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	300	NAP	C5B-O5B-PA-O2A
2	B	300	NAP	C5B-O5B-PA-O1A
2	B	300	NAP	C5B-O5B-PA-O2A
2	C	300	NAP	C5B-O5B-PA-O2A
2	C	300	NAP	C5B-O5B-PA-O3



There are no ring outliers.



13 monomers are involved in 64 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	300	NAP	2	0
2	G	300	NAP	17	0
3	E	301	DVP	3	0
3	G	301	DVP	10	0
3	H	301	DVP	3	0
3	B	301	DVP	2	0
3	C	301	DVP	3	0
3	F	301	DVP	5	0
2	H	300	NAP	11	0
2	D	300	NAP	1	0
3	A	301	DVP	6	0
3	D	301	DVP	4	0
2	B	300	NAP	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	273/288 (94%)	-1.48	0 100 100	5, 16, 33, 62	1 (0%)
1	B	273/288 (94%)	-1.46	0 100 100	7, 16, 32, 60	1 (0%)
1	C	259/288 (89%)	-1.53	0 100 100	7, 15, 41, 69	2 (0%)
1	D	263/288 (91%)	-1.54	0 100 100	7, 16, 45, 90	2 (0%)
1	E	266/288 (92%)	-1.37	0 100 100	7, 16, 39, 63	4 (1%)
1	F	270/288 (93%)	-1.36	0 100 100	7, 17, 39, 81	1 (0%)
1	G	261/288 (90%)	-0.97	0 100 100	12, 38, 63, 93	0
1	H	263/288 (91%)	-1.02	0 100 100	9, 37, 53, 71	0
All	All	2128/2304 (92%)	-1.34	0 100 100	5, 18, 50, 93	11 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	DVP	H	301	32/32	0.96	0.11	86,113,128,129	0
3	DVP	G	301	32/32	0.97	0.08	70,91,106,106	0
2	NAP	G	300	48/48	0.98	0.04	18,37,49,50	0
3	DVP	B	301	32/32	0.99	0.05	18,48,62,63	0
3	DVP	C	301	32/32	0.99	0.05	18,48,62,63	0
3	DVP	D	301	32/32	0.99	0.04	18,48,62,63	0
3	DVP	E	301	32/32	0.99	0.06	18,48,62,63	0
3	DVP	F	301	32/32	0.99	0.05	18,48,62,63	0
2	NAP	H	300	48/48	0.99	0.04	13,41,52,60	0
3	DVP	A	301	32/32	0.99	0.04	18,48,62,63	0
2	NAP	E	300	48/48	1.00	0.03	4,12,16,17	0
2	NAP	F	300	48/48	1.00	0.03	5,12,16,17	0
2	NAP	A	300	48/48	1.00	0.02	4,12,16,17	0
2	NAP	B	300	48/48	1.00	0.02	3,12,16,17	0
2	NAP	C	300	48/48	1.00	0.02	4,12,16,17	0
2	NAP	D	300	48/48	1.00	0.03	4,12,16,17	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around DVP G 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP G 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DVP B 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DVP C 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DVP D 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around DVP E 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DVP F 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP H 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around DVP A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP E 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP F 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP A 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NAP B 300:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP C 300:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.