



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:20 PM UTC

PDB ID : 6H4U / pdb_00006h4u
Title : Crystal structure of human KDM4A in complex with compound 34b
Authors : Le Bihan, Y.V.; van Montfort, R.L.M.
Deposited on : 2018-07-23
Resolution : 2.21 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.21 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7682 (2.24-2.20)
Clashscore	190562	8402 (2.24-2.20)
Ramachandran outliers	187476	8303 (2.24-2.20)
Sidechain outliers	187428	8304 (2.24-2.20)
RSRZ outliers	180081	7683 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	
1	B	360	
1	C	360	
1	D	360	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	GOL	D	404	-	-	X	-

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 11815 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific demethylase 4A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	2	0
			2756	1789	455	497	15			
1	B	348	Total	C	N	O	S	0	1	0
			2736	1782	438	501	15			
1	C	346	Total	C	N	O	S	0	5	1
			2741	1783	448	494	16			
1	D	345	Total	C	N	O	S	0	2	0
			2730	1776	443	495	16			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP O75164
B	0	SER	-	expression tag	UNP O75164
C	0	SER	-	expression tag	UNP O75164
D	0	SER	-	expression tag	UNP O75164

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 8-[4-(1-methylpiperidin-4-yl)pyrazol-1-yl]-3 {H}-pyrido[3,4-d]pyrimidin-4-one (CCD ID: FO2) (formula: C₁₆H₁₈N₆O).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			23	16	6	1		
3	B	1	Total	C	N	O	0	0
			23	16	6	1		
3	C	1	Total	C	N	O	0	0
			22	15	6	1		
3	D	1	Total	C	N	O	0	0
			19	13	5	1		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O S 4 2 1 1	0	0
4	B	1	Total C O S 4 2 1 1	0	0
4	B	1	Total C O S 4 2 1 1	0	0
4	C	1	Total C O S 4 2 1 1	0	0
4	C	1	Total C O S 4 2 1 1	0	0

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	D	1	Total C O 6 3 3	0	0

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total Cl 1 1	0	0
6	B	1	Total Cl 1 1	0	0
6	C	1	Total Cl 1 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	D	1	Total	Cl	0	0
			1	1		

- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	237	Total	O	0	2
			239	239		
7	B	166	Total	O	0	3
			169	169		
7	C	167	Total	O	0	1
			168	168		
7	D	141	Total	O	0	4
			145	145		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Lysine-specific demethylase 4A

Chain A: 





- Molecule 1: Lysine-specific demethylase 4A

Chain B: 





- Molecule 1: Lysine-specific demethylase 4A

Chain C: 



- Molecule 1: Lysine-specific demethylase 4A

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.97Å 104.22Å 145.03Å 90.00° 99.57° 90.00°	Depositor
Resolution (Å)	49.30 – 2.21 49.30 – 2.21	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.30-2.21) 100.0 (49.30-2.21)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.39 (at 2.20Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.176 , 0.208 0.181 , 0.212	Depositor DCC
R_{free} test set	4176 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	49.3	Xtriage
Anisotropy	0.116	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 61.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11815	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, DMS, GOL, CL, FO2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.95	0/2841	1.26	3/3857 (0.1%)
1	B	0.91	0/2823	1.26	4/3846 (0.1%)
1	C	0.93	1/2826 (0.0%)	1.25	1/3843 (0.0%)
1	D	0.90	0/2817	1.26	3/3835 (0.1%)
All	All	0.92	1/11307 (0.0%)	1.26	11/15381 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	15	MET	SD-CE	-5.01	1.67	1.79

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	184	SER	N-CA-C	6.66	119.29	109.24
1	C	184	SER	N-CA-C	6.26	118.69	109.24
1	D	284	ASN	CA-CB-CG	6.14	118.74	112.60
1	B	184	SER	N-CA-C	6.13	119.03	109.52
1	A	184	SER	N-CA-C	6.03	118.87	109.52
1	A	160	VAL	CA-C-N	6.03	128.35	120.28
1	A	160	VAL	C-N-CA	6.03	128.35	120.28
1	D	189	THR	N-CA-C	-5.62	101.88	110.14
1	B	215	HIS	CA-C-N	5.22	125.88	120.03
1	B	215	HIS	C-N-CA	5.22	125.88	120.03
1	B	66	VAL	N-CA-C	5.17	115.75	108.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2756	0	2578	4	0
1	B	2736	0	2486	6	0
1	C	2741	0	2486	5	0
1	D	2730	0	2499	6	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	23	0	0	0	0
3	B	23	0	0	1	0
3	C	22	0	0	1	0
3	D	19	0	0	1	0
4	A	4	0	6	0	0
4	B	8	0	12	1	0
4	C	8	0	12	1	0
5	A	6	0	8	3	0
5	D	6	0	8	5	0
6	A	1	0	0	0	0
6	B	1	0	0	0	0
6	C	1	0	0	0	0
6	D	1	0	0	0	0
7	A	239	0	0	0	0
7	B	169	0	0	0	0
7	C	168	0	0	0	0
7	D	145	0	0	0	0
All	All	11815	0	10095	24	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (24) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:THR:HB	5:A:405:GOL:H2	1.42	1.00
1:D:126:THR:HB	5:D:404:GOL:H2	1.46	0.97

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:76:THR:HA	5:D:404:GOL:H32	1.59	0.84
1:A:76:THR:HA	5:A:405:GOL:H31	1.63	0.79
1:D:74:LEU:HD13	1:D:87:ILE:HD12	1.76	0.67
1:B:126:THR:HB	5:D:404:GOL:H12	1.89	0.55
1:B:126:THR:CB	5:D:404:GOL:H12	2.41	0.49
3:D:403:FO2:N2	3:D:403:FO2:C9	2.75	0.49
1:B:153:LEU:HD11	1:B:197:ILE:HG21	1.96	0.48
1:C:151:GLY:HA2	1:C:174:PRO:HG3	1.96	0.48
3:B:403:FO2:C9	3:B:403:FO2:N2	2.77	0.47
1:B:126:THR:OG1	5:D:404:GOL:H12	2.15	0.47
1:A:82:PHE:HB2	1:A:244:LEU:HB2	1.96	0.46
1:C:74:LEU:HD13	1:C:87:ILE:HD12	1.97	0.46
1:C:82:PHE:HB2	1:C:244:LEU:HB2	1.97	0.46
1:D:21:MET:HE1	1:D:53:TRP:HB2	1.98	0.46
1:D:150:ILE:HG22	1:D:174:PRO:HB3	1.99	0.44
3:C:403:FO2:N2	3:C:403:FO2:C9	2.81	0.43
1:C:150:ILE:HG22	1:C:174:PRO:HB3	2.00	0.43
1:B:114:PHE:HB2	4:B:404:DMS:H13	2.01	0.43
1:D:222:LEU:HG	1:D:226:PHE:CE2	2.55	0.42
1:C:225:GLY:HA3	4:C:404:DMS:H21	2.01	0.41
1:D:153:LEU:HD11	1:D:197:ILE:HG21	2.01	0.41
1:A:126:THR:CB	5:A:405:GOL:H2	2.32	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	337/360 (94%)	331 (98%)	6 (2%)	0	100	100
1	B	347/360 (96%)	341 (98%)	6 (2%)	0	100	100
1	C	345/360 (96%)	338 (98%)	7 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	345/360 (96%)	342 (99%)	3 (1%)	0	100	100
All	All	1374/1440 (95%)	1352 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/316 (88%)	272 (98%)	5 (2%)	51	66
1	B	265/316 (84%)	260 (98%)	5 (2%)	50	64
1	C	264/316 (84%)	258 (98%)	6 (2%)	44	58
1	D	265/316 (84%)	258 (97%)	7 (3%)	40	53
All	All	1071/1264 (85%)	1048 (98%)	23 (2%)	45	61

All (23) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	74	LEU
1	A	137	ASN
1	A	182	LYS
1	A	184	SER
1	A	270	THR
1	B	74	LEU
1	B	168	ILE
1	B	175	TYR
1	B	182	LYS
1	B	184	SER
1	C	7	THR
1	C	71	ILE
1	C	74	LEU
1	C	147	GLU
1	C	175	TYR
1	C	182	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	74	LEU
1	D	175	TYR
1	D	182	LYS
1	D	184	SER
1	D	248	LEU
1	D	249	MET
1	D	289	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	84	GLN
1	B	26	ASN
1	B	137	ASN
1	B	215	HIS
1	B	325	GLN
1	C	215	HIS
1	D	26	ASN
1	D	72	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 23 ligands modelled in this entry, 12 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FO2	C	403	2	24,25,26	0.52	1 (4%)	26,35,37	0.77	1 (3%)
4	DMS	C	404	-	3,3,3	0.31	0	3,3,3	0.24	0
3	FO2	A	403	2	25,26,26	0.60	1 (4%)	30,37,37	0.82	2 (6%)
5	GOL	D	404	-	5,5,5	0.16	0	5,5,5	0.47	0
4	DMS	A	404	-	3,3,3	0.29	0	3,3,3	0.27	0
3	FO2	D	403	2	20,21,26	0.62	1 (5%)	24,29,37	0.85	1 (4%)
4	DMS	C	405	-	3,3,3	0.28	0	3,3,3	0.24	0
5	GOL	A	405	-	5,5,5	0.22	0	5,5,5	0.61	0
4	DMS	B	404	-	3,3,3	0.23	0	3,3,3	0.22	0
4	DMS	B	405	-	3,3,3	0.31	0	3,3,3	0.30	0
3	FO2	B	403	2	25,26,26	0.48	0	30,37,37	0.82	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FO2	C	403	2	-	5/8/16/18	0/4/4/4
3	FO2	A	403	2	-	5/8/18/18	0/4/4/4
5	GOL	D	404	-	-	1/4/4/4	-
5	GOL	A	405	-	-	2/4/4/4	-
3	FO2	D	403	2	-	4/7/7/18	0/3/3/4
3	FO2	B	403	2	-	6/8/18/18	0/4/4/4

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	FO2	C5-N1	-2.45	1.33	1.38
3	D	403	FO2	C5-N1	-2.28	1.34	1.38
3	C	403	FO2	C5-N1	-2.11	1.34	1.38

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	403	FO2	C3-C2-N	2.84	124.45	118.55
3	B	403	FO2	C3-C2-N	2.81	124.37	118.55
3	C	403	FO2	C3-C2-N	2.58	123.89	118.55
3	A	403	FO2	C3-C2-N	2.25	123.21	118.55
3	A	403	FO2	C10-C8-C9	2.12	131.52	127.47

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	FO2	C3-C2-N3-N4
3	A	403	FO2	N-C2-N3-N4
3	B	403	FO2	C3-C2-N3-N4
3	B	403	FO2	N-C2-N3-N4
3	B	403	FO2	N-C2-N3-C9
3	C	403	FO2	C3-C2-N3-N4
3	C	403	FO2	N-C2-N3-N4
3	C	403	FO2	N-C2-N3-C9
3	D	403	FO2	C3-C2-N3-N4
3	D	403	FO2	N-C2-N3-N4
5	A	405	GOL	O1-C1-C2-C3
3	A	403	FO2	N-C2-N3-C9
3	D	403	FO2	N-C2-N3-C9
3	A	403	FO2	C3-C2-N3-C9
3	B	403	FO2	C3-C2-N3-C9
3	C	403	FO2	C3-C2-N3-C9
3	D	403	FO2	C3-C2-N3-C9
3	A	403	FO2	C11-C10-C8-C7
3	B	403	FO2	C14-C10-C8-C7
5	D	404	GOL	C1-C2-C3-O3
5	A	405	GOL	O1-C1-C2-O2
3	B	403	FO2	C11-C10-C8-C9
3	C	403	FO2	C11-C10-C8-C9

There are no ring outliers.

7 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	403	FO2	1	0
4	C	404	DMS	1	0
5	D	404	GOL	5	0
3	D	403	FO2	1	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	405	GOL	3	0
4	B	404	DMS	1	0
3	B	403	FO2	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand FO2 A 403



Ligand FO2 D 403





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/360 (94%)	-0.26	6 (1%) 67 65	20, 49, 76, 113	2 (0%)
1	B	348/360 (96%)	-0.00	6 (1%) 69 67	24, 59, 103, 127	1 (0%)
1	C	346/360 (96%)	-0.02	11 (3%) 50 47	21, 59, 101, 123	5 (1%)
1	D	345/360 (95%)	0.09	8 (2%) 61 59	30, 63, 102, 129	2 (0%)
All	All	1380/1440 (95%)	-0.05	31 (2%) 62 60	20, 58, 99, 129	10 (0%)

All (31) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	10	PRO	5.1
1	C	11[A]	SER	4.0
1	C	154[A]	ARG	3.9
1	C	167	THR	3.9
1	D	354	LEU	3.7
1	B	166	ILE	3.6
1	C	163	GLU	3.6
1	D	311	ASP	3.3
1	A	235[A]	GLU	3.0
1	D	11	SER	2.9
1	A	169	GLU	2.8
1	D	309	ARG	2.7
1	D	310	LYS	2.7
1	C	354	LEU	2.6
1	C	8	LEU	2.5
1	C	6	GLU	2.5
1	A	162	LYS	2.5
1	A	163	GLU	2.4
1	A	164	SER	2.4
1	B	7	THR	2.4
1	D	331	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	9	ASN	2.2
1	B	168	ILE	2.2
1	B	235	GLU	2.1
1	C	52	GLU	2.1
1	A	313	VAL	2.1
1	B	353	PHE	2.1
1	C	313	VAL	2.1
1	B	331	LEU	2.0
1	C	316	SER	2.0
1	D	12	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	DMS	B	404	4/4	0.70	0.30	59,61,62,62	4
4	DMS	A	404	4/4	0.73	0.24	132,132,132,132	0
4	DMS	C	405	4/4	0.75	0.33	74,75,75,75	4
4	DMS	C	404	4/4	0.76	0.22	132,132,132,132	0
4	DMS	B	405	4/4	0.85	0.21	111,111,112,112	0
5	GOL	A	405	6/6	0.88	0.14	31,35,37,38	6
6	CL	D	405	1/1	0.88	0.15	75,75,75,75	0
5	GOL	D	404	6/6	0.89	0.16	29,35,39,40	6
3	FO2	D	403	19/23	0.90	0.12	51,54,62,63	19
3	FO2	B	403	23/23	0.91	0.14	50,54,72,73	23
3	FO2	C	403	22/23	0.92	0.11	44,50,69,69	22
6	CL	B	406	1/1	0.92	0.17	86,86,86,86	0
3	FO2	A	403	23/23	0.92	0.11	35,38,62,64	23

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	CL	A	406	1/1	0.93	0.17	89,89,89,89	0
6	CL	C	406	1/1	0.96	0.10	98,98,98,98	0
2	ZN	D	402	1/1	0.99	0.05	55,55,55,55	1
2	ZN	A	401	1/1	0.99	0.04	38,38,38,38	1
2	ZN	B	402	1/1	0.99	0.03	59,59,59,59	1
2	ZN	C	402	1/1	0.99	0.03	52,52,52,52	1
2	ZN	A	402	1/1	1.00	0.03	47,47,47,47	1
2	ZN	C	401	1/1	1.00	0.04	44,44,44,44	1
2	ZN	B	401	1/1	1.00	0.03	50,50,50,50	1
2	ZN	D	401	1/1	1.00	0.02	50,50,50,50	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FO2 D 403:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FO2 B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FO2 C 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FO2 A 403:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.