



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 11:17 AM UTC

PDB ID : 6H4V / pdb_00006h4v
Title : Crystal structure of human KDM4A in complex with compound 34g
Authors : Le Bihan, Y.V.; van Montfort, R.L.M.
Deposited on : 2018-07-23
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	3% 92% • 5%
1	B	360	2% 91% • 6%
1	C	360	5% 90% 5% 5%
1	D	360	5% 93% • 5%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11784 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific demethylase 4A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	341	Total	C	N	O	S	0	2	0
			2725	1778	439	494	14			
1	B	340	Total	C	N	O	S	0	4	0
			2777	1803	464	494	16			
1	C	343	Total	C	N	O	S	0	5	0
			2743	1779	449	499	16			
1	D	343	Total	C	N	O	S	0	4	0
			2715	1764	445	492	14			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP O75164
B	0	SER	-	expression tag	UNP O75164
C	0	SER	-	expression tag	UNP O75164
D	0	SER	-	expression tag	UNP O75164

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		
2	B	2	Total	Zn	0	0
			2	2		
2	C	2	Total	Zn	0	0
			2	2		
2	D	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 8-[4-(1-cyclopentylpiperidin-4-yl)pyrazol-1-yl]-3 {H}-pyrido[3,4-d]pyrimidin-4-one (CCD ID: FQN) (formula: C₂₀H₂₄N₆O) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			22	15	6	1		
3	B	1	Total	C	N	O	0	0
			27	20	6	1		
3	C	1	Total	C	N	O	0	0
			22	15	6	1		
3	D	1	Total	C	N	O	0	0
			27	20	6	1		

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	C	1	Total	Cl	0	0
			1	1		
4	D	1	Total	Cl	0	0
			1	1		

- Molecule 5 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	B	1	Total	C	O	S	0	0
			4	2	1	1		
5	D	1	Total	C	O	S	0	0
			4	2	1	1		
5	D	1	Total	C	O	S	0	0
			4	2	1	1		
5	D	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	175	Total	O	0	3
			178	178		
6	B	208	Total	O	0	1
			209	209		
6	C	161	Total	O	0	5
			166	166		
6	D	134	Total	O	0	4
			138	138		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lysine-specific demethylase 4A




- Molecule 1: Lysine-specific demethylase 4A



- Molecule 1: Lysine-specific demethylase 4A



- Molecule 1: Lysine-specific demethylase 4A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.83Å 103.78Å 144.03Å 90.00° 99.45° 90.00°	Depositor
Resolution (Å)	49.07 – 2.15 49.07 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.07-2.15) 100.0 (49.07-2.15)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.59 (at 2.16Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.176 , 0.211 0.180 , 0.210	Depositor DCC
R_{free} test set	4446 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	47.5	Xtriage
Anisotropy	0.112	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 72.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11784	wwPDB-VP
Average B, all atoms (Å ²)	61.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 24.14 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.0601e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FQN, DMS, ZN, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	0/2811	1.24	1/3825 (0.0%)
1	B	0.95	0/2863	1.23	5/3886 (0.1%)
1	C	0.90	1/2828 (0.0%)	1.26	5/3848 (0.1%)
1	D	0.91	0/2801	1.24	3/3820 (0.1%)
All	All	0.92	1/11303 (0.0%)	1.24	14/15379 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	211	VAL	CA-C	5.50	1.56	1.52

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	184	SER	N-CA-C	6.55	119.67	109.52
1	D	184	SER	N-CA-C	6.49	119.58	109.52
1	B	184	SER	N-CA-C	6.38	119.42	109.52
1	A	184	SER	N-CA-C	6.09	118.96	109.52
1	D	129	PRO	N-CA-C	5.47	117.37	110.70
1	B	215	HIS	CA-C-N	5.43	126.01	119.98
1	B	215	HIS	C-N-CA	5.43	126.01	119.98
1	C	186	ALA	N-CA-C	5.28	116.91	110.41
1	C	165	GLY	CA-C-N	5.20	128.29	120.69
1	C	165	GLY	C-N-CA	5.20	128.29	120.69
1	C	189	THR	N-CA-C	-5.17	102.54	110.14
1	D	186	ALA	N-CA-C	5.15	117.33	110.53
1	B	149	ASN	CA-C-N	5.04	128.89	120.62
1	B	149	ASN	C-N-CA	5.04	128.89	120.62

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2725	0	2503	4	0
1	B	2777	0	2616	2	0
1	C	2743	0	2498	4	0
1	D	2715	0	2447	5	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	22	0	0	1	0
3	B	27	0	0	1	0
3	C	22	0	0	1	0
3	D	27	0	0	1	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	8	0	12	0	0
5	B	4	0	6	0	0
5	D	12	0	18	0	0
6	A	178	0	0	0	0
6	B	209	0	0	0	0
6	C	166	0	0	0	0
6	D	138	0	0	0	0
All	All	11784	0	10100	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ARG:HH21	1:A:218:ARG:HG2	1.29	0.94
1:D:15:MET:CE	1:D:17:PHE:HZ	2.05	0.69

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:15:MET:CE	1:D:17:PHE:CZ	2.81	0.63
1:A:218:ARG:HG2	1:A:218:ARG:NH2	2.06	0.62
3:D:403:FQN:C9	3:D:403:FQN:N2	2.73	0.51
1:C:74:LEU:HD23	1:C:87:ILE:HD12	1.91	0.51
3:C:403:FQN:C9	3:C:403:FQN:N2	2.73	0.51
1:B:79:SER:O	1:B:249:MET:HE3	2.13	0.49
1:B:82:PHE:HB2	1:B:244:LEU:HB2	1.93	0.49
3:B:403:FQN:C9	3:B:403:FQN:N2	2.77	0.48
1:D:15:MET:HE2	1:D:17:PHE:CZ	2.50	0.47
1:D:15:MET:HE3	1:D:17:PHE:CZ	2.49	0.46
3:A:403:FQN:C9	3:A:403:FQN:N2	2.78	0.46
1:A:153:LEU:HD11	1:A:197:ILE:HG21	2.00	0.44
1:C:150:ILE:HG22	1:C:174:PRO:HB3	2.01	0.43
1:C:306:CYS:HB3	1:C:312:MET:HG3	2.00	0.42
1:C:166:ILE:H	1:C:166:ILE:HG13	1.71	0.41
1:D:150:ILE:HG22	1:D:174:PRO:HB3	2.03	0.40
1:A:82:PHE:HB2	1:A:244:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/360 (94%)	333 (98%)	6 (2%)	0	100	100
1	B	340/360 (94%)	336 (99%)	4 (1%)	0	100	100
1	C	344/360 (96%)	337 (98%)	7 (2%)	0	100	100
1	D	343/360 (95%)	337 (98%)	6 (2%)	0	100	100
All	All	1366/1440 (95%)	1343 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/316 (84%)	260 (98%)	5 (2%)	50	56
1	B	279/316 (88%)	274 (98%)	5 (2%)	51	58
1	C	266/316 (84%)	258 (97%)	8 (3%)	36	38
1	D	259/316 (82%)	256 (99%)	3 (1%)	63	70
All	All	1069/1264 (85%)	1048 (98%)	21 (2%)	48	54

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	54	LYS
1	A	171	VAL
1	A	184	SER
1	A	218	ARG
1	A	345	LEU
1	B	65	LEU
1	B	74	LEU
1	B	137	ASN
1	B	182	LYS
1	B	345	LEU
1	C	58	SER
1	C	65	LEU
1	C	175	TYR
1	C	182	LYS
1	C	184	SER
1	C	248	LEU
1	C	249	MET
1	C	338	ASN
1	D	104	ASP
1	D	182	LYS
1	D	184	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	137	ASN
1	A	325	GLN
1	B	124	ASN
1	B	128	ASN
1	C	84	GLN
1	C	124	ASN
1	D	84	GLN
1	D	215	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 21 ligands modelled in this entry, 11 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	DMS	D	405	-	3,3,3	0.30	0	3,3,3	0.22	0
3	FQN	A	403	2	24,25,31	0.47	0	26,35,44	0.83	1 (3%)
5	DMS	A	406	-	3,3,3	0.27	0	3,3,3	0.26	0
3	FQN	C	403	2	24,25,31	0.57	1 (4%)	26,35,44	0.85	1 (3%)
3	FQN	D	403	2	30,31,31	0.51	1 (3%)	37,44,44	0.84	2 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	DMS	B	404	-	3,3,3	0.27	0	3,3,3	0.20	0
3	FQN	B	403	2	30,31,31	0.57	1 (3%)	37,44,44	0.92	2 (5%)
5	DMS	D	407	-	3,3,3	0.33	0	3,3,3	0.28	0
5	DMS	D	406	-	3,3,3	0.28	0	3,3,3	0.24	0
5	DMS	A	405	-	3,3,3	0.31	0	3,3,3	0.35	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FQN	B	403	2	-	8/12/29/29	0/5/5/5
3	FQN	A	403	2	-	5/8/16/29	0/4/4/5
3	FQN	C	403	2	-	5/8/16/29	0/4/4/5
3	FQN	D	403	2	-	9/12/29/29	0/5/5/5

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	403	FQN	C5-N1	-2.61	1.33	1.38
3	C	403	FQN	C5-N1	-2.37	1.33	1.38
3	D	403	FQN	C5-N1	-2.36	1.33	1.38

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	403	FQN	C3-C2-N	3.11	125.00	118.55
3	B	403	FQN	C3-C2-N	3.09	124.97	118.55
3	D	403	FQN	C3-C2-N	2.94	124.64	118.55
3	A	403	FQN	C3-C2-N	2.86	124.48	118.55
3	D	403	FQN	C19-C15-C16	-2.72	98.47	104.74
3	B	403	FQN	C19-C15-C16	-2.34	99.35	104.74

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	FQN	C3-C2-N3-N4
3	A	403	FQN	N-C2-N3-C9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	403	FQN	N-C2-N3-N4
3	B	403	FQN	C3-C2-N3-N4
3	B	403	FQN	N-C2-N3-C9
3	B	403	FQN	N-C2-N3-N4
3	B	403	FQN	C16-C15-N5-C13
3	B	403	FQN	C19-C15-N5-C13
3	B	403	FQN	C16-C15-N5-C12
3	B	403	FQN	C19-C15-N5-C12
3	C	403	FQN	C11-C10-C8-C9
3	C	403	FQN	C3-C2-N3-N4
3	C	403	FQN	N-C2-N3-C9
3	C	403	FQN	N-C2-N3-N4
3	D	403	FQN	C11-C10-C8-C9
3	D	403	FQN	C3-C2-N3-N4
3	D	403	FQN	N-C2-N3-C9
3	D	403	FQN	N-C2-N3-N4
3	D	403	FQN	C16-C15-N5-C13
3	D	403	FQN	C19-C15-N5-C13
3	D	403	FQN	C16-C15-N5-C12
3	D	403	FQN	C19-C15-N5-C12
3	A	403	FQN	C3-C2-N3-C9
3	B	403	FQN	C3-C2-N3-C9
3	C	403	FQN	C3-C2-N3-C9
3	D	403	FQN	C3-C2-N3-C9
3	A	403	FQN	C14-C10-C8-C9


There are no ring outliers.


4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	403	FQN	1	0
3	C	403	FQN	1	0
3	D	403	FQN	1	0
3	B	403	FQN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	341/360 (94%)	0.20	10 (2%) 53 57	22, 56, 98, 117	3 (0%)
1	B	340/360 (94%)	0.02	6 (1%) 67 71	19, 50, 77, 118	4 (1%)
1	C	343/360 (95%)	0.43	19 (5%) 30 34	21, 62, 106, 127	5 (1%)
1	D	343/360 (95%)	0.36	18 (5%) 33 36	19, 61, 108, 124	4 (1%)
All	All	1367/1440 (94%)	0.25	53 (3%) 43 48	19, 57, 100, 127	16 (1%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	161	GLU	5.0
1	B	161	GLU	4.3
1	C	354	LEU	4.2
1	C	311	ASP	3.7
1	A	7	THR	3.4
1	B	353	PHE	3.4
1	C	166	ILE	3.4
1	B	7	THR	3.3
1	C	170	GLY	3.3
1	C	11	SER	3.2
1	D	170	GLY	3.2
1	D	354	LEU	3.1
1	D	11[A]	SER	2.9
1	C	313	VAL	2.9
1	C	144	HIS	2.7
1	D	168	ILE	2.7
1	D	312	MET	2.6
1	A	6	GLU	2.6
1	A	8	LEU	2.6
1	C	226	PHE	2.6
1	C	171	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	167	THR	2.6
1	D	8	LEU	2.6
1	D	6	GLU	2.5
1	D	171	VAL	2.5
1	A	354	LEU	2.5
1	D	311	ASP	2.4
1	C	312	MET	2.4
1	B	354	LEU	2.4
1	D	227	PHE	2.4
1	D	10	PRO	2.4
1	A	329	TYR	2.3
1	D	226	PHE	2.3
1	B	170	GLY	2.3
1	C	309	ARG	2.3
1	C	12	ALA	2.3
1	A	161	GLU	2.2
1	A	351	ALA	2.2
1	D	329	TYR	2.2
1	C	106	TYR	2.2
1	D	113	GLU	2.1
1	C	104	ASP	2.1
1	C	227	PHE	2.1
1	A	170	GLY	2.1
1	B	312	MET	2.0
1	D	160	VAL	2.0
1	A	303	ALA	2.0
1	D	12	ALA	2.0
1	A	125	LEU	2.0
1	C	105	LYS	2.0
1	D	333	LYS	2.0
1	C	175	TYR	2.0
1	C	329	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	DMS	D	407	4/4	0.57	0.22	111,111,111,111	0
5	DMS	A	406	4/4	0.60	0.38	71,73,73,73	4
5	DMS	D	406	4/4	0.64	0.36	79,80,81,81	4
5	DMS	B	404	4/4	0.69	0.29	102,102,103,103	4
5	DMS	A	405	4/4	0.73	0.23	138,138,138,139	0
5	DMS	D	405	4/4	0.75	0.21	94,94,95,95	4
4	CL	D	404	1/1	0.86	0.17	126,126,126,126	0
3	FQN	D	403	27/27	0.89	0.14	49,62,83,84	27
3	FQN	A	403	22/27	0.89	0.14	43,47,68,69	22
4	CL	A	404	1/1	0.90	0.12	90,90,90,90	0
4	CL	C	404	1/1	0.90	0.17	83,83,83,83	0
3	FQN	C	403	22/27	0.90	0.13	43,46,60,62	22
3	FQN	B	403	27/27	0.93	0.12	37,48,63,63	27
2	ZN	C	401	1/1	0.98	0.06	47,47,47,47	1
2	ZN	D	402	1/1	0.98	0.04	57,57,57,57	1
2	ZN	A	402	1/1	0.98	0.04	58,58,58,58	1
2	ZN	C	402	1/1	0.99	0.04	56,56,56,56	1
2	ZN	B	401	1/1	0.99	0.06	39,39,39,39	1
2	ZN	A	401	1/1	0.99	0.04	44,44,44,44	1
2	ZN	B	402	1/1	1.00	0.03	43,43,43,43	1
2	ZN	D	401	1/1	1.00	0.03	43,43,43,43	1




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around FQN D 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FQN A 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around FQN C 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around FQN B 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.