



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:36 PM UTC

PDB ID : 5H6M / pdb_00005h6m
Title : DNA targeting ADP-ribosyltransferase Pierisin-1
Authors : Oda, T.; Hirabayashi, H.; Shikauchi, G.; Takamura, R.; Hiraga, K.; Minami, H.; Hashimoto, H.; Yamamoto, M.; Wakabayashi, K.; Sugimura, T.; Shimizu, T.; Sato, M.
Deposited on : 2016-11-14
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	271	14% 82% 18%
1	B	271	15% 82% 18%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	B	303	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pierisin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	223	Total	C	N	O	S	0	0	0
			1833	1162	337	330	4			
1	B	222	Total	C	N	O	S	0	0	0
			1822	1155	334	329	4			

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-37	GLY	-	expression tag	UNP H3JU00
A	-36	SER	-	expression tag	UNP H3JU00
A	-35	GLY	-	expression tag	UNP H3JU00
A	-34	MET	-	expression tag	UNP H3JU00
A	-33	LYS	-	expression tag	UNP H3JU00
A	-32	GLU	-	expression tag	UNP H3JU00
A	-31	THR	-	expression tag	UNP H3JU00
A	-30	ALA	-	expression tag	UNP H3JU00
A	-29	ALA	-	expression tag	UNP H3JU00
A	-28	ALA	-	expression tag	UNP H3JU00
A	-27	LYS	-	expression tag	UNP H3JU00
A	-26	PHE	-	expression tag	UNP H3JU00
A	-25	GLU	-	expression tag	UNP H3JU00
A	-24	ARG	-	expression tag	UNP H3JU00
A	-23	GLN	-	expression tag	UNP H3JU00
A	-22	HIS	-	expression tag	UNP H3JU00
A	-21	MET	-	expression tag	UNP H3JU00
A	-20	ASP	-	expression tag	UNP H3JU00
A	-19	SER	-	expression tag	UNP H3JU00
A	-18	PRO	-	expression tag	UNP H3JU00
A	-17	ASP	-	expression tag	UNP H3JU00
A	-16	LEU	-	expression tag	UNP H3JU00
A	-15	GLY	-	expression tag	UNP H3JU00
A	-14	THR	-	expression tag	UNP H3JU00
A	-13	ASP	-	expression tag	UNP H3JU00

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	-12	ASP	-	expression tag	UNP H3JU00
A	-11	ASP	-	expression tag	UNP H3JU00
A	-10	ASP	-	expression tag	UNP H3JU00
A	-9	LYS	-	expression tag	UNP H3JU00
A	-8	ALA	-	expression tag	UNP H3JU00
A	-7	MET	-	expression tag	UNP H3JU00
A	-6	ALA	-	expression tag	UNP H3JU00
A	-5	ASP	-	expression tag	UNP H3JU00
A	-4	ILE	-	expression tag	UNP H3JU00
A	-3	GLY	-	expression tag	UNP H3JU00
A	-2	SER	-	expression tag	UNP H3JU00
A	-1	GLU	-	expression tag	UNP H3JU00
A	0	PHE	-	expression tag	UNP H3JU00
A	165	GLN	GLU	engineered mutation	UNP H3JU00
B	-37	GLY	-	expression tag	UNP H3JU00
B	-36	SER	-	expression tag	UNP H3JU00
B	-35	GLY	-	expression tag	UNP H3JU00
B	-34	MET	-	expression tag	UNP H3JU00
B	-33	LYS	-	expression tag	UNP H3JU00
B	-32	GLU	-	expression tag	UNP H3JU00
B	-31	THR	-	expression tag	UNP H3JU00
B	-30	ALA	-	expression tag	UNP H3JU00
B	-29	ALA	-	expression tag	UNP H3JU00
B	-28	ALA	-	expression tag	UNP H3JU00
B	-27	LYS	-	expression tag	UNP H3JU00
B	-26	PHE	-	expression tag	UNP H3JU00
B	-25	GLU	-	expression tag	UNP H3JU00
B	-24	ARG	-	expression tag	UNP H3JU00
B	-23	GLN	-	expression tag	UNP H3JU00
B	-22	HIS	-	expression tag	UNP H3JU00
B	-21	MET	-	expression tag	UNP H3JU00
B	-20	ASP	-	expression tag	UNP H3JU00
B	-19	SER	-	expression tag	UNP H3JU00
B	-18	PRO	-	expression tag	UNP H3JU00
B	-17	ASP	-	expression tag	UNP H3JU00
B	-16	LEU	-	expression tag	UNP H3JU00
B	-15	GLY	-	expression tag	UNP H3JU00
B	-14	THR	-	expression tag	UNP H3JU00
B	-13	ASP	-	expression tag	UNP H3JU00
B	-12	ASP	-	expression tag	UNP H3JU00
B	-11	ASP	-	expression tag	UNP H3JU00
B	-10	ASP	-	expression tag	UNP H3JU00

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-9	LYS	-	expression tag	UNP H3JU00
B	-8	ALA	-	expression tag	UNP H3JU00
B	-7	MET	-	expression tag	UNP H3JU00
B	-6	ALA	-	expression tag	UNP H3JU00
B	-5	ASP	-	expression tag	UNP H3JU00
B	-4	ILE	-	expression tag	UNP H3JU00
B	-3	GLY	-	expression tag	UNP H3JU00
B	-2	SER	-	expression tag	UNP H3JU00
B	-1	GLU	-	expression tag	UNP H3JU00
B	0	PHE	-	expression tag	UNP H3JU00
B	165	GLN	GLU	engineered mutation	UNP H3JU00

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0
2	B	1	Total 4	C 2	O 2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	138	Total 138	O 138	0	0
3	B	136	Total 136	O 136	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Pierisin-1



• Molecule 1: Pierisin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	134.72Å 134.72Å 73.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.54 – 1.90 49.54 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (49.54-1.90) 99.9 (49.54-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.91 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.8.0103	Depositor
R, R_{free}	0.203 , 0.219 (Not available) , 0.226	Depositor DCC
R_{free} test set	3075 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	21.5	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 43.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.019 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4053	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/1885	0.69	0/2573
1	B	0.51	0/1874	0.70	0/2559
All	All	0.51	0/3759	0.69	0/5132

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1833	0	1792	0	0
1	B	1822	0	1773	0	0
2	A	84	0	126	0	0
2	B	40	0	60	0	0
3	A	138	0	0	0	0
3	B	136	0	0	0	0
All	All	4053	0	3751	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/271 (82%)	221 (100%)	0	0	100	100
1	B	218/271 (80%)	217 (100%)	1 (0%)	0	100	100
All	All	439/542 (81%)	438 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/237 (84%)	199 (100%)	0	100	100
1	B	198/237 (84%)	197 (100%)	1 (0%)	81	84
All	All	397/474 (84%)	396 (100%)	1 (0%)	86	88

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	92	ASN
1	A	131	ASN
1	A	189	GLN
1	B	119	GLN
1	B	139	GLN
1	B	189	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

31 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	305	-	3,3,3	0.43	0	2,2,2	0.29	0
2	EDO	A	306	-	3,3,3	0.45	0	2,2,2	0.24	0
2	EDO	A	307	-	3,3,3	0.44	0	2,2,2	0.31	0
2	EDO	A	304	-	3,3,3	0.36	0	2,2,2	0.44	0
2	EDO	A	310	-	3,3,3	0.47	0	2,2,2	0.25	0
2	EDO	B	305	-	3,3,3	0.45	0	2,2,2	0.21	0
2	EDO	A	320	-	3,3,3	0.40	0	2,2,2	0.37	0
2	EDO	B	301	-	3,3,3	0.37	0	2,2,2	0.45	0
2	EDO	A	317	-	3,3,3	0.43	0	2,2,2	0.25	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	B	304	-	3,3,3	0.46	0	2,2,2	0.15	0
2	EDO	A	318	-	3,3,3	0.41	0	2,2,2	0.35	0
2	EDO	B	302	-	3,3,3	0.45	0	2,2,2	0.24	0
2	EDO	A	309	-	3,3,3	0.42	0	2,2,2	0.33	0
2	EDO	A	314	-	3,3,3	0.42	0	2,2,2	0.36	0
2	EDO	A	319	-	3,3,3	0.42	0	2,2,2	0.35	0
2	EDO	A	308	-	3,3,3	0.42	0	2,2,2	0.35	0
2	EDO	A	303	-	3,3,3	0.46	0	2,2,2	0.09	0
2	EDO	A	311	-	3,3,3	0.44	0	2,2,2	0.32	0
2	EDO	A	301	-	3,3,3	0.44	0	2,2,2	0.25	0
2	EDO	A	315	-	3,3,3	0.42	0	2,2,2	0.36	0
2	EDO	B	306	-	3,3,3	0.44	0	2,2,2	0.31	0
2	EDO	B	309	-	3,3,3	0.45	0	2,2,2	0.28	0
2	EDO	B	310	-	3,3,3	0.35	0	2,2,2	0.32	0
2	EDO	B	303	-	3,3,3	0.43	0	2,2,2	0.35	0
2	EDO	A	312	-	3,3,3	0.43	0	2,2,2	0.26	0
2	EDO	B	307	-	3,3,3	0.41	0	2,2,2	0.36	0
2	EDO	A	313	-	3,3,3	0.46	0	2,2,2	0.26	0
2	EDO	B	308	-	3,3,3	0.45	0	2,2,2	0.31	0
2	EDO	A	321	-	3,3,3	0.43	0	2,2,2	0.34	0
2	EDO	A	316	-	3,3,3	0.44	0	2,2,2	0.29	0
2	EDO	A	302	-	3,3,3	0.42	0	2,2,2	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	305	-	-	0/1/1/1	-
2	EDO	A	306	-	-	0/1/1/1	-
2	EDO	A	307	-	-	1/1/1/1	-
2	EDO	A	304	-	-	1/1/1/1	-
2	EDO	A	310	-	-	0/1/1/1	-
2	EDO	B	305	-	-	1/1/1/1	-
2	EDO	A	320	-	-	1/1/1/1	-
2	EDO	B	301	-	-	0/1/1/1	-
2	EDO	A	317	-	-	0/1/1/1	-
2	EDO	B	304	-	-	1/1/1/1	-
2	EDO	A	318	-	-	0/1/1/1	-
2	EDO	B	302	-	-	0/1/1/1	-
2	EDO	A	309	-	-	1/1/1/1	-
2	EDO	A	314	-	-	1/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	319	-	-	0/1/1/1	-
2	EDO	A	308	-	-	1/1/1/1	-
2	EDO	A	303	-	-	1/1/1/1	-
2	EDO	A	311	-	-	1/1/1/1	-
2	EDO	A	301	-	-	0/1/1/1	-
2	EDO	A	315	-	-	1/1/1/1	-
2	EDO	B	306	-	-	1/1/1/1	-
2	EDO	B	309	-	-	0/1/1/1	-
2	EDO	B	310	-	-	1/1/1/1	-
2	EDO	B	303	-	-	0/1/1/1	-
2	EDO	A	312	-	-	0/1/1/1	-
2	EDO	B	307	-	-	0/1/1/1	-
2	EDO	A	313	-	-	1/1/1/1	-
2	EDO	B	308	-	-	1/1/1/1	-
2	EDO	A	321	-	-	0/1/1/1	-
2	EDO	A	316	-	-	0/1/1/1	-
2	EDO	A	302	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	304	EDO	O1-C1-C2-O2
2	A	307	EDO	O1-C1-C2-O2
2	A	314	EDO	O1-C1-C2-O2
2	B	310	EDO	O1-C1-C2-O2
2	B	306	EDO	O1-C1-C2-O2
2	B	308	EDO	O1-C1-C2-O2
2	A	315	EDO	O1-C1-C2-O2
2	A	303	EDO	O1-C1-C2-O2
2	A	311	EDO	O1-C1-C2-O2
2	A	308	EDO	O1-C1-C2-O2
2	A	309	EDO	O1-C1-C2-O2
2	A	313	EDO	O1-C1-C2-O2
2	A	320	EDO	O1-C1-C2-O2
2	B	305	EDO	O1-C1-C2-O2
2	A	302	EDO	O1-C1-C2-O2
2	B	304	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	223/271 (82%)	0.71	37 (16%) 4 4	13, 20, 36, 56	0
1	B	222/271 (81%)	0.81	40 (18%) 3 3	13, 20, 41, 59	0
All	All	445/542 (82%)	0.76	77 (17%) 4 4	13, 20, 39, 59	0

All (77) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	124	LYS	6.9
1	B	121	ASN	6.1
1	B	229	GLN	5.7
1	A	209	SER	5.5
1	A	229	GLN	5.1
1	B	125	TYR	4.8
1	B	207	VAL	4.8
1	B	210	SER	4.7
1	A	208	SER	4.6
1	B	209	SER	4.5
1	B	130	ARG	4.5
1	B	206	ILE	4.5
1	A	92	ASN	4.3
1	B	6	PRO	4.3
1	B	211	ARG	4.1
1	A	207	VAL	4.1
1	B	128	THR	4.1
1	A	210	SER	4.1
1	B	131	ASN	4.0
1	A	187	ARG	3.8
1	B	184	HIS	3.7
1	A	38	MET	3.6
1	B	127	TRP	3.6
1	B	24	ASP	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	22	ALA	3.5
1	A	211	ARG	3.3
1	B	208	SER	3.3
1	B	92	ASN	3.2
1	B	8	MET	3.2
1	A	131	ASN	3.1
1	A	24	ASP	3.1
1	B	22	ALA	3.1
1	A	219	MET	3.0
1	A	18	GLU	3.0
1	A	7	TYR	2.9
1	A	213	PRO	2.9
1	B	126	VAL	2.8
1	B	120	ARG	2.8
1	B	18	GLU	2.8
1	A	8	MET	2.8
1	B	133	ASN	2.8
1	B	204	GLU	2.8
1	A	23	LEU	2.7
1	B	58	ASN	2.7
1	B	95	TRP	2.7
1	B	219	MET	2.7
1	A	206	ILE	2.7
1	B	26	GLU	2.6
1	B	220	ASN	2.6
1	A	26	GLU	2.6
1	A	134	ARG	2.6
1	B	21	ARG	2.6
1	A	123	LYS	2.6
1	A	220	ASN	2.6
1	B	212	THR	2.6
1	A	188	ILE	2.5
1	A	43	THR	2.5
1	A	212	THR	2.5
1	A	155	SER	2.5
1	B	108	HIS	2.4
1	B	152	ASP	2.4
1	A	91	GLU	2.4
1	B	134	ARG	2.3
1	A	58	ASN	2.3
1	A	121	ASN	2.3
1	A	184	HIS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	185	ASN	2.3
1	A	120	ARG	2.3
1	B	23	LEU	2.3
1	A	95	TRP	2.2
1	B	129	PRO	2.2
1	B	7	TYR	2.2
1	B	71	TRP	2.1
1	A	156	ASP	2.1
1	A	47	ARG	2.0
1	B	132	ALA	2.0
1	A	27	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	B	303	4/4	0.42	0.40	56,57,57,57	0
2	EDO	A	314	4/4	0.62	0.25	48,48,48,49	0
2	EDO	A	308	4/4	0.62	0.38	37,37,38,39	0
2	EDO	A	315	4/4	0.66	0.39	54,54,54,54	0
2	EDO	B	302	4/4	0.68	0.24	39,39,39,39	0
2	EDO	A	302	4/4	0.68	0.27	40,40,40,41	0
2	EDO	A	311	4/4	0.70	0.27	51,51,51,52	0
2	EDO	A	313	4/4	0.70	0.31	43,44,44,44	0
2	EDO	A	306	4/4	0.70	0.31	45,45,45,45	0
2	EDO	B	306	4/4	0.70	0.28	58,58,58,58	0
2	EDO	A	317	4/4	0.71	0.29	40,40,41,41	0
2	EDO	A	305	4/4	0.71	0.27	51,51,51,52	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	320	4/4	0.72	0.24	31,33,34,35	0
2	EDO	A	307	4/4	0.75	0.22	42,42,43,43	0
2	EDO	B	307	4/4	0.75	0.23	40,40,41,41	0
2	EDO	A	310	4/4	0.76	0.23	45,46,46,46	0
2	EDO	B	308	4/4	0.77	0.25	36,36,36,36	0
2	EDO	A	318	4/4	0.78	0.20	48,48,49,49	0
2	EDO	A	319	4/4	0.78	0.34	58,58,58,59	0
2	EDO	A	321	4/4	0.79	0.20	46,46,47,47	0
2	EDO	B	305	4/4	0.80	0.27	48,48,48,48	0
2	EDO	A	316	4/4	0.81	0.32	58,58,58,58	0
2	EDO	A	312	4/4	0.82	0.19	45,46,46,46	0
2	EDO	B	304	4/4	0.83	0.18	25,26,27,27	0
2	EDO	B	301	4/4	0.84	0.17	29,30,30,31	0
2	EDO	B	309	4/4	0.84	0.24	53,53,54,54	0
2	EDO	B	310	4/4	0.84	0.23	25,27,27,28	0
2	EDO	A	303	4/4	0.85	0.18	26,28,28,29	0
2	EDO	A	301	4/4	0.86	0.21	39,39,40,40	0
2	EDO	A	309	4/4	0.87	0.17	35,35,36,36	0
2	EDO	A	304	4/4	0.90	0.12	23,23,23,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.