



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 07:11 PM UTC

PDB ID : 5H6S / pdb_00005h6s
Title : Crystal structure of Hydrazidase S179A mutant complexed with a substrate
Authors : Akiyama, T.; Ishii, M.; Takuwa, A.; Oinuma, K.; Sasaki, Y.; Takaya, N.; Yajima, S.
Deposited on : 2016-11-15
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	2% 92% • 5%
1	B	481	% 91% • 5%
1	C	481	4% 91% • 5%
1	D	481	3% 93% • 5%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 14559 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	456	Total	C	N	O	S	0	2	0
			3447	2148	622	663	14			
1	B	456	Total	C	N	O	S	0	2	0
			3447	2148	622	663	14			
1	C	457	Total	C	N	O	S	0	1	0
			3447	2148	621	664	14			
1	D	456	Total	C	N	O	S	0	2	0
			3447	2147	621	665	14			

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-20	GLY	-	expression tag	UNP A0A170QJP8
A	-19	SER	-	expression tag	UNP A0A170QJP8
A	-18	SER	-	expression tag	UNP A0A170QJP8
A	-17	HIS	-	expression tag	UNP A0A170QJP8
A	-16	HIS	-	expression tag	UNP A0A170QJP8
A	-15	HIS	-	expression tag	UNP A0A170QJP8
A	-14	HIS	-	expression tag	UNP A0A170QJP8
A	-13	HIS	-	expression tag	UNP A0A170QJP8
A	-12	HIS	-	expression tag	UNP A0A170QJP8
A	-11	SER	-	expression tag	UNP A0A170QJP8
A	-10	SER	-	expression tag	UNP A0A170QJP8
A	-9	GLY	-	expression tag	UNP A0A170QJP8
A	-8	LEU	-	expression tag	UNP A0A170QJP8
A	-7	VAL	-	expression tag	UNP A0A170QJP8
A	-6	PRO	-	expression tag	UNP A0A170QJP8
A	-5	ARG	-	expression tag	UNP A0A170QJP8
A	-4	GLY	-	expression tag	UNP A0A170QJP8
A	-3	SER	-	expression tag	UNP A0A170QJP8
A	-2	HIS	-	expression tag	UNP A0A170QJP8
A	-1	MET	-	expression tag	UNP A0A170QJP8
A	0	ALA	-	expression tag	UNP A0A170QJP8

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP A0A170QJP8
B	-20	GLY	-	expression tag	UNP A0A170QJP8
B	-19	SER	-	expression tag	UNP A0A170QJP8
B	-18	SER	-	expression tag	UNP A0A170QJP8
B	-17	HIS	-	expression tag	UNP A0A170QJP8
B	-16	HIS	-	expression tag	UNP A0A170QJP8
B	-15	HIS	-	expression tag	UNP A0A170QJP8
B	-14	HIS	-	expression tag	UNP A0A170QJP8
B	-13	HIS	-	expression tag	UNP A0A170QJP8
B	-12	HIS	-	expression tag	UNP A0A170QJP8
B	-11	SER	-	expression tag	UNP A0A170QJP8
B	-10	SER	-	expression tag	UNP A0A170QJP8
B	-9	GLY	-	expression tag	UNP A0A170QJP8
B	-8	LEU	-	expression tag	UNP A0A170QJP8
B	-7	VAL	-	expression tag	UNP A0A170QJP8
B	-6	PRO	-	expression tag	UNP A0A170QJP8
B	-5	ARG	-	expression tag	UNP A0A170QJP8
B	-4	GLY	-	expression tag	UNP A0A170QJP8
B	-3	SER	-	expression tag	UNP A0A170QJP8
B	-2	HIS	-	expression tag	UNP A0A170QJP8
B	-1	MET	-	expression tag	UNP A0A170QJP8
B	0	ALA	-	expression tag	UNP A0A170QJP8
B	1	SER	-	expression tag	UNP A0A170QJP8
C	-20	GLY	-	expression tag	UNP A0A170QJP8
C	-19	SER	-	expression tag	UNP A0A170QJP8
C	-18	SER	-	expression tag	UNP A0A170QJP8
C	-17	HIS	-	expression tag	UNP A0A170QJP8
C	-16	HIS	-	expression tag	UNP A0A170QJP8
C	-15	HIS	-	expression tag	UNP A0A170QJP8
C	-14	HIS	-	expression tag	UNP A0A170QJP8
C	-13	HIS	-	expression tag	UNP A0A170QJP8
C	-12	HIS	-	expression tag	UNP A0A170QJP8
C	-11	SER	-	expression tag	UNP A0A170QJP8
C	-10	SER	-	expression tag	UNP A0A170QJP8
C	-9	GLY	-	expression tag	UNP A0A170QJP8
C	-8	LEU	-	expression tag	UNP A0A170QJP8
C	-7	VAL	-	expression tag	UNP A0A170QJP8
C	-6	PRO	-	expression tag	UNP A0A170QJP8
C	-5	ARG	-	expression tag	UNP A0A170QJP8
C	-4	GLY	-	expression tag	UNP A0A170QJP8
C	-3	SER	-	expression tag	UNP A0A170QJP8
C	-2	HIS	-	expression tag	UNP A0A170QJP8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	MET	-	expression tag	UNP A0A170QJP8
C	0	ALA	-	expression tag	UNP A0A170QJP8
C	1	SER	-	expression tag	UNP A0A170QJP8
D	-20	GLY	-	expression tag	UNP A0A170QJP8
D	-19	SER	-	expression tag	UNP A0A170QJP8
D	-18	SER	-	expression tag	UNP A0A170QJP8
D	-17	HIS	-	expression tag	UNP A0A170QJP8
D	-16	HIS	-	expression tag	UNP A0A170QJP8
D	-15	HIS	-	expression tag	UNP A0A170QJP8
D	-14	HIS	-	expression tag	UNP A0A170QJP8
D	-13	HIS	-	expression tag	UNP A0A170QJP8
D	-12	HIS	-	expression tag	UNP A0A170QJP8
D	-11	SER	-	expression tag	UNP A0A170QJP8
D	-10	SER	-	expression tag	UNP A0A170QJP8
D	-9	GLY	-	expression tag	UNP A0A170QJP8
D	-8	LEU	-	expression tag	UNP A0A170QJP8
D	-7	VAL	-	expression tag	UNP A0A170QJP8
D	-6	PRO	-	expression tag	UNP A0A170QJP8
D	-5	ARG	-	expression tag	UNP A0A170QJP8
D	-4	GLY	-	expression tag	UNP A0A170QJP8
D	-3	SER	-	expression tag	UNP A0A170QJP8
D	-2	HIS	-	expression tag	UNP A0A170QJP8
D	-1	MET	-	expression tag	UNP A0A170QJP8
D	0	ALA	-	expression tag	UNP A0A170QJP8
D	1	SER	-	expression tag	UNP A0A170QJP8

- Molecule 2 is 4-oxidanylbenzohydrazide (CCD ID: HDH) (formula: $C_7H_8N_2O_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			11	7	2	2		
2	B	1	Total	C	N	O	0	0
			11	7	2	2		
2	C	1	Total	C	N	O	0	0
			11	7	2	2		
2	D	1	Total	C	N	O	0	0
			11	7	2	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	153	Total	O	0	0
			153	153		
3	B	198	Total	O	0	0
			198	198		
3	C	172	Total	O	0	0
			172	172		
3	D	204	Total	O	0	0
			204	204		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Amidase




- Molecule 1: Amidase




- Molecule 1: Amidase



- Molecule 1: Amidase





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	73.04Å 126.48Å 123.19Å 90.00° 105.66° 90.00°	Depositor
Resolution (Å)	40.60 – 1.80 40.60 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.6 (40.60-1.80) 98.0 (40.60-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.33 (at 1.79Å)	Xtriage
Refinement program	REFMAC 5.8.0155	Depositor
R, R_{free}	0.172 , 0.192 0.183 , 0.202	Depositor DCC
R_{free} test set	9889 reflections (4.91%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.077	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14559	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.97 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.9113e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HDH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/3506	0.79	2/4768 (0.0%)
1	B	0.65	0/3506	0.78	0/4768
1	C	0.65	0/3506	0.80	1/4768 (0.0%)
1	D	0.64	0/3506	0.80	0/4769
All	All	0.65	0/14024	0.79	3/19073 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	82	SER	N-CA-C	5.40	117.92	111.71
1	A	179	SER	N-CA-C	5.15	118.67	112.38
1	C	82	SER	N-CA-C	5.02	117.48	111.71

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3447	0	3428	6	0
1	B	3447	0	3428	9	0
1	C	3447	0	3427	8	0
1	D	3447	0	3421	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	11	0	0	0	0
2	B	11	0	0	0	0
2	C	11	0	0	0	0
2	D	11	0	0	1	0
3	A	153	0	0	1	0
3	B	198	0	0	0	0
3	C	172	0	0	0	0
3	D	204	0	0	1	0
All	All	14559	0	13704	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80[B]:LYS:NZ	1:B:129:CYS:SG	2.37	0.96
1:C:80:LYS:NZ	1:C:129:CYS:SG	2.52	0.82
1:C:208:ARG:O	1:C:210:ARG:NH1	2.13	0.78
1:A:80[B]:LYS:NZ	1:A:129:CYS:SG	2.61	0.74
1:B:355:THR:OG1	1:D:320:ARG:NH1	2.22	0.73
1:B:433:LYS:NZ	1:D:245:THR:O	2.26	0.68
1:B:80[B]:LYS:NZ	1:B:82:SER:OG	2.33	0.62
1:B:260:ARG:HD2	1:B:374:TYR:CE1	2.36	0.60
1:D:246:MET:HE2	1:D:246:MET:HA	1.84	0.59
1:C:208:ARG:C	1:C:210:ARG:HH12	2.10	0.56
1:C:334:ARG:HE	1:C:335:MET:HE3	1.71	0.56
1:C:85:VAL:HG22	1:C:109:THR:HG21	1.89	0.54
1:A:260:ARG:NH2	1:A:295:GLU:OE2	2.36	0.54
1:D:448:ARG:NH2	3:D:602:HOH:O	2.36	0.50
1:D:245:THR:HG23	1:D:246:MET:HE3	1.95	0.47
1:D:185:SER:HB3	1:D:426:GLY:HA2	1.98	0.46
1:B:256:ILE:HD11	1:B:432:ARG:NH2	2.32	0.45
1:C:185:SER:HB3	1:C:426:GLY:HA2	1.98	0.45
1:B:320:ARG:O	1:B:324:GLU:HG3	2.18	0.42
1:C:256:ILE:HD11	1:C:259:PRO:HB3	2.02	0.42
1:A:80[B]:LYS:NZ	1:A:82:SER:OG	2.53	0.41
1:B:216:LEU:HA	1:B:246:MET:HE1	2.02	0.41
1:A:245:THR:HG21	1:C:364:GLN:HG2	2.02	0.41
1:D:129:CYS:O	2:D:501:HDH:N17	2.54	0.41
1:A:99:ASP:HB2	3:A:800:HOH:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:VAL:HG23	1:A:246:MET:HE1	2.03	0.40
1:B:185:SER:HB3	1:B:426:GLY:HA2	2.04	0.40
1:D:125:ILE:N	1:D:125:ILE:HD13	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	456/481 (95%)	440 (96%)	15 (3%)	1 (0%)	43	31
1	B	456/481 (95%)	442 (97%)	13 (3%)	1 (0%)	43	31
1	C	456/481 (95%)	439 (96%)	16 (4%)	1 (0%)	43	31
1	D	456/481 (95%)	440 (96%)	15 (3%)	1 (0%)	43	31
All	All	1824/1924 (95%)	1761 (96%)	59 (3%)	4 (0%)	43	31

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	381	VAL
1	C	381	VAL
1	B	381	VAL
1	D	381	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	358/377 (95%)	356 (99%)	2 (1%)	78	77
1	B	358/377 (95%)	355 (99%)	3 (1%)	73	70
1	C	358/377 (95%)	353 (99%)	5 (1%)	59	52
1	D	358/377 (95%)	356 (99%)	2 (1%)	78	77
All	All	1432/1508 (95%)	1420 (99%)	12 (1%)	73	70

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	82	SER
1	A	125	ILE
1	B	43	ASN
1	B	125	ILE
1	B	208	ARG
1	C	101	LEU
1	C	122	HIS
1	C	125	ILE
1	C	247	SER
1	C	256	ILE
1	D	18	SER
1	D	125	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	29	HIS
1	B	219	GLN
1	B	266	HIS
1	B	394	HIS
1	B	406	GLN
1	C	29	HIS
1	C	219	GLN
1	D	219	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HDH	B	601	-	11,11,11	5.05	2 (18%)	14,14,14	2.17	3 (21%)
2	HDH	A	600	-	11,11,11	4.98	3 (27%)	14,14,14	1.96	2 (14%)
2	HDH	D	501	-	11,11,11	5.63	2 (18%)	14,14,14	1.88	2 (14%)
2	HDH	C	501	-	11,11,11	7.40	3 (27%)	14,14,14	2.94	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HDH	B	601	-	-	6/6/6/6	0/1/1/1
2	HDH	A	600	-	-	4/6/6/6	0/1/1/1
2	HDH	D	501	-	-	6/6/6/6	0/1/1/1
2	HDH	C	501	-	-	4/6/6/6	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	HDH	N17-N9	-23.71	1.07	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	501	HDH	N17-N9	-18.06	1.15	1.41
2	B	601	HDH	N17-N9	-16.07	1.18	1.41
2	A	600	HDH	N17-N9	-15.23	1.19	1.41
2	C	501	HDH	C4-C8	-5.59	1.37	1.50
2	A	600	HDH	C4-C8	-5.17	1.38	1.50
2	D	501	HDH	C4-C8	-4.18	1.41	1.50
2	B	601	HDH	C4-C8	-3.90	1.41	1.50
2	A	600	HDH	C8-N9	3.51	1.37	1.33
2	C	501	HDH	C8-N9	2.36	1.36	1.33

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	501	HDH	C4-C8-N9	-6.40	109.23	116.33
2	C	501	HDH	O10-C8-N9	6.35	130.09	122.46
2	B	601	HDH	C4-C8-N9	5.83	122.80	116.33
2	D	501	HDH	C4-C8-N9	5.78	122.75	116.33
2	A	600	HDH	O10-C8-N9	4.72	128.13	122.46
2	B	601	HDH	O10-C8-N9	-4.56	116.98	122.46
2	A	600	HDH	C4-C8-N9	-4.56	111.27	116.33
2	C	501	HDH	C5-C4-C3	3.74	123.32	118.57
2	D	501	HDH	O10-C8-N9	-3.42	118.36	122.46
2	C	501	HDH	C6-C5-C4	-3.21	117.37	120.80
2	B	601	HDH	C8-N9-N17	2.73	125.79	121.50
2	C	501	HDH	C6-C1-C2	2.16	123.20	119.77

There are no chirality outliers.

All (20) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	601	HDH	C4-C8-N9-N17
2	B	601	HDH	O10-C8-N9-N17
2	D	501	HDH	C4-C8-N9-N17
2	D	501	HDH	O10-C8-N9-N17
2	C	501	HDH	C5-C4-C8-N9
2	C	501	HDH	C5-C4-C8-O10
2	C	501	HDH	C3-C4-C8-O10
2	C	501	HDH	C3-C4-C8-N9
2	A	600	HDH	C5-C4-C8-O10
2	A	600	HDH	C5-C4-C8-N9
2	D	501	HDH	C5-C4-C8-O10
2	D	501	HDH	C3-C4-C8-O10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	600	HDH	C3-C4-C8-O10
2	D	501	HDH	C5-C4-C8-N9
2	D	501	HDH	C3-C4-C8-N9
2	B	601	HDH	C3-C4-C8-O10
2	A	600	HDH	C3-C4-C8-N9
2	B	601	HDH	C5-C4-C8-O10
2	B	601	HDH	C5-C4-C8-N9
2	B	601	HDH	C3-C4-C8-N9

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	501	HDH	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	456/481 (94%)	-0.07	11 (2%) 59 60	6, 16, 29, 54	2 (0%)
1	B	456/481 (94%)	-0.12	6 (1%) 75 75	6, 16, 28, 51	2 (0%)
1	C	457/481 (95%)	-0.03	17 (3%) 45 45	7, 16, 29, 64	1 (0%)
1	D	456/481 (94%)	-0.02	14 (3%) 51 51	7, 17, 29, 60	2 (0%)
All	All	1825/1924 (94%)	-0.06	48 (2%) 57 57	6, 16, 29, 64	7 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	207	GLY	8.4
1	C	208	ARG	6.6
1	D	207	GLY	5.7
1	D	208	ARG	5.7
1	A	208	ARG	4.4
1	A	257	GLU	4.3
1	B	257	GLU	3.7
1	D	398	GLU	3.6
1	B	207	GLY	3.5
1	D	206	GLY	3.4
1	C	207	GLY	3.3
1	B	327	GLU	3.2
1	A	327	GLU	3.2
1	C	460	ILE	3.1
1	B	208	ARG	3.0
1	B	209	SER	2.9
1	D	257	GLU	2.9
1	A	209	SER	2.9
1	D	324	GLU	2.8
1	D	209	SER	2.8
1	C	4	GLU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	206	GLY	2.8
1	D	210	ARG	2.8
1	A	210	ARG	2.7
1	C	64	PHE	2.7
1	D	176	LEU	2.6
1	C	209	SER	2.5
1	C	210	ARG	2.5
1	D	215	GLU	2.5
1	B	365	MET	2.5
1	D	26[A]	GLU	2.4
1	D	452	ARG	2.4
1	C	99	ASP	2.4
1	D	4	GLU	2.3
1	A	206	GLY	2.2
1	A	285	GLU	2.2
1	D	19	GLU	2.2
1	A	36	ASP	2.2
1	C	327	GLU	2.1
1	C	174	GLY	2.1
1	A	99	ASP	2.1
1	A	256	ILE	2.1
1	C	398	GLU	2.1
1	C	452	ARG	2.1
1	C	370	LEU	2.0
1	C	18	SER	2.0
1	C	256	ILE	2.0
1	C	323	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	HDH	C	501	11/11	0.76	0.15	23,24,30,31	0
2	HDH	B	601	11/11	0.82	0.12	22,23,26,28	0
2	HDH	D	501	11/11	0.82	0.13	23,25,29,31	0
2	HDH	A	600	11/11	0.85	0.12	22,23,27,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.