



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 12:28 AM UTC

PDB ID : 8H7H / pdb_00008h7h
Title : The crystal structure of human abl1 kinase domain in complex with abl1-A-EBA
Authors : Zhu, C.J.; Zhang, Z.M.
Deposited on : 2022-10-20
Resolution : 2.28 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.28 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	9078 (2.30-2.26)
Clashscore	190562	9802 (2.30-2.26)
Ramachandran outliers	187476	9690 (2.30-2.26)
Sidechain outliers	187428	9691 (2.30-2.26)
RSRZ outliers	180081	9085 (2.30-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	5% 89% 7% • •
1	B	277	15% 86% 6% • 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4737 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Tyrosine-protein kinase ABL1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	P	S	0	0	0
			2164	1394	351	401	1	17			
1	B	258	Total	C	N	O	P	S	0	0	0
			2031	1306	328	381	1	15			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	224	GLY	-	expression tag	UNP P00519
A	225	ALA	-	expression tag	UNP P00519
A	226	MET	-	expression tag	UNP P00519
A	227	GLY	-	expression tag	UNP P00519
A	228	SER	-	expression tag	UNP P00519
B	224	GLY	-	expression tag	UNP P00519
B	225	ALA	-	expression tag	UNP P00519
B	226	MET	-	expression tag	UNP P00519
B	227	GLY	-	expression tag	UNP P00519
B	228	SER	-	expression tag	UNP P00519

- Molecule 2 is 5-[3-(6-methoxyisoquinolin-7-yl)-1H-pyrrolo[2,3-b]pyridin-5-yl]-N-methyl-N-prop-2-ynyl-pyridine-3-carboxamide (CCD ID: QH9) (formula: C₂₇H₂₁N₅O₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			34	27	5	2		
2	B	1	Total	C	N	O	0	0
			34	27	5	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	276	Total	O	0	0
			276	276		
3	B	198	Total	O	0	0
			198	198		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Tyrosine-protein kinase ABL1



• Molecule 1: Tyrosine-protein kinase ABL1



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 ₁ 2 ₁ 2 ₁	Depositor
Cell constants a, b, c, α , β , γ	56.77Å 104.78Å 133.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.19 – 2.28 26.19 – 2.28	Depositor EDS
% Data completeness (in resolution range)	94.2 (26.19-2.28) 94.1 (26.19-2.28)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.9_1692+SVN	Depositor
R, R_{free}	0.182 , 0.221 0.185 , 0.218	Depositor DCC
R_{free} test set	2000 reflections (5.39%)	wwPDB-VP
Wilson B-factor (Å ²)	9.5	Xtriage
Anisotropy	1.445	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 56.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4737	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.39% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: QH9, PTR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.29	0/2202	0.67	0/2983
1	B	0.28	0/2061	0.70	1/2796 (0.0%)
All	All	0.29	0/4263	0.68	1/5779 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	264	TYR	N-CA-C	-6.97	99.67	108.45

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2164	0	2088	17	0
1	B	2031	0	1916	12	0
2	A	34	0	0	0	0
2	B	34	0	0	1	0
3	A	276	0	0	4	0
3	B	198	0	0	5	0
All	All	4737	0	4004	29	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:399:ALA:O	3:B:701:HOH:O	1.83	0.97
2:B:601:QH9:O1	3:B:702:HOH:O	1.93	0.85
1:A:393:PTR:O1P	3:A:701:HOH:O	1.94	0.84
1:A:282:GLU:OE1	3:A:702:HOH:O	2.05	0.74
1:B:332:ARG:O	3:B:704:HOH:O	2.09	0.70
1:A:234:LYS:HD2	1:A:234:LYS:H	1.57	0.69
1:B:305:CYS:HB2	1:B:312:TYR:HB2	1.77	0.66
1:B:262:LYS:C	1:B:264:TYR:H	2.02	0.65
1:A:234:LYS:HD2	1:A:234:LYS:N	2.12	0.64
1:B:257:TYR:OH	3:B:703:HOH:O	2.06	0.63
1:A:249:GLY:HA3	1:A:251:GLY:H	1.72	0.54
1:B:282:GLU:HG3	1:B:283:PHE:N	2.23	0.52
1:A:294:LYS:NZ	3:A:711:HOH:O	2.40	0.52
1:B:356:LYS:HG3	3:B:753:HOH:O	2.10	0.50
1:A:249:GLY:HA3	1:A:251:GLY:N	2.27	0.49
1:A:386:ARG:NH1	1:B:498:GLN:O	2.47	0.47
1:A:334:GLU:OE1	3:A:705:HOH:O	2.20	0.45
1:A:278:MET:HE2	1:A:282:GLU:HB3	1.99	0.44
1:B:232:TYR:O	1:B:236:GLU:HG2	2.17	0.44
1:A:307:ARG:HE	1:A:307:ARG:HB3	1.64	0.44
1:B:262:LYS:C	1:B:264:TYR:N	2.72	0.43
1:A:248:LEU:HD21	1:A:258:GLU:HB2	2.01	0.43
1:A:292:GLU:OE2	1:A:386:ARG:NH2	2.52	0.43
1:B:386:ARG:HD2	1:B:388:MET:SD	2.58	0.42
1:A:320:TYR:HE2	1:A:373:GLU:HA	1.84	0.42
1:B:315:THR:HG21	1:B:382:PHE:HE1	1.85	0.41
1:A:298:LEU:HD23	1:A:379:VAL:HB	2.02	0.41
1:A:400:LYS:HA	1:A:400:LYS:HD3	1.83	0.40
1:A:493:PHE:HA	1:A:496:MET:HE3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	266/277 (96%)	258 (97%)	8 (3%)	0	100	100
1	B	251/277 (91%)	241 (96%)	10 (4%)	0	100	100
All	All	517/554 (93%)	499 (96%)	18 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	225/239 (94%)	222 (99%)	3 (1%)	61	76
1	B	205/239 (86%)	202 (98%)	3 (2%)	57	72
All	All	430/478 (90%)	424 (99%)	6 (1%)	59	74

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	294	LYS
1	A	356	LYS
1	A	462	GLU
1	B	241	ASP
1	B	242	ILE
1	B	282	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such

sidechains are listed below:

Mol	Chain	Res	Type
1	A	297	ASN
1	A	477	GLN
1	A	491	GLN
1	B	477	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	PTR	A	393	1	15,16,17	1.21	1 (6%)	17,22,24	0.55	0
1	PTR	B	393	1	15,16,17	1.22	1 (6%)	17,22,24	0.62	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	A	393	1	-	1/10/11/13	0/1/1/1
1	PTR	B	393	1	-	0/10/11/13	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	393	PTR	OH-CZ	-4.39	1.30	1.40
1	A	393	PTR	OH-CZ	-4.35	1.30	1.40

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	393	PTR	O2P-P-OH	2.04	111.36	105.32

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	393	PTR	C-CA-CB-CG

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	393	PTR	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	QH9	A	601	1	38,38,38	2.47	15 (39%)	51,54,54	2.17	20 (39%)
2	QH9	B	601	1	38,38,38	2.54	16 (42%)	51,54,54	1.99	17 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	QH9	A	601	1	-	4/20/21/21	0/5/5/5
2	QH9	B	601	1	-	0/20/21/21	0/5/5/5

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	QH9	C22-N5	6.56	1.45	1.34
2	B	601	QH9	C22-N5	6.49	1.45	1.34
2	B	601	QH9	C13-C10	5.04	1.42	1.37
2	B	601	QH9	C23-C25	5.00	1.54	1.46
2	B	601	QH9	C14-C15	4.93	1.48	1.39
2	A	601	QH9	C13-C10	4.93	1.42	1.37
2	A	601	QH9	C23-C25	4.92	1.54	1.46
2	A	601	QH9	C14-C15	4.81	1.48	1.39
2	B	601	QH9	C14-C11	4.31	1.46	1.39
2	A	601	QH9	C14-C11	4.10	1.46	1.39
2	B	601	QH9	C18-C17	3.87	1.46	1.39
2	A	601	QH9	C18-C17	3.76	1.46	1.39
2	B	601	QH9	C12-N3	3.73	1.40	1.35
2	A	601	QH9	C12-N3	3.43	1.40	1.35
2	B	601	QH9	C8-C10	3.42	1.55	1.49
2	A	601	QH9	C8-C10	3.14	1.54	1.49
2	B	601	QH9	C2-C3	3.09	1.48	1.42
2	A	601	QH9	C2-C3	3.05	1.48	1.42
2	B	601	QH9	C17-C15	2.97	1.56	1.49
2	B	601	QH9	C6-C7	2.89	1.41	1.36
2	A	601	QH9	C6-C7	2.82	1.41	1.36
2	A	601	QH9	C17-C15	2.74	1.55	1.49
2	A	601	QH9	C8-C7	2.72	1.45	1.40
2	B	601	QH9	C8-C7	2.71	1.45	1.40
2	B	601	QH9	C2-C1	2.30	1.39	1.36
2	B	601	QH9	C5-N1	2.29	1.36	1.32
2	A	601	QH9	C19-C22	2.25	1.53	1.50
2	A	601	QH9	C2-C1	2.24	1.39	1.36
2	A	601	QH9	C5-N1	2.17	1.36	1.32
2	B	601	QH9	C16-C15	2.06	1.43	1.39
2	B	601	QH9	C20-C19	2.01	1.42	1.39

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	QH9	C12-C11-C10	-5.03	104.48	106.22
2	A	601	QH9	O2-C7-C6	-4.49	119.47	125.16

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	QH9	C12-C11-C10	-4.34	104.72	106.22
2	B	601	QH9	C12-N2-C13	4.32	110.52	108.48
2	B	601	QH9	O2-C7-C6	-4.09	119.98	125.16
2	A	601	QH9	C11-C12-N2	4.06	109.91	108.42
2	B	601	QH9	C21-N4-C20	3.70	122.54	117.51
2	A	601	QH9	C19-C22-N5	3.69	125.13	118.92
2	A	601	QH9	C12-N2-C13	3.57	110.16	108.48
2	B	601	QH9	C11-C12-N3	-3.47	122.10	126.31
2	A	601	QH9	C21-N4-C20	3.43	122.17	117.51
2	A	601	QH9	C11-C12-N3	-3.40	122.20	126.31
2	B	601	QH9	C16-N3-C12	3.29	122.63	116.19
2	A	601	QH9	C14-C15-C16	3.22	120.56	117.10
2	B	601	QH9	C11-C12-N2	3.20	109.59	108.42
2	B	601	QH9	C10-C13-N2	-3.12	108.45	110.46
2	A	601	QH9	C17-C18-C19	-3.10	117.72	121.04
2	B	601	QH9	C14-C11-C10	3.06	137.53	131.82
2	A	601	QH9	C16-N3-C12	3.03	122.12	116.19
2	A	601	QH9	O2-C7-C8	3.01	120.92	116.55
2	B	601	QH9	C15-C16-N3	-3.00	119.51	124.28
2	A	601	QH9	C10-C13-N2	-2.99	108.53	110.46
2	A	601	QH9	C16-C15-C17	-2.92	116.69	121.62
2	A	601	QH9	C14-C11-C10	2.87	137.19	131.82
2	B	601	QH9	C17-C18-C19	-2.84	118.00	121.04
2	A	601	QH9	C15-C16-N3	-2.82	119.80	124.28
2	B	601	QH9	C14-C15-C16	2.70	120.00	117.10
2	B	601	QH9	O2-C7-C8	2.58	120.30	116.55
2	A	601	QH9	C18-C17-C21	2.47	119.75	117.10
2	A	601	QH9	C27-O2-C7	-2.39	114.00	117.51
2	A	601	QH9	O1-C22-N5	-2.24	118.44	122.21
2	B	601	QH9	C18-C17-C21	2.23	119.49	117.10
2	A	601	QH9	C1-N1-C5	2.18	121.55	117.19
2	B	601	QH9	C1-N1-C5	2.13	121.46	117.19
2	B	601	QH9	C19-C22-N5	2.12	122.50	118.92
2	A	601	QH9	C11-C14-C15	-2.07	116.98	120.82
2	B	601	QH9	O1-C22-N5	-2.03	118.79	122.21

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	QH9	C11-C10-C8-C9
2	A	601	QH9	C25-C23-N5-C24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	601	QH9	C6-C7-O2-C27
2	A	601	QH9	C8-C7-O2-C27


There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	QH9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand QH9 A 601





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/277 (96%)	-0.04	15 (5%) 30 31	5, 17, 45, 58	0
1	B	257/277 (92%)	0.64	42 (16%) 4 5	10, 25, 73, 83	0
All	All	525/554 (94%)	0.29	57 (10%) 10 11	5, 21, 65, 83	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	242	ILE	6.9
1	B	312	TYR	6.4
1	B	232	TYR	5.3
1	B	397	ALA	5.1
1	B	262	LYS	4.9
1	A	232	TYR	4.8
1	B	264	TYR	4.7
1	B	240	THR	4.6
1	B	233	ASP	4.5
1	B	282	GLU	4.3
1	B	260	VAL	4.2
1	A	462	GLU	4.2
1	B	257	TYR	4.2
1	B	261	TRP	4.2
1	A	249	GLY	4.1
1	B	238	GLU	4.1
1	B	244	MET	4.0
1	A	274	LYS	3.9
1	B	265	SER	3.8
1	A	389	THR	3.8
1	B	396	HIS	3.8
1	B	273	LEU	3.7
1	B	241	ASP	3.7
1	B	266	LEU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	245	LYS	3.4
1	A	277	THR	3.3
1	B	279	GLU	3.3
1	A	275	GLU	3.3
1	B	246	HIS	3.2
1	B	243	THR	3.2
1	A	251	GLY	3.2
1	B	280	VAL	3.2
1	B	272	THR	3.2
1	B	444	ASP	3.1
1	B	248	LEU	2.9
1	B	239	ARG	2.8
1	B	237	MET	2.7
1	A	233	ASP	2.6
1	A	252	GLN	2.5
1	B	263	LYS	2.4
1	B	251	GLY	2.4
1	A	248	LEU	2.4
1	B	235	TRP	2.4
1	A	250	GLY	2.3
1	B	255	GLU	2.3
1	B	303	GLY	2.3
1	B	462	GLU	2.2
1	B	305	CYS	2.2
1	A	254	GLY	2.2
1	B	390	GLY	2.2
1	A	500	SER	2.2
1	B	256	VAL	2.1
1	B	281	GLU	2.1
1	B	259	GLY	2.1
1	B	267	THR	2.0
1	A	234	LYS	2.0
1	B	236	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	PTR	A	393	16/17	0.93	0.10	10,18,50,56	0
1	PTR	B	393	16/17	0.93	0.10	22,25,60,60	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	QH9	B	601	34/34	0.90	0.12	11,27,55,57	0
2	QH9	A	601	34/34	0.93	0.11	6,14,52,54	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around QH9 B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around QH9 A 601:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.