



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:27 AM UTC

PDB ID : 1H8X / pdb_00001h8x
Title : Domain-swapped Dimer of a Human Pancreatic Ribonuclease Variant
Authors : Canals, A.; Pous, J.; Guasch, A.; Benito, A.; Ribo, M.; Vilanova, M.; Coll, M.
Deposited on : 2001-02-16
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	128	
1	B	128	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2327 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	126	Total	C	N	O	S	18	3	1
			998	601	194	190	13			
1	B	126	Total	C	N	O	S	54	3	1
			989	595	189	192	13			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	104	ALA	ARG	engineered mutation	UNP P07998
B	104	ALA	ARG	engineered mutation	UNP P07998
A	106	ALA	LYS	engineered mutation	UNP P07998
B	106	ALA	LYS	engineered mutation	UNP P07998
A	109	GLU	GLN	engineered mutation	UNP P07998
B	109	GLU	GLN	engineered mutation	UNP P07998
A	116	GLY	ASP	engineered mutation	UNP P07998
B	116	GLY	ASP	engineered mutation	UNP P07998
A	117	ASN	SER	engineered mutation	UNP P07998
B	117	ASN	SER	engineered mutation	UNP P07998
A	201	GLN	PRO	engineered mutation	UNP P07998
B	201	GLN	PRO	engineered mutation	UNP P07998

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	162	Total	O	0	0
			162	162		
2	B	178	Total	O	0	0
			178	178		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: RIBONUCLEASE 1

Chain A:  80% 15% ...



- Molecule 1: RIBONUCLEASE 1

Chain B:  74% 20% ...



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.61Å 61.43Å 75.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	12.00 – 2.00	Depositor
% Data completeness (in resolution range)	99.6 (12.00-2.00)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.196 , 0.243	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2327	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.47	1/1032 (0.1%)	1.15	12/1387 (0.9%)
1	B	0.46	1/1024 (0.1%)	1.02	8/1379 (0.6%)
All	All	0.46	2/2056 (0.1%)	1.09	20/2766 (0.7%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	225	GLU	C-N	-5.81	1.25	1.33
1	B	225	GLU	C-N	-5.71	1.25	1.33

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	191[A]	ARG	CG-CD-NE	-10.57	88.73	112.00
1	A	191[B]	ARG	CG-CD-NE	-10.57	88.73	112.00
1	B	192	TYR	N-CA-C	-8.78	97.24	109.71
1	B	205	HIS	N-CA-C	-7.05	100.04	110.48
1	B	223	SER	N-CA-C	-6.77	99.18	109.95
1	A	187	THR	N-CA-C	-6.75	101.44	110.55
1	A	174[A]	LYS	CG-CD-CE	6.74	126.81	111.30
1	A	174[B]	LYS	CG-CD-CE	6.74	126.81	111.30
1	A	161	GLU	N-CA-C	6.65	118.73	109.15
1	A	141	LYS	CA-C-N	6.62	125.86	118.97
1	A	141	LYS	C-N-CA	6.62	125.86	118.97
1	A	115	SER	N-CA-C	6.58	118.45	111.28
1	A	112	HIS	N-CA-C	5.99	121.51	113.72
1	A	205	HIS	N-CA-C	-5.88	101.78	110.48
1	B	193	PRO	N-CA-C	5.66	126.80	112.10
1	B	112	HIS	N-CA-C	5.58	121.87	114.12
1	B	189	GLY	N-CA-C	5.54	126.30	113.18
1	B	141	LYS	CA-C-N	5.11	126.22	119.84
1	B	141	LYS	C-N-CA	5.11	126.22	119.84
1	A	134	ASN	N-CA-C	5.09	118.78	112.58

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	998	0	955	21	0
1	B	989	0	933	25	0
2	A	162	0	0	1	0
2	B	178	0	0	9	0
All	All	2327	0	1888	45	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (45) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:191[B]:ARG:CD	1:A:191[B]:ARG:CG	1.76	1.60
1:A:191[B]:ARG:CG	1:A:191[B]:ARG:NE	2.26	0.98
1:A:174[A]:LYS:HD2	1:A:224:VAL:HG21	1.63	0.80
1:A:191[A]:ARG:HH21	1:A:191[A]:ARG:HB3	1.54	0.73
1:A:162:LYS:NZ	1:A:171:ASN:HD22	1.89	0.71
1:B:189:GLY:O	2:B:2140:HOH:O	2.10	0.68
1:A:162:LYS:HZ3	1:A:171:ASN:HD22	1.42	0.66
1:B:167:ASN:HB3	2:B:2075:HOH:O	1.97	0.62
1:B:191:ARG:O	1:B:192:TYR:C	2.42	0.62
1:A:160:GLN:HG2	1:A:176:ASN:ND2	2.16	0.61
1:B:162:LYS:NZ	1:B:171:ASN:HD22	1.99	0.61
1:B:166:LYS:HE3	2:B:2086:HOH:O	2.02	0.60
1:A:191[B]:ARG:CG	1:A:191[B]:ARG:HE	2.16	0.57
1:A:191[A]:ARG:HB3	1:A:191[A]:ARG:NH2	2.18	0.56
1:B:209:ALA:HB3	1:B:219[B]:HIS:HB3	1.89	0.54
1:B:131:ARG:NH2	2:B:2038:HOH:O	2.41	0.54
1:A:174[A]:LYS:HE3	2:A:2113:HOH:O	2.07	0.54
1:B:209:ALA:HB3	1:B:219[A]:HIS:HB2	1.89	0.54
1:B:224:VAL:O	1:B:225:GLU:HB2	2.08	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:162:LYS:HZ1	1:B:171:ASN:HD22	1.55	0.53
1:B:139:ARG:HA	1:B:192:TYR:CD1	2.44	0.52
1:B:167:ASN:CB	2:B:2075:HOH:O	2.57	0.50
1:A:191[B]:ARG:CD	1:A:191[B]:ARG:HB2	2.39	0.49
1:B:221:ASP:O	1:B:222:ALA:HB2	2.14	0.47
1:B:187:THR:HB	1:B:198:ARG:HG3	1.96	0.47
1:A:163:VAL:HG11	1:A:174[B]:LYS:HE2	1.97	0.47
1:A:191[B]:ARG:NE	1:A:191[B]:ARG:HG2	2.23	0.46
1:A:133:ARG:O	1:A:134:ASN:HB2	2.15	0.45
1:B:186:LEU:HD12	1:B:196:ALA:O	2.16	0.45
1:B:174:LYS:HE3	1:B:224:VAL:HG21	1.98	0.45
1:A:148:HIS:ND1	1:B:117:ASN:O	2.50	0.44
1:A:161:GLU:HG2	1:A:163:VAL:HG13	2.00	0.43
1:B:152:VAL:HG13	2:B:2079:HOH:O	2.19	0.43
1:A:192:TYR:CD1	1:A:193:PRO:HA	2.54	0.43
1:A:161:GLU:OE1	1:A:174[A]:LYS:HE2	2.20	0.42
1:B:205:HIS:HE1	2:B:2157:HOH:O	2.03	0.42
1:B:218:VAL:O	1:B:219[A]:HIS:ND1	2.53	0.41
1:B:166:LYS:NZ	2:B:2110:HOH:O	2.54	0.41
1:B:171:ASN:O	1:B:209:ALA:HA	2.20	0.41
1:A:203:GLU:C	1:A:204:ARG:HG2	2.46	0.41
1:B:189:GLY:N	2:B:2141:HOH:O	2.54	0.41
1:B:191:ARG:O	1:B:193:PRO:N	2.54	0.40
1:B:216:VAL:HB	1:B:217:PRO:CD	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	127/128 (99%)	122 (96%)	5 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	127/128 (99%)	113 (89%)	13 (10%)	1 (1%)	16	11
All	All	254/256 (99%)	235 (92%)	18 (7%)	1 (0%)	30	27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	193	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	116/116 (100%)	113 (97%)	3 (3%)	40	44
1	B	116/116 (100%)	115 (99%)	1 (1%)	70	78
All	All	232/232 (100%)	228 (98%)	4 (2%)	61	60

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	188	ASN
1	A	191[A]	ARG
1	A	191[B]	ARG
1	B	194	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	111	GLN
1	A	155	GLN
1	A	171	ASN
1	A	205	HIS
1	B	111	GLN
1	B	171	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	176	ASN
1	B	205	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.