



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 07:32 AM UTC

PDB ID : 4H8S / pdb_00004h8s
Title : Crystal structure of human APPL2BARPH domain
Authors : Martin, J.L.; King, G.J.
Deposited on : 2012-09-23
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1085 (3.54-3.46)
Clashscore	190562	1140 (3.54-3.46)
Ramachandran outliers	187476	1113 (3.54-3.46)
Sidechain outliers	187428	1114 (3.54-3.46)
RSRZ outliers	180081	1084 (3.54-3.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	407	
1	B	407	
1	C	407	
1	D	407	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 23866 atoms, of which 11908 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DCC-interacting protein 13-beta.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	378	Total	C	H	N	O	S	0	0	0
			5982	1880	2982	514	585	21			
1	B	374	Total	C	H	N	O	S	0	0	0
			5926	1859	2959	509	578	21			
1	C	382	Total	C	H	N	O	S	0	0	0
			6053	1905	3017	519	591	21			
1	D	373	Total	C	H	N	O	S	0	0	0
			5905	1850	2950	508	576	21			

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	MET	-	expression tag	UNP Q8NEU8
A	-21	HIS	-	expression tag	UNP Q8NEU8
A	-20	HIS	-	expression tag	UNP Q8NEU8
A	-19	HIS	-	expression tag	UNP Q8NEU8
A	-18	HIS	-	expression tag	UNP Q8NEU8
A	-17	HIS	-	expression tag	UNP Q8NEU8
A	-16	HIS	-	expression tag	UNP Q8NEU8
A	-15	SER	-	expression tag	UNP Q8NEU8
A	-14	SER	-	expression tag	UNP Q8NEU8
A	-13	GLY	-	expression tag	UNP Q8NEU8
A	-12	VAL	-	expression tag	UNP Q8NEU8
A	-11	ASP	-	expression tag	UNP Q8NEU8
A	-10	LEU	-	expression tag	UNP Q8NEU8
A	-9	GLY	-	expression tag	UNP Q8NEU8
A	-8	THR	-	expression tag	UNP Q8NEU8
A	-7	GLU	-	expression tag	UNP Q8NEU8
A	-6	ASN	-	expression tag	UNP Q8NEU8
A	-5	LEU	-	expression tag	UNP Q8NEU8
A	-4	TYR	-	expression tag	UNP Q8NEU8
A	-3	PHE	-	expression tag	UNP Q8NEU8
A	-2	GLN	-	expression tag	UNP Q8NEU8

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	expression tag	UNP Q8NEU8
A	0	ASN	-	expression tag	UNP Q8NEU8
A	1	ALA	-	expression tag	UNP Q8NEU8
B	-22	MET	-	expression tag	UNP Q8NEU8
B	-21	HIS	-	expression tag	UNP Q8NEU8
B	-20	HIS	-	expression tag	UNP Q8NEU8
B	-19	HIS	-	expression tag	UNP Q8NEU8
B	-18	HIS	-	expression tag	UNP Q8NEU8
B	-17	HIS	-	expression tag	UNP Q8NEU8
B	-16	HIS	-	expression tag	UNP Q8NEU8
B	-15	SER	-	expression tag	UNP Q8NEU8
B	-14	SER	-	expression tag	UNP Q8NEU8
B	-13	GLY	-	expression tag	UNP Q8NEU8
B	-12	VAL	-	expression tag	UNP Q8NEU8
B	-11	ASP	-	expression tag	UNP Q8NEU8
B	-10	LEU	-	expression tag	UNP Q8NEU8
B	-9	GLY	-	expression tag	UNP Q8NEU8
B	-8	THR	-	expression tag	UNP Q8NEU8
B	-7	GLU	-	expression tag	UNP Q8NEU8
B	-6	ASN	-	expression tag	UNP Q8NEU8
B	-5	LEU	-	expression tag	UNP Q8NEU8
B	-4	TYR	-	expression tag	UNP Q8NEU8
B	-3	PHE	-	expression tag	UNP Q8NEU8
B	-2	GLN	-	expression tag	UNP Q8NEU8
B	-1	SER	-	expression tag	UNP Q8NEU8
B	0	ASN	-	expression tag	UNP Q8NEU8
B	1	ALA	-	expression tag	UNP Q8NEU8
C	-22	MET	-	expression tag	UNP Q8NEU8
C	-21	HIS	-	expression tag	UNP Q8NEU8
C	-20	HIS	-	expression tag	UNP Q8NEU8
C	-19	HIS	-	expression tag	UNP Q8NEU8
C	-18	HIS	-	expression tag	UNP Q8NEU8
C	-17	HIS	-	expression tag	UNP Q8NEU8
C	-16	HIS	-	expression tag	UNP Q8NEU8
C	-15	SER	-	expression tag	UNP Q8NEU8
C	-14	SER	-	expression tag	UNP Q8NEU8
C	-13	GLY	-	expression tag	UNP Q8NEU8
C	-12	VAL	-	expression tag	UNP Q8NEU8
C	-11	ASP	-	expression tag	UNP Q8NEU8
C	-10	LEU	-	expression tag	UNP Q8NEU8
C	-9	GLY	-	expression tag	UNP Q8NEU8
C	-8	THR	-	expression tag	UNP Q8NEU8



Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-7	GLU	-	expression tag	UNP Q8NEU8
C	-6	ASN	-	expression tag	UNP Q8NEU8
C	-5	LEU	-	expression tag	UNP Q8NEU8
C	-4	TYR	-	expression tag	UNP Q8NEU8
C	-3	PHE	-	expression tag	UNP Q8NEU8
C	-2	GLN	-	expression tag	UNP Q8NEU8
C	-1	SER	-	expression tag	UNP Q8NEU8
C	0	ASN	-	expression tag	UNP Q8NEU8
C	1	ALA	-	expression tag	UNP Q8NEU8
D	-22	MET	-	expression tag	UNP Q8NEU8
D	-21	HIS	-	expression tag	UNP Q8NEU8
D	-20	HIS	-	expression tag	UNP Q8NEU8
D	-19	HIS	-	expression tag	UNP Q8NEU8
D	-18	HIS	-	expression tag	UNP Q8NEU8
D	-17	HIS	-	expression tag	UNP Q8NEU8
D	-16	HIS	-	expression tag	UNP Q8NEU8
D	-15	SER	-	expression tag	UNP Q8NEU8
D	-14	SER	-	expression tag	UNP Q8NEU8
D	-13	GLY	-	expression tag	UNP Q8NEU8
D	-12	VAL	-	expression tag	UNP Q8NEU8
D	-11	ASP	-	expression tag	UNP Q8NEU8
D	-10	LEU	-	expression tag	UNP Q8NEU8
D	-9	GLY	-	expression tag	UNP Q8NEU8
D	-8	THR	-	expression tag	UNP Q8NEU8
D	-7	GLU	-	expression tag	UNP Q8NEU8
D	-6	ASN	-	expression tag	UNP Q8NEU8
D	-5	LEU	-	expression tag	UNP Q8NEU8
D	-4	TYR	-	expression tag	UNP Q8NEU8
D	-3	PHE	-	expression tag	UNP Q8NEU8
D	-2	GLN	-	expression tag	UNP Q8NEU8
D	-1	SER	-	expression tag	UNP Q8NEU8
D	0	ASN	-	expression tag	UNP Q8NEU8
D	1	ALA	-	expression tag	UNP Q8NEU8

- Molecule 1: DCC-interacting protein 13-beta





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	97.96Å 208.01Å 218.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.95 – 3.50 46.95 – 3.50	Depositor EDS
% Data completeness (in resolution range)	95.4 (46.95-3.50) 94.4 (46.95-3.50)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 3.48Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1031)	Depositor
R, R_{free}	0.209 , 0.256 0.219 , 0.260	Depositor DCC
R_{free} test set	2810 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	97.7	Xtriage
Anisotropy	0.357	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 94.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.025 for -h,l,k	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	23866	wwPDB-VP
Average B, all atoms (Å ²)	119.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.39	0/3043	0.77	5/4100 (0.1%)
1	B	0.37	0/3009	0.75	1/4052 (0.0%)
1	C	0.37	0/3080	0.71	0/4151
1	D	0.37	0/2996	0.73	0/4034
All	All	0.38	0/12128	0.74	6/16337 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	323	GLN	N-CA-C	6.64	119.37	111.33
1	A	282	GLY	N-CA-C	6.39	120.43	111.10
1	A	346	THR	CA-C-N	-5.96	113.54	119.56
1	A	346	THR	C-N-CA	-5.96	113.54	119.56
1	A	280	LYS	CA-C-N	5.12	130.91	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	280	LYS	Peptide
1	B	12	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3000	2982	2990	17	0
1	B	2967	2959	2964	17	0
1	C	3036	3017	3027	22	0
1	D	2955	2950	2955	19	0
All	All	11958	11908	11936	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 69 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:372:ASN:OD1	1:C:375:ARG:NH2	2.19	0.76
1:C:8:LEU:CB	1:C:9:LEU:HA	2.17	0.75
1:A:280:LYS:HA	1:A:281:ALA:CB	2.27	0.65
1:A:324:ASP:HB3	1:A:325:LEU:HA	1.77	0.64
1:C:8:LEU:HB2	1:C:9:LEU:HA	1.80	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	374/407 (92%)	346 (92%)	23 (6%)	5 (1%)	9	39
1	B	372/407 (91%)	344 (92%)	24 (6%)	4 (1%)	11	43

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	378/407 (93%)	346 (92%)	24 (6%)	8 (2%)	5	31
1	D	371/407 (91%)	336 (91%)	29 (8%)	6 (2%)	7	36
All	All	1495/1628 (92%)	1372 (92%)	100 (7%)	23 (2%)	8	37

5 of 23 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	281	ALA
1	C	5	ASP
1	C	22	LEU
1	C	293	VAL
1	C	325	LEU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/355 (93%)	322 (98%)	8 (2%)	43	64
1	B	326/355 (92%)	314 (96%)	12 (4%)	30	56
1	C	334/355 (94%)	319 (96%)	15 (4%)	24	50
1	D	325/355 (92%)	313 (96%)	12 (4%)	30	56
All	All	1315/1420 (93%)	1268 (96%)	47 (4%)	31	57

5 of 47 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	317	VAL
1	D	10	GLU
1	C	323	GLN
1	C	346	THR
1	D	102	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 14 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	363	ASN
1	D	72	ASN
1	D	312	GLN
1	D	276	ASN
1	D	308	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	378/407 (92%)	0.18	14 (3%)	45 25	62, 109, 162, 303	0
1	B	374/407 (91%)	0.16	18 (4%)	35 19	71, 114, 191, 287	0
1	C	382/407 (93%)	0.20	18 (4%)	36 20	71, 116, 172, 267	0
1	D	373/407 (91%)	0.22	19 (5%)	33 19	55, 116, 189, 272	0
All	All	1507/1628 (92%)	0.19	69 (4%)	37 20	55, 114, 179, 303	0

The worst 5 of 69 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	265	ASP	6.6
1	A	377	ILE	4.8
1	B	14	GLN	4.5
1	C	378	TYR	4.4
1	D	265	ASP	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.