



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:25 PM UTC

PDB ID : 5H8V / pdb_00005h8v
Title : Crystal structure of the complex between maize Sulfite Reductase and ferredoxin in the form-1 crystal
Authors : Kurisu, G.; Nakayama, M.; Hase, T.
Deposited on : 2015-12-24
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	583	
1	B	583	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SRM	A	702	X	-	-	-
3	SRM	B	702	X	-	-	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 9287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sulfite reductase [ferredoxin], chloroplastic.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	564	Total	C	N	O	S	0	0	0
			4455	2830	772	836	17			
1	B	560	Total	C	N	O	S	0	0	0
			4421	2808	767	830	16			

- Molecule 2 is IRON/SULFUR CLUSTER (CCD ID: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is SIROHEME (CCD ID: SRM) (formula: C₄₂H₄₄FeN₄O₁₆).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 63	C 42	Fe 1	N 4	O 16	0	0
3	B	1	Total 63	C 42	Fe 1	N 4	O 16	0	0

- Molecule 4 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total 5	O 4	P 1	0	0
4	B	1	Total 5	O 4	P 1	0	0

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total 2	Mg 2	0	0
5	B	2	Total 2	Mg 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	115	Total 115	O 115	0	0
6	B	140	Total 140	O 140	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sulfite reductase [ferredoxin], chloroplastic



- Molecule 1: Sulfite reductase [ferredoxin], chloroplastic



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.35Å 103.35Å 255.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.04 – 2.20 48.04 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.4 (48.04-2.20) 99.4 (48.04-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.23 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.200 , 0.239 0.205 , 0.239	Depositor DCC
R_{free} test set	2129 reflections (3.00%)	wwPDB-VP
Wilson B-factor (Å ²)	24.3	Xtriage
Anisotropy	0.062	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 25.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9287	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.3896e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, MG, SRM, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.11	3/4550 (0.1%)	1.02	1/6162 (0.0%)
1	B	1.13	6/4516 (0.1%)	1.03	4/6118 (0.1%)
All	All	1.12	9/9066 (0.1%)	1.02	5/12280 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	235	GLY	N-CA	6.17	1.55	1.45
1	A	575	LEU	C-O	-6.10	1.16	1.24
1	B	575	LEU	C-O	-6.03	1.16	1.24
1	A	162	LEU	N-CA	-5.69	1.39	1.46
1	B	546	ARG	C-O	5.62	1.26	1.23
1	A	396	PHE	C-O	-5.38	1.17	1.23
1	B	360	ARG	C-O	5.37	1.30	1.23
1	B	212	GLN	CA-CB	5.33	1.61	1.53
1	B	606	GLY	C-O	-5.13	1.17	1.23

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	160	PHE	CB-CA-C	-5.79	99.75	109.48
1	B	500	CYS	CA-C-N	-5.78	113.86	119.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	500	CYS	C-N-CA	-5.78	113.86	119.87
1	B	160	PHE	CB-CA-C	-5.60	100.08	109.48
1	B	193	ARG	NE-CZ-NH1	-5.45	116.05	121.50

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	234	ASP	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4455	0	4448	31	0
1	B	4421	0	4409	29	0
2	A	8	0	0	0	0
2	B	8	0	0	0	0
3	A	63	0	34	5	0
3	B	63	0	34	7	0
4	A	5	0	0	1	0
4	B	5	0	0	0	0
5	A	2	0	0	0	0
5	B	2	0	0	0	0
6	A	115	0	0	1	0
6	B	140	0	0	0	0
All	All	9287	0	8925	65	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (65) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:155:THR:HB	3:A:702:SRM:HAB2	1.29	1.10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:155:THR:HB	3:B:702:SRM:HAB2	1.34	1.09
1:A:124:ARG:HE	1:A:161:GLN:HE21	1.19	0.87
1:B:124:ARG:HE	1:B:161:GLN:HE21	1.23	0.87
1:B:116:GLN:O	1:B:117:LYS:HB2	1.83	0.77
1:B:355:GLN:HA	1:B:372:MET:HE1	1.71	0.72
1:B:155:THR:HB	3:B:702:SRM:CAB	2.20	0.68
1:B:116:GLN:O	1:B:117:LYS:CB	2.43	0.66
1:A:103:HIS:CD2	1:A:232:TRP:HE1	2.14	0.65
1:B:103:HIS:CD2	1:B:232:TRP:HE1	2.15	0.64
1:B:72:GLU:HB2	1:B:271:GLN:HE22	1.62	0.63
1:A:72:GLU:HB2	1:A:271:GLN:HE22	1.64	0.62
1:A:155:THR:HB	3:A:702:SRM:CAB	2.18	0.61
1:A:161:GLN:NE2	3:A:702:SRM:HMB3	2.16	0.59
1:A:123:LEU:CD1	1:A:174:LEU:HD11	2.35	0.56
3:A:702:SRM:HBB2	3:A:702:SRM:CHC	2.35	0.56
1:A:186:GLY:HA2	1:A:191:LEU:HD12	1.87	0.56
1:A:424:VAL:H	1:A:455:ASN:ND2	2.05	0.55
3:B:702:SRM:HBB2	3:B:702:SRM:CHC	2.36	0.54
1:B:424:VAL:H	1:B:455:ASN:ND2	2.06	0.54
1:B:186:GLY:HA2	1:B:191:LEU:HD12	1.89	0.54
1:A:101:LYS:O	1:A:188:CYS:HB2	2.08	0.53
1:B:101:LYS:O	1:B:188:CYS:HB2	2.10	0.51
1:B:193:ARG:HG2	1:B:193:ARG:HH11	1.75	0.51
1:A:193:ARG:HG2	1:A:193:ARG:HH11	1.76	0.50
1:B:598:TYR:OH	1:B:620:LYS:HE3	2.12	0.50
1:B:303:ASP:OD1	1:B:303:ASP:C	2.56	0.49
1:A:123:LEU:HD13	1:A:174:LEU:HD11	1.94	0.49
1:A:598:TYR:OH	1:A:620:LYS:HE3	2.13	0.49
1:A:303:ASP:OD1	1:A:303:ASP:C	2.55	0.49
1:B:321:ARG:HH22	1:B:323:HIS:HD2	1.61	0.49
1:B:424:VAL:H	1:B:455:ASN:HD22	1.59	0.49
3:A:702:SRM:HBB2	3:A:702:SRM:HHC	1.94	0.48
1:B:323:HIS:HE1	1:B:556:GLY:O	1.96	0.48
1:A:77:LEU:O	1:A:103:HIS:HE1	1.96	0.48
1:A:519:ARG:NH1	1:A:529:ASP:HA	2.29	0.48
1:A:587:ASP:OD1	1:A:590:LYS:HE2	2.14	0.48
1:B:519:ARG:NH1	1:B:529:ASP:HA	2.29	0.48
1:A:572:GLN:HA	6:A:819:HOH:O	2.14	0.47
1:B:512:LEU:HB2	1:B:513:PRO:HD3	1.95	0.47
1:A:124:ARG:NE	1:A:161:GLN:HE21	2.01	0.47
1:A:424:VAL:H	1:A:455:ASN:HD22	1.61	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:161:GLN:NE2	3:B:702:SRM:HMB3	2.30	0.47
1:B:117:LYS:HE3	1:B:117:LYS:HA	1.96	0.46
1:A:114:ARG:HG2	1:A:114:ARG:HH11	1.80	0.46
1:A:123:LEU:HD11	1:A:174:LEU:HD11	1.98	0.46
3:B:702:SRM:HAB2	3:B:702:SRM:O3B	2.16	0.45
1:B:319:MET:HE2	1:B:319:MET:HA	1.98	0.44
1:A:319:MET:HA	1:A:319:MET:HE2	1.99	0.44
1:A:276:LYS:NZ	4:A:703:PO4:O4	2.47	0.43
1:B:453:ASN:HA	1:B:492:MET:SD	2.58	0.43
3:B:702:SRM:HBA1	3:B:702:SRM:HBB	2.01	0.43
1:A:124:ARG:HE	1:A:161:GLN:NE2	2.01	0.42
1:A:116:GLN:HG3	1:A:118:ASN:OD1	2.20	0.42
1:A:372:MET:CE	1:A:376:TRP:CZ3	3.02	0.42
1:B:156:THR:OG1	3:B:702:SRM:HAB1	2.19	0.42
1:B:124:ARG:HE	1:B:161:GLN:NE2	2.04	0.42
1:B:321:ARG:HD2	1:B:330:PRO:HD3	2.01	0.42
1:B:627:LYS:HB3	1:B:627:LYS:NZ	2.35	0.42
1:B:124:ARG:HH21	1:B:161:GLN:NE2	2.19	0.41
1:A:77:LEU:HD13	1:A:102:PHE:HB3	2.01	0.41
1:A:512:LEU:HB2	1:A:513:PRO:HD3	2.01	0.41
1:A:123:LEU:HD11	1:A:184:THR:HG21	2.03	0.41
1:A:598:TYR:HE2	1:A:616:THR:HG21	1.86	0.40
1:B:77:LEU:O	1:B:103:HIS:HE1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	560/583 (96%)	550 (98%)	9 (2%)	1 (0%)	43	51
1	B	556/583 (95%)	545 (98%)	10 (2%)	1 (0%)	43	51

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	1116/1166 (96%)	1095 (98%)	19 (2%)	2 (0%)	43	51

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	116	GLN
1	B	117	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	481/493 (98%)	470 (98%)	11 (2%)	44	59
1	B	477/493 (97%)	466 (98%)	11 (2%)	44	59
All	All	958/986 (97%)	936 (98%)	22 (2%)	44	59

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	63	LYS
1	A	114	ARG
1	A	238	ILE
1	A	242	GLU
1	A	255	SER
1	A	324	ARG
1	A	411	GLN
1	A	435	LYS
1	A	482	LYS
1	A	516	LYS
1	A	627	LYS
1	B	113	VAL
1	B	114	ARG
1	B	116	GLN
1	B	117	LYS
1	B	123	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	138	LEU
1	B	324	ARG
1	B	411	GLN
1	B	482	LYS
1	B	516	LYS
1	B	627	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	103	HIS
1	A	127	ASN
1	A	135	GLN
1	A	161	GLN
1	A	169	ASN
1	A	180	ASN
1	A	271	GLN
1	A	403	GLN
1	A	405	ASN
1	A	455	ASN
1	A	463	GLN
1	A	488	ASN
1	A	572	GLN
1	B	103	HIS
1	B	116	GLN
1	B	118	ASN
1	B	135	GLN
1	B	161	GLN
1	B	180	ASN
1	B	256	HIS
1	B	271	GLN
1	B	323	HIS
1	B	411	GLN
1	B	426	ASN
1	B	455	ASN
1	B	488	ASN
1	B	572	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SF4	B	701	1	0,12,12	-	-	-		
4	PO4	B	703	-	4,4,4	1.24	0	6,6,6	0.85	0
3	SRM	B	702	1	68,70,70	2.81	28 (41%)	86,112,112	3.90	38 (44%)
3	SRM	A	702	1	68,70,70	2.74	29 (42%)	86,112,112	3.66	36 (41%)
4	PO4	A	703	-	4,4,4	1.22	1 (25%)	6,6,6	0.64	0
2	SF4	A	701	1	0,12,12	-	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	B	701	1	-	-	0/6/5/5
3	SRM	A	702	1	3/3/19/23	11/38/126/126	-
3	SRM	B	702	1	3/3/19/23	12/38/126/126	-
2	SF4	A	701	1	-	-	0/6/5/5

All (58) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	SRM	C4B-NB	-10.41	1.27	1.35
3	A	702	SRM	CAB-CBB	6.41	1.72	1.52
3	B	702	SRM	CHD-C4C	6.11	1.50	1.38
3	A	702	SRM	CAB-C3B	-5.80	1.41	1.54
3	B	702	SRM	CHC-C1C	5.66	1.49	1.38
3	A	702	SRM	C4B-NB	-5.57	1.31	1.35
3	B	702	SRM	CAB-CBB	5.55	1.69	1.52
3	A	702	SRM	CHC-C1C	5.53	1.49	1.38
3	A	702	SRM	CHD-C4C	5.21	1.48	1.38
3	A	702	SRM	CDC-CEC	-5.18	1.43	1.51
3	A	702	SRM	CBB-CCB	4.97	1.62	1.50
3	B	702	SRM	C3C-C2C	4.75	1.46	1.36
3	A	702	SRM	C3B-C4B	-4.69	1.41	1.51
3	B	702	SRM	C3B-C4B	-4.58	1.41	1.51
3	B	702	SRM	C1D-ND	-4.51	1.31	1.39
3	A	702	SRM	C3C-C2C	4.46	1.46	1.36
3	A	702	SRM	FE-NA	4.45	2.08	1.94
3	A	702	SRM	CHD-C1D	4.42	1.49	1.39
3	B	702	SRM	CAB-C3B	-4.41	1.44	1.54
3	B	702	SRM	CBB-CCB	4.23	1.60	1.50
3	B	702	SRM	C4C-NC	-4.22	1.31	1.39
3	B	702	SRM	CHA-C4D	4.12	1.48	1.39
3	B	702	SRM	C2B-C3B	-4.02	1.44	1.55
3	A	702	SRM	FE-NC	3.91	2.08	1.95
3	A	702	SRM	CHA-C4D	3.90	1.48	1.39
3	B	702	SRM	CDC-CEC	-3.87	1.45	1.51
3	A	702	SRM	C1D-ND	-3.82	1.32	1.39
3	A	702	SRM	C2B-C3B	-3.74	1.45	1.55
3	A	702	SRM	CHA-C1A	3.70	1.47	1.37
3	B	702	SRM	FE-ND	3.62	2.07	1.95
3	A	702	SRM	C2A-C3A	-3.43	1.46	1.55
3	B	702	SRM	O4D-CED	-3.40	1.19	1.30
3	B	702	SRM	FE-NA	3.40	2.05	1.94
3	A	702	SRM	C4A-NA	-3.27	1.32	1.35
3	B	702	SRM	C3A-C4A	-3.19	1.44	1.51
3	A	702	SRM	FE-ND	3.19	2.05	1.95
3	A	702	SRM	C1C-NC	-3.18	1.33	1.39
3	B	702	SRM	FE-NC	3.15	2.05	1.95
3	B	702	SRM	C2A-C3A	-2.98	1.47	1.55
3	B	702	SRM	CHD-C1D	2.94	1.46	1.39
3	A	702	SRM	O4C-CEC	-2.84	1.21	1.30
3	A	702	SRM	C4C-NC	-2.64	1.34	1.39
3	B	702	SRM	FE-NB	2.63	2.03	1.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	SRM	C4D-ND	-2.61	1.34	1.39
3	B	702	SRM	CHA-C1A	2.53	1.44	1.37
3	A	702	SRM	C3A-C4A	-2.51	1.45	1.51
3	B	702	SRM	C1D-C2D	2.47	1.49	1.44
3	B	702	SRM	C1C-NC	-2.29	1.35	1.39
3	A	702	SRM	C1D-C2D	2.28	1.49	1.44
3	B	702	SRM	CHB-C4A	2.28	1.45	1.39
3	A	702	SRM	O4A-CEA	-2.28	1.23	1.30
3	A	702	SRM	O2D-CCD	-2.27	1.23	1.30
4	A	703	PO4	P-O4	-2.21	1.48	1.54
3	A	702	SRM	CDA-CEA	2.15	1.57	1.50
3	A	702	SRM	CHB-C4A	2.11	1.45	1.39
3	B	702	SRM	CDD-CED	-2.08	1.47	1.51
3	B	702	SRM	O1B-CCB	2.04	1.28	1.22
3	A	702	SRM	CDC-C2C	-2.02	1.48	1.51

All (74) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	702	SRM	CMB-C2B-C3B	13.05	136.13	112.05
3	A	702	SRM	C2B-C3B-C4B	12.47	116.69	100.89
3	B	702	SRM	CAB-CBB-CCB	11.41	142.94	112.49
3	A	702	SRM	CAA-C3A-C4A	11.26	130.83	111.19
3	A	702	SRM	CAB-CBB-CCB	11.24	142.47	112.49
3	B	702	SRM	CAA-C3A-C4A	11.22	130.76	111.19
3	A	702	SRM	CMB-C2B-C3B	10.76	131.91	112.05
3	B	702	SRM	C2B-C3B-C4B	10.70	114.45	100.89
3	B	702	SRM	CBB-CAB-C3B	-10.08	86.07	114.65
3	A	702	SRM	CBB-CAB-C3B	-8.67	90.06	114.65
3	B	702	SRM	CMA-C2A-C3A	-8.57	96.23	112.05
3	A	702	SRM	O2B-CCB-O1B	-7.20	104.83	123.33
3	B	702	SRM	C2B-C1B-CHB	-6.83	117.41	123.54
3	B	702	SRM	CAB-C3B-C4B	6.37	122.30	111.19
3	A	702	SRM	CMA-C2A-C3A	-6.35	100.31	112.05
3	A	702	SRM	CAB-C3B-C4B	5.94	121.55	111.19
3	B	702	SRM	C3B-C2B-C1B	-5.83	91.03	100.92
3	A	702	SRM	C2B-C1B-CHB	-5.81	118.32	123.54
3	B	702	SRM	CMB-C2B-CDB	-5.72	101.02	110.74
3	A	702	SRM	CDB-C2B-C3B	-5.71	93.30	108.37
3	B	702	SRM	CDB-C2B-C3B	-5.60	93.58	108.37
3	B	702	SRM	CDA-C2A-C3A	5.59	123.13	108.37
3	A	702	SRM	C3B-C2B-C1B	-5.57	91.47	100.92

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	SRM	CDA-C2A-C3A	5.43	122.70	108.37
3	B	702	SRM	C3C-C4C-NC	5.35	115.49	110.32
3	A	702	SRM	C2A-C3A-C4A	5.22	107.50	100.89
3	A	702	SRM	C4C-C3C-C2C	-4.96	101.30	106.84
3	B	702	SRM	C2A-C3A-C4A	4.90	107.10	100.89
3	B	702	SRM	CAC-CBC-CCC	-4.87	100.75	113.67
3	B	702	SRM	O2B-CCB-O1B	-4.83	110.92	123.33
3	A	702	SRM	C3C-C4C-NC	4.28	114.45	110.32
3	B	702	SRM	CDA-C2A-C1A	4.23	120.21	107.11
3	B	702	SRM	CHD-C4C-NC	-4.16	119.92	124.45
3	B	702	SRM	C4A-CHB-C1B	-4.02	119.93	125.84
3	B	702	SRM	C1D-C2D-C3D	-4.00	102.62	106.81
3	A	702	SRM	C4D-C3D-C2D	-3.92	102.70	106.81
3	B	702	SRM	C4C-C3C-C2C	-3.89	102.49	106.84
3	A	702	SRM	CDA-C2A-C1A	3.84	118.99	107.11
3	A	702	SRM	O2B-CCB-CBB	3.74	125.83	114.00
3	B	702	SRM	CBD-CAD-C2D	-3.64	102.47	112.53
3	A	702	SRM	CBD-CAD-C2D	-3.33	103.33	112.53
3	B	702	SRM	O3C-CEC-CDC	-3.29	112.74	122.11
3	B	702	SRM	CAB-C3B-C2B	3.29	123.23	114.19
3	A	702	SRM	CMB-C2B-C1B	3.25	125.73	109.59
3	B	702	SRM	C4D-C3D-C2D	-3.22	103.43	106.81
3	A	702	SRM	C3D-C4D-ND	3.14	113.64	110.15
3	B	702	SRM	C1C-C2C-C3C	-3.12	103.35	106.84
3	B	702	SRM	CMB-C2B-C1B	3.09	124.92	109.59
3	A	702	SRM	CAC-C3C-C4C	2.92	130.80	124.85
3	B	702	SRM	O4C-CEC-O3C	2.92	130.84	123.33
3	A	702	SRM	CAB-C3B-C2B	2.76	121.77	114.19
3	A	702	SRM	CAC-CBC-CCC	-2.72	106.46	113.67
3	A	702	SRM	O4C-CEC-O3C	2.72	130.32	123.33
3	A	702	SRM	C4A-CHB-C1B	-2.71	121.86	125.84
3	B	702	SRM	C2D-C1D-ND	2.64	113.08	110.15
3	A	702	SRM	CEC-CDC-C2C	2.64	120.40	113.76
3	B	702	SRM	C2C-C1C-NC	2.62	112.85	110.32
3	B	702	SRM	O2A-CCA-CBA	2.60	122.22	114.00
3	B	702	SRM	O2A-CCA-O1A	-2.59	116.67	123.33
3	B	702	SRM	CHC-C1C-NC	-2.58	121.65	124.45
3	B	702	SRM	C3A-C4A-CHB	-2.48	118.06	123.33
3	A	702	SRM	C1D-C2D-C3D	-2.46	104.23	106.81
3	B	702	SRM	CAA-C3A-C2A	2.44	120.90	114.19
3	B	702	SRM	C3D-C4D-ND	2.37	112.78	110.15
3	A	702	SRM	C2D-C1D-ND	2.35	112.76	110.15

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	702	SRM	CDB-C2B-C1B	-2.35	99.84	107.11
3	A	702	SRM	CMB-C2B-CDB	-2.31	106.80	110.74
3	A	702	SRM	O2D-CCD-CBD	2.21	120.97	114.00
3	A	702	SRM	CHC-C1C-NC	-2.20	122.06	124.45
3	B	702	SRM	O2B-CCB-CBB	2.20	120.94	114.00
3	A	702	SRM	C3A-C2A-C1A	-2.12	97.32	100.92
3	B	702	SRM	CAD-C2D-C3D	2.06	131.72	125.30
3	A	702	SRM	CBA-CAA-C3A	2.06	120.49	114.65
3	A	702	SRM	CAA-C3A-C2A	2.05	119.83	114.19

All (6) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
3	A	702	SRM	C3B
3	A	702	SRM	C3A
3	A	702	SRM	NC
3	B	702	SRM	C3B
3	B	702	SRM	C3A
3	B	702	SRM	NC

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	702	SRM	C1A-C2A-CDA-CEA
3	A	702	SRM	CMA-C2A-CDA-CEA
3	A	702	SRM	C2A-C3A-CAA-CBA
3	B	702	SRM	C1A-C2A-CDA-CEA
3	B	702	SRM	CMA-C2A-CDA-CEA
3	B	702	SRM	C3A-C2A-CDA-CEA
3	B	702	SRM	C2A-C3A-CAA-CBA
3	A	702	SRM	C2B-C3B-CAB-CBB
3	B	702	SRM	C3C-CAC-CBC-CCC
3	B	702	SRM	C2B-C3B-CAB-CBB
3	A	702	SRM	C3B-C2B-CDB-CEB
3	B	702	SRM	CAA-CBA-CCA-O2A
3	A	702	SRM	C3C-CAC-CBC-CCC
3	A	702	SRM	CAA-CBA-CCA-O1A
3	A	702	SRM	C3A-C2A-CDA-CEA
3	B	702	SRM	C3B-C2B-CDB-CEB
3	B	702	SRM	CAA-CBA-CCA-O1A
3	A	702	SRM	CAA-CBA-CCA-O2A
3	B	702	SRM	CAC-CBC-CCC-O2C

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	702	SRM	CAC-CBC-CCC-O2C
3	A	702	SRM	CAC-CBC-CCC-O1C
3	B	702	SRM	CAC-CBC-CCC-O1C
3	B	702	SRM	CAB-CBB-CCB-O1B


There are no ring outliers.

3 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	702	SRM	7	0
3	A	702	SRM	5	0
4	A	703	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	564/583 (96%)	-0.18	14 (2%) 58 55	15, 24, 42, 70	0
1	B	560/583 (96%)	-0.28	7 (1%) 75 73	15, 23, 38, 67	0
All	All	1124/1166 (96%)	-0.23	21 (1%) 66 63	15, 24, 40, 70	0

All (21) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	235	GLY	6.8
1	A	239	MET	4.9
1	B	114	ARG	4.5
1	A	324	ARG	3.3
1	B	627	LYS	3.0
1	A	627	LYS	3.0
1	A	305	ALA	2.9
1	A	237	LYS	2.9
1	A	116	GLN	2.6
1	B	116	GLN	2.5
1	A	114	ARG	2.4
1	A	242	GLU	2.4
1	A	63	LYS	2.4
1	B	63	LYS	2.4
1	B	242	GLU	2.3
1	A	238	ILE	2.3
1	A	306	GLY	2.2
1	B	324	ARG	2.1
1	A	623	GLU	2.1
1	A	235	GLY	2.1
1	A	115	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	PO4	A	703	5/5	0.95	0.12	29,30,35,41	0
5	MG	A	705	1/1	0.95	0.06	30,30,30,30	0
4	PO4	B	703	5/5	0.96	0.10	29,30,39,41	0
3	SRM	A	702	63/63	0.96	0.07	15,20,24,31	0
3	SRM	B	702	63/63	0.97	0.07	16,19,25,27	0
2	SF4	B	701	8/8	0.97	0.04	18,18,19,19	0
5	MG	B	705	1/1	0.97	0.06	27,27,27,27	0
5	MG	B	704	1/1	0.98	0.08	8,8,8,8	0
2	SF4	A	701	8/8	0.98	0.04	16,18,19,20	0
5	MG	A	704	1/1	0.99	0.08	8,8,8,8	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around SRM A 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around SRM B 702:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.