



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:24 PM UTC

PDB ID : 8H8Y / pdb_00008h8y
Title : Crystal structure of AbHheG from Acidimicrobiia bacterium
Authors : Zhou, C.H.; Chen, X.; Han, X.; Liu, W.D.; Wu, Q.Q.; Zhu, D.M.; Ma, Y.H.
Deposited on : 2022-10-24
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
Ramachandran outliers	187476	2153 (1.56-1.56)
Sidechain outliers	187428	2150 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	246	7% 93% 7%
1	B	246	8% 93% 7%
1	C	246	% 93% 6%
1	D	246	13% 89% 9%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8082 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called alpha/beta hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	0	0	0
			1841	1156	321	352	12			
1	B	245	Total	C	N	O	S	0	0	0
			1833	1151	320	351	11			
1	C	244	Total	C	N	O	S	0	0	0
			1827	1148	319	349	11			
1	D	244	Total	C	N	O	S	0	0	0
			1827	1148	319	349	11			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	C	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	194	Total	O	0	0
			194	194		
3	B	180	Total	O	0	0
			180	180		
3	C	216	Total	O	0	0
			216	216		
3	D	146	Total	O	0	0
			146	146		



3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: alpha/beta hydrolase





- Molecule 1: alpha/beta hydrolase



- Molecule 1: alpha/beta hydrolase



- Molecule 1: alpha/beta hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	60.89Å 62.41Å 71.42Å 69.24° 82.88° 62.19°	Depositor
Resolution (Å)	52.01 – 1.55 52.01 – 1.55	Depositor EDS
% Data completeness (in resolution range)	95.7 (52.01-1.55) 95.7 (52.01-1.55)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 1.55Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.172 , 0.200 0.183 , 0.207	Depositor DCC
R_{free} test set	5949 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.257	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 36.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8082	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	1/1877 (0.1%)	1.15	0/2546
1	B	1.02	1/1869 (0.1%)	1.15	0/2536
1	C	1.04	0/1863	1.14	0/2528
1	D	1.01	0/1863	1.20	2/2528 (0.1%)
All	All	1.03	2/7472 (0.0%)	1.16	2/10138 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	19	ASP	CG-OD2	5.57	1.35	1.25
1	A	19	ASP	CG-OD1	5.50	1.35	1.25

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	206	SER	N-CA-C	-5.51	102.73	110.50
1	D	206	SER	CA-C-O	-5.03	116.08	121.56

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1841	0	1818	8	0
1	B	1833	0	1806	11	0
1	C	1827	0	1801	9	0
1	D	1827	0	1801	11	0
2	B	6	0	8	0	0
2	C	6	0	8	0	0
2	D	6	0	8	0	0
3	A	194	0	0	1	0
3	B	180	0	0	2	0
3	C	216	0	0	1	0
3	D	146	0	0	0	0
All	All	8082	0	7250	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:190:LEU:HA	1:A:195:ALA:HB3	1.66	0.77
1:A:157:ALA:HB2	1:B:157:ALA:HB2	1.79	0.63
1:A:195:ALA:O	1:A:201:ARG:HB2	2.00	0.62
1:C:157:ALA:HB2	1:D:157:ALA:HB2	1.81	0.61
1:B:90:ILE:HG22	3:B:401:HOH:O	2.07	0.55
1:B:190:LEU:HA	1:B:195:ALA:HB3	1.91	0.52
1:A:205:GLU:O	1:A:211:ARG:HA	2.09	0.52
1:B:197:ASP:OD2	1:B:200:VAL:HG23	2.09	0.52
1:C:50:ARG:HD3	3:C:421:HOH:O	2.10	0.50
1:D:113:ALA:HB3	1:D:114:PRO:HD3	1.94	0.49
1:B:233:PHE:HB3	1:D:241:TYR:CE2	2.49	0.48
1:B:232:ARG:O	1:D:210:MET:HE1	2.14	0.47
1:D:162:VAL:HG13	1:D:176:VAL:HG12	1.97	0.47
1:B:210:MET:HE1	1:D:232:ARG:O	2.15	0.46
1:B:119:ARG:HA	1:B:119:ARG:HD2	1.77	0.45
1:C:91:VAL:HG21	1:C:106:VAL:CG1	2.46	0.45
1:C:91:VAL:HG21	1:C:106:VAL:HG12	1.97	0.45
1:B:135:ILE:O	1:B:178:ALA:HA	2.18	0.43
1:A:232:ARG:O	1:C:210:MET:HE1	2.19	0.43
1:D:183:PHE:HE2	1:D:245:TRP:CG	2.35	0.43
1:C:162:VAL:HG13	1:C:176:VAL:HG12	2.01	0.43
1:C:3:ASN:CG	1:C:79:ARG:HH11	2.28	0.42
1:D:7:ALA:HB3	1:D:31:LEU:HD23	2.02	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:3:ASN:ND2	1:C:79:ARG:HH11	2.18	0.42
1:A:128:ARG:HD2	3:A:431:HOH:O	2.19	0.42
1:C:43:GLU:O	1:C:47:MET:HG3	2.19	0.42
1:D:186:PHE:CE2	1:D:188:ALA:HB3	2.55	0.42
1:A:6:VAL:HG21	1:A:77:PHE:CB	2.50	0.41
1:D:135:ILE:O	1:D:178:ALA:HA	2.21	0.41
1:B:124:VAL:O	1:B:128:ARG:HD2	2.20	0.41
1:A:189:PHE:O	1:A:192:ALA:O	2.38	0.41
1:B:119:ARG:NE	3:B:403:HOH:O	2.53	0.40
1:D:206:SER:O	1:D:207:GLN:CB	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/246 (99%)	237 (97%)	7 (3%)	0	100	100
1	B	243/246 (99%)	237 (98%)	6 (2%)	0	100	100
1	C	242/246 (98%)	236 (98%)	6 (2%)	0	100	100
1	D	242/246 (98%)	235 (97%)	7 (3%)	0	100	100
All	All	971/984 (99%)	945 (97%)	26 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	194/194 (100%)	188 (97%)	6 (3%)	35	8
1	B	193/194 (100%)	190 (98%)	3 (2%)	55	28
1	C	192/194 (99%)	189 (98%)	3 (2%)	55	28
1	D	192/194 (99%)	184 (96%)	8 (4%)	26	4
All	All	771/776 (99%)	751 (97%)	20 (3%)	40	13

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	TYR
1	A	128	ARG
1	A	191	LYS
1	A	199	GLU
1	A	206	SER
1	A	207	GLN
1	B	50	ARG
1	B	119	ARG
1	B	203	LYS
1	C	175	GLN
1	C	191	LYS
1	C	203	LYS
1	D	3	ASN
1	D	89	GLN
1	D	128	ARG
1	D	171	ARG
1	D	191	LYS
1	D	203	LYS
1	D	206	SER
1	D	207	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	67	GLN
1	B	105	GLN
1	D	3	ASN
1	D	67	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GOL	C	301	-	5,5,5	0.08	0	5,5,5	0.64	0
2	GOL	B	301	-	5,5,5	0.07	0	5,5,5	0.35	0
2	GOL	D	301	-	5,5,5	0.10	0	5,5,5	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	301	-	-	2/4/4/4	-
2	GOL	B	301	-	-	2/4/4/4	-
2	GOL	D	301	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	301	GOL	O1-C1-C2-C3
2	C	301	GOL	O1-C1-C2-O2
2	B	301	GOL	O2-C2-C3-O3
2	B	301	GOL	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	246/246 (100%)	0.25	17 (6%) 23 26	10, 16, 43, 70	0
1	B	245/246 (99%)	0.30	19 (7%) 19 23	10, 17, 51, 76	0
1	C	244/246 (99%)	0.08	3 (1%) 76 83	10, 16, 30, 39	0
1	D	244/246 (99%)	0.65	31 (12%) 8 9	11, 21, 49, 74	0
All	All	979/984 (99%)	0.32	70 (7%) 21 25	10, 17, 44, 76	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	202	ALA	5.8
1	D	200	VAL	5.7
1	D	202	ALA	5.3
1	A	206	SER	5.2
1	D	192	ALA	5.2
1	A	199	GLU	5.1
1	D	195	ALA	5.0
1	A	192	ALA	4.8
1	D	188	ALA	4.7
1	A	200	VAL	4.5
1	B	195	ALA	4.4
1	D	190	LEU	4.3
1	A	14	TYR	4.3
1	D	14	TYR	4.3
1	D	189	PHE	4.3
1	C	3	ASN	4.0
1	B	194	GLY	3.9
1	A	195	ALA	3.9
1	D	197	ASP	3.8
1	D	198	PRO	3.5
1	B	206	SER	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	191	LYS	3.4
1	D	203	LYS	3.4
1	D	204	ILE	3.3
1	D	194	GLY	3.3
1	B	193	SER	3.3
1	B	200	VAL	3.2
1	B	207	GLN	3.2
1	B	198	PRO	3.1
1	D	187	PRO	3.0
1	D	199	GLU	3.0
1	B	39	ASN	3.0
1	B	204	ILE	3.0
1	A	196	ASN	3.0
1	B	190	LEU	2.9
1	C	14	TYR	2.8
1	D	28	GLY	2.8
1	A	202	ALA	2.8
1	D	39	ASN	2.8
1	A	198	PRO	2.8
1	D	196	ASN	2.8
1	D	208	VAL	2.7
1	B	90	ILE	2.7
1	D	46	SER	2.7
1	D	206	SER	2.7
1	D	201	ARG	2.7
1	A	204	ILE	2.7
1	B	188	ALA	2.6
1	D	207	GLN	2.6
1	A	188	ALA	2.6
1	A	207	GLN	2.5
1	D	38	PRO	2.5
1	D	3	ASN	2.5
1	A	190	LEU	2.4
1	B	46	SER	2.4
1	C	204	ILE	2.3
1	A	193	SER	2.3
1	A	208	VAL	2.3
1	D	41	VAL	2.3
1	D	193	SER	2.2
1	B	196	ASN	2.2
1	A	187	PRO	2.2
1	B	203	LYS	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	2	SER	2.2
1	D	128	ARG	2.2
1	D	186	PHE	2.2
1	A	205	GLU	2.1
1	B	205	GLU	2.1
1	B	189	PHE	2.1
1	D	49	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	B	301	6/6	0.87	0.11	27,28,31,31	0
2	GOL	D	301	6/6	0.87	0.12	28,29,32,32	0
2	GOL	C	301	6/6	0.89	0.10	25,28,29,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.