



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:58 PM UTC

PDB ID : 9H8C / pdb_00009h8c
Title : Human CDK8/Cyclin-C complex with inhibitor 2-9
Authors : Somers, D.O.
Deposited on : 2024-10-29
Resolution : 2.57 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4770 (2.60-2.56)
Clashscore	190562	5124 (2.60-2.56)
Ramachandran outliers	187476	5046 (2.60-2.56)
Sidechain outliers	187428	5046 (2.60-2.56)
RSRZ outliers	180081	4770 (2.60-2.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	414	
2	B	313	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 5344 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclin-dependent kinase 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2782	1789	488	488	17	0	1	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	GLY	-	expression tag	UNP P49336
A	-9	PRO	-	expression tag	UNP P49336
A	-8	MET	-	expression tag	UNP P49336
A	-7	ASP	-	expression tag	UNP P49336
A	-6	TYR	-	expression tag	UNP P49336
A	-5	LYS	-	expression tag	UNP P49336
A	-4	ASP	-	expression tag	UNP P49336
A	-3	ASP	-	expression tag	UNP P49336
A	-2	ASP	-	expression tag	UNP P49336
A	-1	ASP	-	expression tag	UNP P49336
A	0	LYS	-	expression tag	UNP P49336

- Molecule 2 is a protein called Cyclin-C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	266	2236	1465	368	386	17	0	2	0

There are 30 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	-29	GLY	-	expression tag	UNP P24863
B	-28	SER	-	expression tag	UNP P24863
B	-27	GLY	-	expression tag	UNP P24863
B	-26	MET	-	expression tag	UNP P24863
B	-25	LYS	-	expression tag	UNP P24863

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	-24	GLU	-	expression tag	UNP P24863
B	-23	THR	-	expression tag	UNP P24863
B	-22	ALA	-	expression tag	UNP P24863
B	-21	ALA	-	expression tag	UNP P24863
B	-20	ALA	-	expression tag	UNP P24863
B	-19	LYS	-	expression tag	UNP P24863
B	-18	PHE	-	expression tag	UNP P24863
B	-17	GLU	-	expression tag	UNP P24863
B	-16	ARG	-	expression tag	UNP P24863
B	-15	GLN	-	expression tag	UNP P24863
B	-14	HIS	-	expression tag	UNP P24863
B	-13	MET	-	expression tag	UNP P24863
B	-12	ASP	-	expression tag	UNP P24863
B	-11	SER	-	expression tag	UNP P24863
B	-10	PRO	-	expression tag	UNP P24863
B	-9	ASP	-	expression tag	UNP P24863
B	-8	LEU	-	expression tag	UNP P24863
B	-7	GLY	-	expression tag	UNP P24863
B	-6	THR	-	expression tag	UNP P24863
B	-5	ASP	-	expression tag	UNP P24863
B	-4	ASP	-	expression tag	UNP P24863
B	-3	ASP	-	expression tag	UNP P24863
B	-2	ASP	-	expression tag	UNP P24863
B	-1	LYS	-	expression tag	UNP P24863
B	0	ALA	-	expression tag	UNP P24863

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is 4-[(3 {S})-3-[2-[(3 {R})-3-fluoranylpyrrolidin-1-yl]pyrimidin-4-yl]piperidin-1-yl]carbonyl-1 {H}-pyrrole-2-carbonitrile (CCD ID: A1IS8) (formula: $C_{19}H_{21}FN_6O$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	A	1	Total	C	F	N	O	0	0
			27	19	1	6	1		

- Molecule 6 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	B	1	Total	C	O	S	0	0
			4	2	1	1		
6	B	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	147	Total 147	O 147	0	0
7	B	128	Total 128	O 128	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	71.53Å 69.21Å 168.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.00 – 2.57 64.00 – 2.57	Depositor EDS
% Data completeness (in resolution range)	73.2 (64.00-2.57) 73.2 (64.00-2.57)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 2.58Å)	Xtriage
Refinement program	BUSTER 2.11.8 (26-JUL-2023)	Depositor
R, R_{free}	0.189 , 0.262 0.182 , 0.245	Depositor DCC
R_{free} test set	933 reflections (3.42%)	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.018 for k,h,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5344	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, GOL, A1IS8, DMS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/2853	1.09	2/3841 (0.1%)
2	B	0.65	0/2295	1.11	5/3109 (0.2%)
All	All	0.66	0/5148	1.10	7/6950 (0.1%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	244	PHE	CA-CB-CG	6.12	119.92	113.80
2	B	250	MET	N-CA-C	5.44	118.13	111.82
2	B	92	PHE	CA-CB-CG	5.35	119.15	113.80
1	A	245	ASN	CA-CB-CG	5.33	117.93	112.60
1	A	5	PHE	CA-CB-CG	5.28	119.08	113.80
2	B	15	ILE	N-CA-C	-5.16	99.25	106.53
2	B	164	GLN	N-CA-C	-5.13	105.59	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2782	0	2778	21	0
2	B	2236	0	2269	11	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	4	0	6	0	0
4	A	6	0	8	1	0
4	B	6	0	8	1	0
5	A	27	0	0	1	0
6	B	8	0	12	1	0
7	A	147	0	0	1	0
7	B	128	0	0	0	0
All	All	5344	0	5081	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:8:LYS:HZ3	1:A:12:GLU:HG3	1.51	0.76
1:A:8:LYS:NZ	1:A:12:GLU:HG3	2.01	0.75
1:A:52:LYS:NZ	5:A:503:A1IS8:N5	2.47	0.63
2:B:96:LYS:HD3	2:B:102:VAL:HG22	1.81	0.62
2:B:214:ARG:HG3	6:B:302:DMS:H23	1.87	0.57
1:A:8:LYS:HZ3	1:A:12:GLU:CG	2.17	0.56
1:A:139:ILE:HD13	1:A:142:LEU:HD12	1.86	0.56
1:A:76:PRO:HA	4:A:502:GOL:H2	1.93	0.51
1:A:54:ILE:HG21	1:A:63:ALA:HB2	1.92	0.50
2:B:157:ARG:NE	4:B:303:GOL:H11	2.27	0.50
2:B:182:ASP:HB3	2:B:237:LEU:HD21	1.94	0.49
1:A:8:LYS:NZ	1:A:12:GLU:CG	2.73	0.48
2:B:244:PHE:CZ	2:B:246:GLU:HB3	2.48	0.48
1:A:201:ALA:HA	1:A:218:TRP:CD1	2.51	0.46
1:A:47:LYS:NZ	7:A:606:HOH:O	2.49	0.45
2:B:152:VAL:HG12	2:B:154:HIS:CE1	2.51	0.45
2:B:18:LYS:O	2:B:22:LEU:HG	2.16	0.45
2:B:163:VAL:HG21	2:B:173:LEU:HD13	1.99	0.45
1:A:140:HIS:HD2	1:A:327:GLU:HG3	1.83	0.44
2:B:70:LYS:HD2	2:B:189:CYS:SG	2.57	0.44
1:A:217:ILE:HD13	1:A:220:ILE:HD12	2.00	0.44
1:A:57:THR:HB	1:A:91[B]:ARG:HB2	2.00	0.43
1:A:54:ILE:HD11	1:A:95:LEU:HD11	1.99	0.43
1:A:151:ASP:CG	1:A:153:LYS:HZ1	2.26	0.43
1:A:276:HIS:O	1:A:280:MET:HG2	2.18	0.42
1:A:306:SER:O	1:A:310:HIS:HD2	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:PRO:HD3	1:A:318:MET:HE3	2.01	0.41
1:A:290:ASN:HB3	1:A:295:LYS:NZ	2.35	0.41
1:A:294:ILE:HD13	1:A:304:PRO:HB3	2.02	0.41
2:B:83:PRO:HA	2:B:86:MET:HB3	2.02	0.41
2:B:246:GLU:HB2	2:B:250:MET:HG2	2.04	0.41
1:A:319:ASP:OD1	1:A:321:ILE:HG13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	329/414 (80%)	306 (93%)	22 (7%)	1 (0%)	36	56
2	B	266/313 (85%)	263 (99%)	3 (1%)	0	100	100
All	All	595/727 (82%)	569 (96%)	25 (4%)	1 (0%)	43	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	125	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	299/368 (81%)	279 (93%)	20 (7%)	15	31
2	B	245/281 (87%)	237 (97%)	8 (3%)	33	59
All	All	544/649 (84%)	516 (95%)	28 (5%)	21	43

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	8	LYS
1	A	19	LEU
1	A	25	CYS
1	A	44	LYS
1	A	45	ASP
1	A	80	SER
1	A	82	GLN
1	A	104	LEU
1	A	123	LEU
1	A	150	ARG
1	A	196	THR
1	A	209	ARG
1	A	264	ASP
1	A	281	LYS
1	A	294	ILE
1	A	298	GLU
1	A	327	GLU
1	A	331	GLN
1	A	360	THR
2	B	40	LEU
2	B	58	ARG
2	B	100	PHE
2	B	140	PHE
2	B	160	LEU
2	B	171	MET
2	B	237	LEU
2	B	252	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	77	ASN
1	A	140	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	210	HIS
1	A	235	HIS
1	A	300	HIS
1	A	310	HIS
2	B	19	GLN
2	B	49	GLN
2	B	59	GLN
2	B	60	GLN
2	B	168	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	501	-	3,3,3	0.25	0	2,2,2	0.28	0
6	DMS	B	301	-	3,3,3	0.71	0	3,3,3	0.28	0
4	GOL	A	502	-	5,5,5	0.12	0	5,5,5	0.26	0
4	GOL	B	303	-	5,5,5	0.18	0	5,5,5	0.34	0
5	A1IS8	A	503	-	29,30,30	0.37	0	35,42,42	0.75	2 (5%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	DMS	B	302	-	3,3,3	0.62	0	3,3,3	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	501	-	-	1/1/1/1	-
4	GOL	A	502	-	-	0/4/4/4	-
4	GOL	B	303	-	-	1/4/4/4	-
5	A1IS8	A	503	-	-	2/16/37/37	0/4/4/4

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	503	A1IS8	C12-C8-C7	-2.02	106.88	111.89
5	A	503	A1IS8	C15-C14-C13	-2.00	119.97	126.78

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	501	EDO	O1-C1-C2-O2
5	A	503	A1IS8	O-C13-C14-C17
5	A	503	A1IS8	N3-C13-C14-C15
4	B	303	GOL	O2-C2-C3-O3


There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	502	GOL	1	0
4	B	303	GOL	1	0
5	A	503	A1IS8	1	0
6	B	302	DMS	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In

addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/414 (81%)	0.11	7 (2%) 63 59	28, 54, 81, 98	1 (0%)
2	B	266/313 (84%)	0.03	1 (0%) 88 87	22, 50, 68, 99	2 (0%)
All	All	602/727 (82%)	0.07	8 (1%) 75 72	22, 51, 77, 99	3 (0%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	123	LEU	4.1
1	A	91[A]	ARG	3.1
1	A	195	VAL	2.3
1	A	246	PRO	2.1
1	A	245	ASN	2.1
1	A	351	ILE	2.1
1	A	117	ASN	2.0
2	B	262	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	303	6/6	0.75	0.18	69,69,69,69	0
6	DMS	B	301	4/4	0.82	0.23	94,94,94,94	0
4	GOL	A	502	6/6	0.90	0.21	63,64,65,65	0
3	EDO	A	501	4/4	0.91	0.17	74,74,74,74	0
6	DMS	B	302	4/4	0.91	0.17	94,95,95,95	0
5	A1IS8	A	503	27/27	0.96	0.07	45,45,46,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around A1IS8 A 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.