



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 09:53 AM UTC

PDB ID : 1H9S / pdb_00001h9s
Title : Molybdate bound complex of Dimop domain of ModE from E.coli
Authors : Gourley, D.G.; Hunter, W.N.
Deposited on : 2001-03-19
Resolution : 1.82 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1148 (1.82-1.82)
Ramachandran outliers	187476	1140 (1.82-1.82)
Sidechain outliers	187428	1140 (1.82-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	140	
2	B	140	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MOO	A	1261	-	-	X	-
3	MOO	B	1261	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2300 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MOLYBDENUM TRANSPORT PROTEIN MODE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	138	Total	C	N	O	S	0	0	0
			1032	640	184	204	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	123	MET	LEU	engineered mutation	UNP P46930


- Molecule 2 is a protein called MOLYBDENUM TRANSPORT PROTEIN MODE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	137	Total	C	N	O	S	0	0	0
			1031	638	182	208	3			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	242	GLU	GLN	conflict	UNP P46930

- Molecule 3 is MOLYBDATE ION (CCD ID: MOO) (formula: MoO₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Mo	O	0	0
			5	1	4		
3	B	1	Total	Mo	O	0	0
			5	1	4		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	118	Total	O	0	0
			118	118		
4	B	109	Total	O	0	0
			109	109		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: MOLYBDENUM TRANSPORT PROTEIN MODE

Chain A:  78% 20% ..



• Molecule 2: MOLYBDENUM TRANSPORT PROTEIN MODE

Chain B:  78% 19% ..



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α , β , γ	72.34Å 73.34Å 49.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.82	Depositor
% Data completeness (in resolution range)	98.1 (20.00-1.82)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.49	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.188 , 0.234	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2300	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MOO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.71	0/1045	1.50	4/1424 (0.3%)
2	B	0.69	0/1044	1.53	6/1423 (0.4%)
All	All	0.70	0/2089	1.52	10/2847 (0.4%)

There are no bond length outliers.

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	ASP	N-CA-C	11.05	122.89	111.07
2	B	217	CYS	CA-C-O	-8.69	110.97	120.36
2	B	252	ASN	CA-CB-CG	-7.70	104.90	112.60
2	B	142	ASP	N-CA-C	6.48	120.92	112.89
1	A	123	MET	CA-C-O	-6.16	110.33	120.80
1	A	216	GLN	OE1-CD-NE2	-6.03	116.57	122.60
2	B	233	VAL	N-CA-CB	-5.59	106.44	112.37
2	B	249	ALA	CA-C-O	-5.58	114.69	121.05
2	B	217	CYS	CB-CA-C	-5.50	99.64	109.71
1	A	212	ARG	CA-CB-CG	-5.32	103.47	114.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1032	0	1023	21	0
2	B	1031	0	1018	19	0
3	A	5	0	0	4	0
3	B	5	0	0	4	0
4	A	118	0	0	3	0
4	B	109	0	0	3	0
All	All	2300	0	2041	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:HIS:HB3	1:A:159:LYS:HE2	1.40	1.00
1:A:260:THR:C	4:A:2118:HOH:O	2.03	0.99
3:B:1261:MOO:MO	3:B:1261:MOO:O3	1.62	0.71
3:A:1261:MOO:O3	3:A:1261:MOO:MO	1.66	0.67
3:B:1261:MOO:MO	3:B:1261:MOO:O4	1.65	0.67
3:A:1261:MOO:MO	3:A:1261:MOO:O2	1.65	0.67
1:A:147:VAL:O	1:A:159:LYS:HE3	1.97	0.65
3:A:1261:MOO:MO	3:A:1261:MOO:O4	1.69	0.63
2:B:183:LYS:HG2	2:B:185:PRO:HD2	1.83	0.61
3:A:1261:MOO:MO	3:A:1261:MOO:O1	1.71	0.60
3:B:1261:MOO:MO	3:B:1261:MOO:O2	1.72	0.60
3:B:1261:MOO:MO	3:B:1261:MOO:O1	1.72	0.60
2:B:159:LYS:N	2:B:159:LYS:HD3	2.18	0.59
2:B:202:GLN:C	2:B:203:LEU:HD12	2.26	0.59
1:A:187:VAL:HG11	1:A:229:LEU:HD21	1.85	0.58
2:B:186:TRP:CE3	2:B:255:SER:O	2.59	0.56
2:B:138:ARG:HG3	2:B:147:VAL:HG12	1.87	0.55
1:A:257:ILE:HD12	2:B:257:ILE:HD12	1.90	0.54
2:B:158:LEU:C	2:B:159:LYS:HD3	2.34	0.53
2:B:217:CYS:CB	4:B:2075:HOH:O	2.57	0.53
1:A:183:LYS:HG2	1:A:185:PRO:HD2	1.92	0.52
2:B:125:THR:H	2:B:130:GLN:HE22	1.59	0.51
1:A:180:ILE:HG23	1:A:256:VAL:HG13	1.93	0.50
1:A:180:ILE:HG23	1:A:256:VAL:CG1	2.43	0.49
2:B:217:CYS:HB2	4:B:2075:HOH:O	2.11	0.49
1:A:229:LEU:HD23	1:A:229:LEU:C	2.38	0.49
2:B:182:LEU:HD21	2:B:187:VAL:HG22	1.95	0.48
2:B:141:ASP:HA	4:B:2019:HOH:O	2.13	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:138:ARG:HD2	2:B:172:LEU:O	2.14	0.48
1:A:135:ILE:HD13	1:A:149:VAL:HG22	1.97	0.46
2:B:125:THR:OG1	2:B:130:GLN:NE2	2.49	0.46
1:A:176:LYS:HE3	4:A:2040:HOH:O	2.15	0.46
1:A:147:VAL:C	1:A:159:LYS:HE3	2.41	0.45
1:A:138:ARG:HD3	1:A:172:LEU:O	2.17	0.45
1:A:124:GLN:HE21	2:B:260:THR:CG2	2.29	0.45
1:A:223:LEU:HD11	1:A:229:LEU:HB2	1.98	0.45
2:B:210:ILE:HG23	2:B:217:CYS:SG	2.58	0.44
1:A:124:GLN:HE21	2:B:260:THR:HG21	1.82	0.44
1:A:197:GLN:HG3	4:A:2053:HOH:O	2.17	0.44
1:A:196:ALA:O	1:A:202:GLN:NE2	2.48	0.43
1:A:260:THR:OG1	2:B:124:GLN:HB3	2.19	0.43
1:A:139:ASP:OD1	1:A:139:ASP:C	2.63	0.42
2:B:169:ARG:HG2	2:B:170:LEU:HD12	2.01	0.41
1:A:223:LEU:HB2	1:A:227:GLN:HB2	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	136/140 (97%)	134 (98%)	2 (2%)	0	100	100
2	B	135/140 (96%)	134 (99%)	1 (1%)	0	100	100
All	All	271/280 (97%)	268 (99%)	3 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	109/113 (96%)	106 (97%)	3 (3%)	38	21
2	B	110/113 (97%)	108 (98%)	2 (2%)	51	39
All	All	219/226 (97%)	214 (98%)	5 (2%)	44	28

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	125	THR
1	A	162	ILE
1	A	202	GLN
2	B	159	LYS
2	B	162	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	GLN
1	A	216	GLN
2	B	124	GLN
2	B	130	GLN
2	B	140	HIS
2	B	191	GLN
2	B	243	GLN
2	B	246	ASN
2	B	252	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MOO	A	1261	-	2,4,4	3.29	1 (50%)	-		
3	MOO	B	1261	-	2,4,4	1.43	0	-		

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1261	MOO	O2-MO	-4.39	1.65	1.75

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1261	MOO	4	0
3	B	1261	MOO	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.