



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:31 PM UTC

PDB ID : 2H9S / pdb_00002h9s
Title : Crystal Structure of Homo-DNA and Nature's Choice of Pentose over Hexose in the Genetic System
Authors : Egli, M.; Pallan, P.S.
Deposited on : 2006-06-11
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3299 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	8	 62% 38%
1	B	8	 75% 25%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 437 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*(XCT)P*(XGU)P*(XAD)P*(XAD)P*(XTH)P*(XTH)P*(XCT)P*(XGU))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	8	Total	C	N	O	P	0	0	0
			169	86	30	46	7			
1	B	8	Total	C	N	O	P	0	3	0
			195	97	32	57	9			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	30	Total	O	1	0
			30	30		
3	B	42	Total	O	0	0
			42	42		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*(XCT)P*(XGU)P*(XAD)P*(XAD)P*(XTH)P*(XTH)P*(XCT)P*(XGU))-3'

Chain A:  62% 38%

 G1 G2 A3 A4 T5 T6 C7 G8

- Molecule 1: 5'-D(*(XCT)P*(XGU)P*(XAD)P*(XAD)P*(XTH)P*(XTH)P*(XCT)P*(XGU))-3'

Chain B:  75% 25%

 C9 G10 A11 A12 T13 T14 C15 G16

4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	38.94Å 38.94Å 133.85Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 1.75 20.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.5 (20.00-1.75) 96.8 (20.00-1.75)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.62 (at 1.74Å)	Xtriage
Refinement program	REFMAC 5.0, CNS	Depositor
R, R_{free}	0.240 , 0.280 0.240 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtriage
Anisotropy	0.139	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 55.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	437	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, XGU, XCT, XTH, XAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	169	0	107	2	0
1	B	195	0	100	2	0
2	A	1	0	0	0	0
3	A	30	0	0	0	0
3	B	42	0	0	1	0
All	All	437	0	207	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:10:XGU:OP2	3:B:57:HOH:O	2.15	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:5:XTH:H3'2	1:A:6:XTH:OP2	2.14	0.47
1:A:6:XTH:H2'1	1:A:7:XCT:C2	2.47	0.45

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

19 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	XAD	B	12	1	21,24,25	0.55	0	26,34,37	1.27	3 (11%)
1	XCT	B	15[B]	1	18,21,22	0.98	1 (5%)	21,29,32	1.31	4 (19%)
1	XGU	A	8	1	22,25,26	0.91	1 (4%)	28,36,39	1.83	8 (28%)
1	XCT	A	7	1	18,21,22	0.78	0	21,29,32	1.20	2 (9%)
1	XTH	B	13[A]	1	19,22,23	0.56	0	23,31,34	1.69	3 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	XAD	B	11	1	21,24,25	0.78	1 (4%)	26,34,37	1.25	3 (11%)
1	XTH	B	14[A]	1	19,22,23	0.48	0	23,31,34	1.49	5 (21%)
1	XCT	A	1	1	18,18,22	0.96	2 (11%)	22,25,32	0.86	2 (9%)
1	XCT	B	15[A]	1	18,21,22	0.90	1 (5%)	21,29,32	1.31	4 (19%)
1	XGU	B	16	1	22,25,26	1.13	2 (9%)	28,36,39	1.72	6 (21%)
1	XGU	A	2	1,2	22,25,26	0.74	0	28,36,39	1.78	8 (28%)
1	XAD	A	4	1	21,24,25	0.73	0	26,34,37	1.10	2 (7%)
1	XCT	B	9	1	18,18,22	0.48	0	22,25,32	1.17	2 (9%)
1	XGU	B	10	1	22,25,26	0.67	0	28,36,39	1.86	7 (25%)
1	XTH	A	5	1	19,22,23	0.56	0	23,31,34	1.70	7 (30%)
1	XTH	B	13[B]	1	19,22,23	0.94	1 (5%)	23,31,34	1.84	4 (17%)
1	XTH	A	6	1	19,22,23	0.67	0	23,31,34	1.71	5 (21%)
1	XTH	B	14[B]	1	19,22,23	0.49	0	23,31,34	1.74	6 (26%)
1	XAD	A	3	1	21,24,25	0.67	0	26,34,37	1.28	3 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	XAD	B	12	1	-	0/7/22/23	0/3/3/3
1	XCT	B	15[B]	1	-	2/7/22/23	0/2/2/2
1	XGU	A	8	1	-	4/7/22/23	0/3/3/3
1	XCT	A	7	1	-	0/7/22/23	0/2/2/2
1	XTH	B	13[A]	1	-	2/7/22/23	0/2/2/2
1	XAD	B	11	1	-	0/7/22/23	0/3/3/3
1	XTH	B	14[A]	1	-	2/7/22/23	1/2/2/2
1	XCT	A	1	1	-	0/6/19/23	0/2/2/2
1	XCT	B	15[A]	1	-	3/7/22/23	0/2/2/2
1	XGU	B	16	1	-	2/7/22/23	0/3/3/3
1	XGU	A	2	1,2	-	1/7/22/23	0/3/3/3
1	XAD	A	4	1	-	0/7/22/23	0/3/3/3
1	XCT	B	9	1	-	0/6/19/23	0/2/2/2
1	XGU	B	10	1	-	2/7/22/23	0/3/3/3
1	XTH	A	5	1	-	4/7/22/23	0/2/2/2
1	XTH	B	13[B]	1	-	2/7/22/23	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	XTH	A	6	1	-	1/7/22/23	0/2/2/2
1	XTH	B	14[B]	1	-	2/7/22/23	0/2/2/2
1	XAD	A	3	1	-	0/7/22/23	0/3/3/3

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	16	XGU	O6'-C6'	-3.70	1.33	1.44
1	B	13[B]	XTH	O4'-C4'	3.40	1.50	1.43
1	A	8	XGU	O6-C6	2.54	1.28	1.23
1	A	1	XCT	C1'-N1	-2.38	1.42	1.48
1	A	1	XCT	C4'-C5'	2.35	1.56	1.52
1	B	11	XAD	O4'-C4'	-2.31	1.38	1.43
1	B	15[A]	XCT	O5'-C1'	2.20	1.45	1.42
1	B	15[B]	XCT	O5'-C1'	2.20	1.45	1.42
1	B	16	XGU	O4'-C4'	-2.15	1.38	1.43

All (84) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	6	XTH	C3'-C4'-C5'	-4.67	103.44	111.03
1	B	14[B]	XTH	C3'-C4'-C5'	-4.67	103.44	111.03
1	B	10	XGU	C2-N3-C4	4.57	120.17	112.30
1	B	13[A]	XTH	C3'-C4'-C5'	-4.53	103.67	111.03
1	B	13[B]	XTH	C3'-C4'-C5'	-4.53	103.67	111.03
1	A	8	XGU	C2-N3-C4	4.20	119.54	112.30
1	A	8	XGU	C5-C4-N3	-4.09	121.89	128.39
1	B	10	XGU	C5-C4-N3	-4.06	121.92	128.39
1	A	3	XAD	C3'-C2'-C1'	-3.95	102.49	109.06
1	A	2	XGU	C2-N3-C4	3.83	118.90	112.30
1	B	13[A]	XTH	O5'-C5'-C4'	-3.79	104.93	110.10
1	B	13[B]	XTH	O5'-C5'-C4'	-3.79	104.93	110.10
1	B	16	XGU	C2-N3-C4	3.76	118.78	112.30
1	A	5	XTH	C3'-C4'-C5'	-3.70	105.02	111.03
1	B	13[B]	XTH	O4'-C4'-C3'	3.64	118.90	109.99
1	B	14[B]	XTH	O5'-C5'-C4'	-3.63	105.15	110.10
1	B	16	XGU	C5-C4-N3	-3.52	122.79	128.39
1	B	14[A]	XTH	C3'-C4'-C5'	-3.36	105.58	111.03
1	B	12	XAD	C3'-C2'-C1'	-3.35	103.49	109.06
1	A	7	XCT	C3'-C4'-C5'	-3.26	105.73	111.03
1	A	2	XGU	C5-C4-N3	-3.26	123.20	128.39
1	A	6	XTH	O5'-C5'-C4'	-3.22	105.71	110.10

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	XAD	C3'-C2'-C1'	-3.11	103.89	109.06
1	A	5	XTH	O5'-C5'-C4'	-3.06	105.92	110.10
1	B	12	XAD	C3'-C4'-C5'	-3.04	106.09	111.03
1	B	11	XAD	C3'-C2'-C1'	-3.02	104.04	109.06
1	A	8	XGU	N9-C4-N3	2.94	131.83	125.95
1	B	10	XGU	N9-C4-N3	2.85	131.65	125.95
1	A	5	XTH	O4'-C4'-C3'	2.85	116.95	109.99
1	A	8	XGU	N9-C8-N7	-2.77	108.26	113.40
1	B	14[A]	XTH	O5'-C5'-C4'	-2.76	106.34	110.10
1	B	15[A]	XCT	C3'-C4'-C5'	-2.73	106.59	111.03
1	B	15[B]	XCT	C3'-C4'-C5'	-2.73	106.59	111.03
1	B	14[B]	XTH	O4-C4-C5	-2.69	121.84	124.92
1	A	8	XGU	C8-N7-C5	2.61	108.91	104.26
1	B	10	XGU	N9-C8-N7	-2.57	108.63	113.40
1	B	14[A]	XTH	C3'-C2'-C1'	-2.57	104.79	109.06
1	B	16	XGU	C6'-C5'-C4'	-2.55	109.79	113.74
1	A	2	XGU	C6'-C5'-C4'	-2.54	109.82	113.74
1	B	16	XGU	N9-C8-N7	-2.51	108.75	113.40
1	B	9	XCT	C3'-C4'-C5'	-2.50	106.97	111.03
1	B	9	XCT	C3'-C2'-C1'	-2.46	104.97	109.06
1	A	5	XTH	C3'-C2'-C1'	-2.46	104.97	109.06
1	A	4	XAD	C3'-C4'-C5'	-2.45	107.05	111.03
1	A	8	XGU	O5'-C5'-C4'	-2.44	106.77	110.10
1	A	3	XAD	N3-C2-N1	-2.40	124.95	128.58
1	B	14[B]	XTH	C4-N3-C2	-2.39	124.21	127.34
1	B	15[A]	XCT	O5'-C5'-C6'	2.39	111.43	106.69
1	B	15[B]	XCT	O5'-C5'-C6'	2.39	111.43	106.69
1	A	2	XGU	O5'-C5'-C6'	2.39	111.43	106.69
1	B	13[A]	XTH	C3'-C2'-C1'	-2.36	105.14	109.06
1	B	13[B]	XTH	C3'-C2'-C1'	-2.36	105.14	109.06
1	A	2	XGU	C2-N1-C6	-2.34	120.88	125.11
1	A	5	XTH	C4-N3-C2	-2.32	124.29	127.34
1	A	2	XGU	N9-C8-N7	-2.32	109.09	113.40
1	A	6	XTH	O5'-C5'-C6'	2.30	111.26	106.69
1	A	8	XGU	C2-N1-C6	-2.30	120.94	125.11
1	A	5	XTH	O5'-C1'-N1	-2.29	103.28	105.77
1	B	11	XAD	C3'-C4'-C5'	-2.29	107.32	111.03
1	B	10	XGU	C2-N1-C6	-2.26	121.00	125.11
1	B	16	XGU	N9-C4-N3	2.25	130.44	125.95
1	B	14[B]	XTH	C3'-C2'-C1'	-2.22	105.36	109.06
1	A	1	XCT	C3'-C2'-C1'	-2.19	105.41	109.06
1	B	14[A]	XTH	C4-N3-C2	-2.19	124.47	127.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	XGU	O6-C6-C5	-2.18	120.79	126.53
1	A	2	XGU	N9-C4-N3	2.17	130.29	125.95
1	B	15[A]	XCT	C6'-C5'-C4'	-2.16	110.40	113.74
1	B	15[B]	XCT	C6'-C5'-C4'	-2.16	110.40	113.74
1	B	10	XGU	O6-C6-C5	-2.16	120.84	126.53
1	A	3	XAD	O5'-C1'-N9	-2.16	103.43	105.77
1	A	6	XTH	O4'-C4'-C3'	2.15	115.25	109.99
1	B	11	XAD	N3-C2-N1	-2.15	125.33	128.58
1	B	16	XGU	O6-C6-C5	-2.14	120.88	126.53
1	B	12	XAD	N3-C2-N1	-2.14	125.34	128.58
1	A	5	XTH	O4-C4-C5	-2.13	122.48	124.92
1	B	14[A]	XTH	O4-C4-C5	-2.12	122.49	124.92
1	B	15[A]	XCT	C3'-C2'-C1'	-2.12	105.54	109.06
1	B	15[B]	XCT	C3'-C2'-C1'	-2.12	105.54	109.06
1	A	6	XTH	C4-N3-C2	-2.12	124.56	127.34
1	B	14[B]	XTH	O4'-C4'-C3'	2.06	115.03	109.99
1	B	10	XGU	C2'-C3'-C4'	-2.05	108.74	111.76
1	A	1	XCT	C3'-C4'-C5'	-2.05	107.69	111.03
1	A	8	XGU	O6-C6-C5	-2.04	121.15	126.53
1	A	7	XCT	C3'-C2'-C1'	-2.02	105.70	109.06

There are no chirality outliers.

All (27) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	5	XTH	C2'-C1'-N1-C2
1	A	5	XTH	C2'-C1'-N1-C6
1	A	8	XGU	C2'-C1'-N9-C4
1	B	10	XGU	C2'-C1'-N9-C4
1	B	14[A]	XTH	C4'-C5'-C6'-O6'
1	B	14[A]	XTH	O5'-C5'-C6'-O6'
1	B	14[B]	XTH	C4'-C5'-C6'-O6'
1	B	14[B]	XTH	O5'-C5'-C6'-O6'
1	B	15[A]	XCT	O5'-C5'-C6'-O6'
1	B	15[A]	XCT	C4'-C5'-C6'-O6'
1	B	15[B]	XCT	O5'-C5'-C6'-O6'
1	B	15[B]	XCT	C4'-C5'-C6'-O6'
1	B	16	XGU	O5'-C5'-C6'-O6'
1	B	16	XGU	C4'-C5'-C6'-O6'
1	A	8	XGU	C4'-C5'-C6'-O6'
1	A	8	XGU	O5'-C5'-C6'-O6'
1	B	13[A]	XTH	C2'-C1'-N1-C2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
1	B	13[B]	XTH	C2'-C1'-N1-C2
1	B	15[A]	XCT	C5'-C6'-O6'-P
1	A	6	XTH	O5'-C5'-C6'-O6'
1	A	5	XTH	O5'-C1'-N1-C6
1	A	2	XGU	C2'-C1'-N9-C8
1	A	8	XGU	C2'-C1'-N9-C8
1	B	10	XGU	C2'-C1'-N9-C8
1	A	5	XTH	O5'-C1'-N1-C2
1	B	13[A]	XTH	O5'-C1'-N1-C2
1	B	13[B]	XTH	O5'-C1'-N1-C2

All (1) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	14[A]	XTH	C1'-C2'-C3'-C4'-C5'-O5'

5 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	7	XCT	1	0
1	B	16	XGU	1	0
1	B	10	XGU	1	0
1	A	5	XTH	1	0
1	A	6	XTH	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	0/8	-	-	-	-
1	B	0/8	-	-	-	-
All	All	0/16	-	-	-	-

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	XTH	B	14[A]	21/22	0.57	0.37	64,78,83,85	21
1	XTH	B	14[B]	21/22	0.57	0.37	39,49,58,59	21
1	XCT	B	15[A]	20/21	0.82	0.17	31,43,68,68	4
1	XCT	B	15[B]	20/21	0.82	0.17	31,43,62,62	4
1	XCT	A	1	17/21	0.84	0.11	31,35,39,40	0
1	XTH	A	6	21/22	0.84	0.14	38,46,56,58	0
1	XGU	A	2	23/24	0.90	0.10	28,30,33,37	0
1	XCT	B	9	17/21	0.90	0.10	28,31,35,39	0
1	XAD	A	4	22/23	0.91	0.10	34,38,46,50	0
1	XCT	A	7	20/21	0.92	0.09	31,33,48,50	0
1	XTH	A	5	21/22	0.92	0.10	37,46,50,56	0
1	XAD	A	3	22/23	0.92	0.11	29,35,39,44	0
1	XTH	B	13[A]	21/22	0.93	0.10	40,45,56,61	1
1	XTH	B	13[B]	21/22	0.93	0.10	40,45,56,58	1
1	XGU	B	10	23/24	0.93	0.09	26,31,35,37	0
1	XGU	B	16	23/24	0.93	0.10	31,35,44,48	0
1	XAD	B	12	22/23	0.94	0.08	31,36,41,42	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	XGU	A	8	23/24	0.95	0.07	30,34,42,48	0
1	XAD	B	11	22/23	0.96	0.08	28,32,36,37	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MG	A	200	1/1	0.73	0.14	30,30,30,30	1

6.5 Other polymers [i](#)

There are no such residues in this entry.