



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 10:15 AM UTC

PDB ID : 5H9W / pdb_00005h9w
Title : Crystal structure of Regnase PIN domain, form II
Authors : Yokogawa, M.; Tsushima, T.; Adachi, W.; Noda, N.N.; Inagaki, F.
Deposited on : 2015-12-29
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	210	4% 63% 10% • 23%
1	B	210	64% 11% • 23%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2694 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonuclease ZC3H12A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	162	Total	C	N	O	S	0	0	0
			1328	847	239	237	5			
1	B	162	Total	C	N	O	S	0	0	0
			1330	847	241	237	5			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	130	GLY	-	expression tag	UNP Q5D1E7
A	131	PRO	-	expression tag	UNP Q5D1E7
A	132	HIS	-	expression tag	UNP Q5D1E7
A	133	MET	-	expression tag	UNP Q5D1E7
B	130	GLY	-	expression tag	UNP Q5D1E7
B	131	PRO	-	expression tag	UNP Q5D1E7
B	132	HIS	-	expression tag	UNP Q5D1E7
B	133	MET	-	expression tag	UNP Q5D1E7

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	15	Total	O	0	0
			15	15		
3	B	19	Total	O	0	0
			19	19		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Ribonuclease ZC3H12A



• Molecule 1: Ribonuclease ZC3H12A



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	113.33Å 113.33Å 78.37Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	26.94 – 2.60 26.94 – 2.60	Depositor EDS
% Data completeness (in resolution range)	92.7 (26.94-2.60) 92.8 (26.94-2.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.68 (at 2.60Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.191 , 0.231 0.200 , 0.240	Depositor DCC
R_{free} test set	863 reflections (4.73%)	wwPDB-VP
Wilson B-factor (Å ²)	54.5	Xtriage
Anisotropy	0.888	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2694	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.03	1/1360 (0.1%)	1.12	8/1837 (0.4%)
1	B	1.08	4/1362 (0.3%)	1.09	5/1840 (0.3%)
All	All	1.06	5/2722 (0.2%)	1.10	13/3677 (0.4%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	177	VAL	N-CA	-6.35	1.38	1.46
1	B	191	VAL	C-O	-6.09	1.16	1.24
1	A	264	LEU	C-O	-5.39	1.17	1.24
1	B	138	VAL	C-O	-5.36	1.18	1.24
1	B	264	LEU	C-O	-5.20	1.17	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	191	VAL	CA-C-N	-6.38	114.08	120.21
1	A	191	VAL	C-N-CA	-6.38	114.08	120.21
1	B	191	VAL	CA-C-N	-6.23	113.25	119.92
1	B	191	VAL	C-N-CA	-6.23	113.25	119.92
1	A	190	ASP	CA-C-N	-6.04	118.24	122.59
1	A	190	ASP	C-N-CA	-6.04	118.24	122.59
1	A	191	VAL	CB-CA-C	5.74	116.85	110.71
1	A	211	THR	CB-CA-C	5.47	117.35	109.09
1	B	175	ILE	CB-CA-C	-5.41	101.73	110.28
1	A	191	VAL	CA-C-O	5.37	122.32	119.15
1	A	148	SER	CB-CA-C	-5.16	102.08	110.85
1	B	187	PRO	CB-CA-C	-5.16	104.63	111.23
1	B	211	THR	CB-CA-C	5.06	116.42	108.63

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1328	0	1325	14	0
1	B	1330	0	1325	9	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	15	0	0	0	0
3	B	19	0	0	0	0
All	All	2694	0	2650	23	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:226:ASP:HB2	1:A:249:LEU:HD11	1.79	0.64
1:A:186:GLN:N	1:A:187:PRO:HD3	2.14	0.61
1:A:294:LYS:C	1:A:295:LYS:HG3	2.28	0.59
1:A:186:GLN:N	1:A:187:PRO:CD	2.72	0.52
1:B:245:THR:O	1:B:250:GLN:NE2	2.43	0.51
1:B:184:LYS:HD3	1:B:210:PHE:CE2	2.46	0.50
1:A:200:ARG:HD2	1:A:200:ARG:O	2.12	0.50
1:A:254:GLN:HE21	1:A:254:GLN:HA	1.79	0.47
1:A:294:LYS:C	1:A:295:LYS:CG	2.88	0.47
1:B:151:ASN:HD22	1:B:153:GLU:H	1.64	0.44
1:A:185:GLU:C	1:A:187:PRO:HD3	2.43	0.44
1:B:182:TRP:HA	1:B:185:GLU:HG2	1.99	0.43
1:B:215:ARG:NH1	1:B:218:GLY:O	2.49	0.43
1:B:168:LEU:HD13	1:B:175:ILE:CD1	2.49	0.43
1:A:250:GLN:HB2	1:A:257:LYS:HG3	2.02	0.41
1:A:168:LEU:HD13	1:A:175:ILE:CD1	2.50	0.41
1:A:283:ARG:HH11	1:A:283:ARG:HA	1.85	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:ASP:CB	1:A:283:ARG:HH12	2.33	0.41
1:B:250:GLN:HB2	1:B:257:LYS:HG3	2.02	0.41
1:A:254:GLN:HE21	1:A:254:GLN:CA	2.34	0.41
1:A:168:LEU:HD12	1:A:168:LEU:HA	1.90	0.40
1:B:168:LEU:HD12	1:B:168:LEU:HA	1.96	0.40
1:B:227:ARG:O	1:B:231:LYS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	160/210 (76%)	154 (96%)	5 (3%)	1 (1%)	21	42
1	B	160/210 (76%)	155 (97%)	5 (3%)	0	100	100
All	All	320/420 (76%)	309 (97%)	10 (3%)	1 (0%)	36	58

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	186	GLN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	148/192 (77%)	131 (88%)	17 (12%)	5	11
1	B	148/192 (77%)	138 (93%)	10 (7%)	14	32
All	All	296/384 (77%)	269 (91%)	27 (9%)	9	19

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	134	ASP
1	A	168	LEU
1	A	185	GLU
1	A	200	ARG
1	A	211	THR
1	A	215	ARG
1	A	219	LYS
1	A	225	ASP
1	A	226	ASP
1	A	231	LYS
1	A	247	ARG
1	A	254	GLN
1	A	255	GLU
1	A	258	ARG
1	A	283	ARG
1	A	288	LEU
1	A	295	LYS
1	B	134	ASP
1	B	151	ASN
1	B	168	LEU
1	B	204	LYS
1	B	211	THR
1	B	225	ASP
1	B	232	LEU
1	B	258	ARG
1	B	283	ARG
1	B	288	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	151	ASN
1	A	254	GLN
1	B	151	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	254	GLN
1	B	271	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	162/210 (77%)	0.11	8 (4%) 35 29	47, 72, 112, 147	0
1	B	162/210 (77%)	-0.06	1 (0%) 85 83	44, 69, 109, 127	0
All	All	324/420 (77%)	0.03	9 (2%) 55 49	44, 70, 110, 147	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	295	LYS	3.2
1	A	231	LYS	3.1
1	A	218	GLY	3.0
1	A	295	LYS	2.9
1	A	223	CYS	2.9
1	A	187	PRO	2.7
1	A	189	PRO	2.7
1	A	224	TYR	2.5
1	A	252	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NA	A	401	1/1	0.98	0.05	61,61,61,61	0
2	NA	B	401	1/1	0.98	0.04	61,61,61,61	0

6.5 Other polymers [i](#)

There are no such residues in this entry.