



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:51 PM UTC

PDB ID : 6H9F / pdb_00006h9f
Title : Structure of glutamate mutase reconstituted with bishomo-coenzyme B12
Authors : Gruber, K.; Csitkovits, V.; Kratky, C.
Deposited on : 2018-08-03
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	137	91% 8% .
1	C	137	2% 91% 9% .
2	B	483	89% 10% .
2	D	483	92% 8%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11200 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate mutase sigma subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	137	Total	C	N	O	S	0	0	0
			1041	661	176	199	5			
1	C	137	Total	C	N	O	S	0	0	0
			1041	661	176	199	5			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	45	ASN	VAL	conflict	UNP P80078
A	60	VAL	LEU	conflict	UNP P80078
C	45	ASN	VAL	conflict	UNP P80078
C	60	VAL	LEU	conflict	UNP P80078


- Molecule 2 is a protein called Glutamate mutase epsilon subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	483	Total	C	N	O	S	0	0	0
			3761	2380	645	714	22			
2	D	483	Total	C	N	O	S	0	0	0
			3761	2380	645	714	22			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	130	HIS	TYR	conflict	UNP P80077
D	130	HIS	TYR	conflict	UNP P80077

- Molecule 3 is COBALAMIN (CCD ID: B12) (formula: $C_{62}H_{89}CoN_{13}O_{14}P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		
3	C	1	Total	C	Co	N	O	P	0	0
			91	62	1	13	14	1		

- Molecule 4 is (2 {R},3 {R},4 {S},5 {R})-2-(6-aminopurin-9-yl)-5-propyl-oxolane-3,4-diol (CCD ID: 8ZB) (formula: C₁₂H₁₇N₅O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	N	O	0	0
			20	12	5	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			20	12	5	3		

- Molecule 5 is D(-)-TARTARIC ACID (CCD ID: TAR) (formula: C₄H₆O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			10	4	6		
5	D	1	Total	C	O	0	0
			10	4	6		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	133	Total	O	0	0
			133	133		
6	B	548	Total	O	0	0
			548	548		
6	C	125	Total	O	0	0
			125	125		
6	D	548	Total	O	0	0
			548	548		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate mutase sigma subunit

Chain A:  91% 8% .




- Molecule 1: Glutamate mutase sigma subunit

Chain C:  2% 91% 9% .




- Molecule 2: Glutamate mutase epsilon subunit

Chain B:  89% 10% .



- Molecule 2: Glutamate mutase epsilon subunit

Chain D:  92% 8%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.75Å 112.29Å 108.01Å 90.00° 95.71° 90.00°	Depositor
Resolution (Å)	35.35 – 2.10 35.35 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (35.35-2.10) 92.0 (35.35-2.10)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.16 (at 2.10Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.158 , 0.203 0.158 , 0.202	Depositor DCC
R_{free} test set	4528 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	8.3	Xtriage
Anisotropy	0.519	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11200	wwPDB-VP
Average B, all atoms (Å ²)	13.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TAR, 8ZB, B12

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.26	0/1058	0.48	0/1427
1	C	0.27	0/1058	0.46	0/1427
2	B	0.30	0/3832	0.49	0/5170
2	D	0.30	0/3832	0.48	0/5170
All	All	0.29	0/9780	0.48	0/13194

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	D	0	1
All	All	0	2

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	92	PRO	Peptide
2	D	92	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1041	0	1052	7	0
1	C	1041	0	1052	7	0
2	B	3761	0	3760	28	0
2	D	3761	0	3760	24	0
3	A	91	0	88	3	0
3	C	91	0	88	6	0
4	B	20	0	0	0	0
4	D	20	0	0	1	0
5	B	10	0	4	0	0
5	D	10	0	4	0	0
6	A	133	0	0	0	0
6	B	548	0	0	2	0
6	C	125	0	0	0	0
6	D	548	0	0	5	0
All	All	11200	0	9808	71	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:55:ALA:HB2	2:B:62:MET:HE2	1.57	0.84
2:D:55:ALA:HB2	2:D:62:MET:HE2	1.69	0.73
2:D:51:LYS:HG2	2:D:62:MET:HE1	1.76	0.68
2:D:33:GLN:NE2	6:D:601:HOH:O	2.21	0.64
2:B:451:GLN:NE2	6:B:601:HOH:O	2.16	0.63
1:A:44:GLU:OE2	1:A:77:LYS:NZ	2.32	0.59
1:C:3:LYS:HE2	1:C:33:ASN:HB2	1.86	0.56
3:C:201:B12:H5R1	3:C:201:B12:H3P1	1.87	0.56
2:D:299:GLN:HG3	2:D:332:ILE:HD11	1.88	0.55
2:B:133:LYS:HE2	2:B:137:LYS:NZ	2.22	0.54
3:A:201:B12:H601	3:A:201:B12:H252	1.89	0.54
2:B:113:GLU:HB3	2:B:121:LEU:HD12	1.92	0.51
2:D:147:GLN:HB2	2:D:169:SER:HB3	1.92	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:96:ASP:O	2:D:100:ARG:HG3	2.11	0.51
2:B:147:GLN:HB2	2:B:169:SER:HB3	1.93	0.50
2:D:59:GLY:HA2	6:D:799:HOH:O	2.10	0.50
1:C:60:VAL:O	1:C:90:VAL:HA	2.12	0.50
2:B:299:GLN:CD	2:B:332:ILE:HD11	2.36	0.50
3:A:201:B12:H552	3:A:201:B12:H531	1.93	0.49
2:D:277:LYS:NZ	6:D:610:HOH:O	2.44	0.49
2:B:95:ILE:HB	2:B:150:HIS:HB3	1.95	0.49
2:D:139:LEU:HD22	2:D:166:GLY:HA3	1.95	0.49
2:B:51:LYS:HG2	2:B:62:MET:HE1	1.95	0.48
2:B:74:GLU:OE1	2:B:74:GLU:N	2.44	0.48
3:C:201:B12:H552	3:C:201:B12:H531	1.95	0.48
2:D:454:ALA:HB2	2:D:461:VAL:HG23	1.96	0.48
2:D:176:SER:HB2	2:D:216:PHE:HB3	1.97	0.47
2:B:138:VAL:HG12	2:B:146:LEU:HD11	1.97	0.47
1:A:10:VAL:HB	1:A:39:VAL:HA	1.97	0.47
2:B:454:ALA:HB2	2:B:461:VAL:HG23	1.97	0.47
3:C:201:B12:H253	3:C:201:B12:H301	1.80	0.46
2:B:183:LYS:HA	2:B:411:SER:HA	1.96	0.46
2:D:116:LYS:HE2	6:D:1025:HOH:O	2.16	0.46
3:C:201:B12:H352	2:D:334:ILE:HD11	1.98	0.45
2:D:94:THR:HG21	4:D:501:8ZB:N48	2.32	0.45
1:C:64:TYR:HB2	3:C:201:B12:C36	2.47	0.45
2:B:94:THR:HA	2:B:149:ARG:O	2.16	0.45
3:C:201:B12:H601	3:C:201:B12:H252	1.99	0.45
2:B:66:ARG:CZ	2:B:324:ILE:HG21	2.47	0.44
2:D:94:THR:HA	2:D:149:ARG:O	2.16	0.44
2:B:259:ILE:HG21	2:B:364:GLU:HG2	1.99	0.44
2:B:136:ARG:HG3	2:B:165:GLY:HA2	1.98	0.44
2:B:482:PRO:O	2:B:483:GLU:HB2	2.18	0.44
2:D:62:MET:HE3	2:D:90:PHE:CE2	2.52	0.44
2:B:221:GLY:HA2	6:B:975:HOH:O	2.17	0.43
2:D:265:LEU:HD11	2:D:288:THR:HB	2.00	0.43
2:B:133:LYS:HE2	2:B:137:LYS:HZ2	1.81	0.43
2:B:216:PHE:CD1	2:B:218:PRO:HD2	2.54	0.43
2:D:219:LEU:HD13	2:D:406:ILE:HD12	1.99	0.43
1:C:64:TYR:CZ	1:C:66:GLN:HB2	2.54	0.43
2:D:51:LYS:HG2	2:D:62:MET:CE	2.48	0.43
2:B:429:ARG:HB2	2:B:450:LEU:HD21	2.01	0.42
2:D:56:LYS:HG2	2:D:287:THR:HG22	2.00	0.42
2:D:297:PHE:CD1	2:D:330:GLU:HG2	2.53	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:1:MET:HE2	6:D:907:HOH:O	2.18	0.42
2:B:292:GLN:O	2:B:293:TRP:C	2.62	0.42
2:B:262:ILE:HG12	2:B:318:ALA:HB2	2.02	0.42
1:A:51:ILE:HD11	1:A:81:ALA:HB1	2.01	0.42
1:C:64:TYR:OH	1:C:70:ASP:OD2	2.34	0.42
1:A:64:TYR:HB2	3:A:201:B12:C36	2.49	0.41
2:D:136:ARG:HG3	2:D:165:GLY:HA2	2.02	0.41
2:D:167:TRP:N	2:D:167:TRP:CD1	2.88	0.41
1:A:60:VAL:O	1:A:90:VAL:HA	2.21	0.41
2:B:167:TRP:CD1	2:B:167:TRP:N	2.88	0.41
2:B:309:VAL:HG13	2:B:350:ALA:HB2	2.03	0.41
1:C:47:ILE:CD1	1:C:77:LYS:HB3	2.50	0.41
2:B:59:GLY:O	2:B:358:ARG:NH2	2.52	0.41
1:A:15:CYS:SG	2:B:181:TYR:HA	2.62	0.40
1:A:64:TYR:CZ	1:A:66:GLN:HB2	2.57	0.40
2:B:32:LEU:HD23	2:B:32:LEU:HA	1.93	0.40
1:C:35:VAL:HG21	1:C:53:THR:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	135/137 (98%)	131 (97%)	4 (3%)	0	100	100
1	C	135/137 (98%)	131 (97%)	4 (3%)	0	100	100
2	B	481/483 (100%)	471 (98%)	7 (2%)	3 (1%)	21	18
2	D	481/483 (100%)	471 (98%)	7 (2%)	3 (1%)	21	18
All	All	1232/1240 (99%)	1204 (98%)	22 (2%)	6 (0%)	24	22

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	293	TRP
2	B	151	GLY
2	D	151	GLY
2	D	293	TRP
2	B	150	HIS
2	D	150	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	111/111 (100%)	109 (98%)	2 (2%)	51	60
1	C	111/111 (100%)	108 (97%)	3 (3%)	39	45
2	B	396/396 (100%)	393 (99%)	3 (1%)	73	81
2	D	396/396 (100%)	395 (100%)	1 (0%)	86	91
All	All	1014/1014 (100%)	1005 (99%)	9 (1%)	70	78

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	LYS
1	A	64	TYR
2	B	62	MET
2	B	297	PHE
2	B	366	GLU
1	C	2	GLU
1	C	21	LYS
1	C	64	TYR
2	D	62	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
2	B	22	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	143	ASN
2	B	340	ASN
2	B	464	GLN
2	D	22	GLN
2	D	64	GLN
2	D	212	ASN
2	D	247	ASN
2	D	340	ASN
2	D	444	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	8ZB	B	501	3	22,22,22	1.55	5 (22%)	32,32,32	2.05	8 (25%)
5	TAR	D	502	-	9,9,9	1.28	0	12,12,12	1.34	2 (16%)
5	TAR	B	502	-	9,9,9	1.22	0	12,12,12	1.43	2 (16%)
3	B12	C	201	4,1	94,101,101	0.82	2 (2%)	149,166,166	1.16	13 (8%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	B12	A	201	4,1	94,101,101	0.81	3 (3%)	149,166,166	1.19	11 (7%)
4	8ZB	D	501	3	22,22,22	1.48	5 (22%)	32,32,32	2.13	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	8ZB	B	501	3	-	1/7/23/23	0/3/3/3
5	TAR	D	502	-	-	4/12/12/12	-
5	TAR	B	502	-	-	4/12/12/12	-
3	B12	C	201	4,1	-	22/56/223/223	0/3/11/11
3	B12	A	201	4,1	-	19/56/223/223	0/3/11/11
4	8ZB	D	501	3	-	1/7/23/23	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	501	8ZB	C49-N48	4.15	1.39	1.31
4	B	501	8ZB	C49-N48	3.74	1.38	1.31
4	B	501	8ZB	C53-N54	3.49	1.40	1.33
4	B	501	8ZB	C51-N52	2.90	1.39	1.34
4	D	501	8ZB	C53-N54	2.83	1.39	1.33
4	B	501	8ZB	C53-N52	2.73	1.38	1.33
4	D	501	8ZB	C53-N52	2.68	1.38	1.33
4	B	501	8ZB	C55-N57	2.55	1.40	1.34
4	D	501	8ZB	C55-N57	2.50	1.40	1.34
4	D	501	8ZB	C51-N52	2.32	1.38	1.34
3	A	201	B12	C53-C15	2.19	1.55	1.50
3	C	201	B12	C53-C15	2.19	1.55	1.50
3	A	201	B12	C2B-N1B	-2.17	1.33	1.37
3	A	201	B12	C35-C5	2.16	1.55	1.50
3	C	201	B12	C2B-N1B	-2.13	1.33	1.37

All (44) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	501	8ZB	N54-C53-N52	-5.71	119.94	128.58
4	B	501	8ZB	N54-C53-N52	-5.64	120.05	128.58
4	D	501	8ZB	N47-C49-N48	-5.23	106.51	113.94

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	201	B12	C9-N22-C6	4.62	110.83	105.28
4	B	501	8ZB	N47-C49-N48	-4.58	107.44	113.94
3	C	201	B12	C9-N22-C6	4.09	110.20	105.28
4	B	501	8ZB	C50-C51-N52	-4.05	121.14	126.72
3	A	201	B12	C30-C3-C4	3.79	118.54	109.66
4	D	501	8ZB	C50-C51-N52	-3.77	121.53	126.72
4	D	501	8ZB	C50-N48-C49	3.53	109.00	103.45
4	D	501	8ZB	C51-N47-C49	3.21	109.10	105.74
4	B	501	8ZB	C51-N47-C49	3.18	109.08	105.74
5	B	502	TAR	C2-C3-C4	-3.06	103.04	109.82
3	C	201	B12	C36-C7-C37	2.91	115.68	110.74
4	D	501	8ZB	C51-C50-N48	-2.90	107.27	110.58
3	C	201	B12	C2-C3-C4	2.90	104.90	101.64
3	C	201	B12	C30-C3-C4	2.87	116.39	109.66
4	B	501	8ZB	C50-N48-C49	2.83	107.90	103.45
4	D	501	8ZB	C53-N52-C51	2.83	118.73	111.83
4	B	501	8ZB	N52-C51-N47	2.79	131.91	127.17
3	A	201	B12	C36-C7-C37	2.73	115.38	110.74
4	B	501	8ZB	C53-N52-C51	2.67	118.34	111.83
3	A	201	B12	C13-C12-C11	-2.61	98.05	100.97
3	C	201	B12	C18-C19-N24	2.56	106.18	102.33
5	D	502	TAR	O41-C4-C3	2.56	120.43	113.31
3	A	201	B12	C19-N24-C16	2.56	110.08	107.29
3	A	201	B12	C2-C3-C4	2.50	104.46	101.64
3	A	201	B12	C2P-C1P-N59	-2.49	109.25	112.92
3	A	201	B12	C8-C7-C6	2.48	105.12	100.92
3	C	201	B12	C2R-C3R-C4R	2.46	107.54	103.24
3	A	201	B12	C2R-C3R-C4R	2.45	107.53	103.24
3	C	201	B12	C12-C11-N23	-2.44	108.46	111.83
3	C	201	B12	C54-C17-C18	-2.41	109.53	112.99
5	D	502	TAR	O11-C1-C2	2.38	119.93	113.31
4	B	501	8ZB	C55-C50-C51	2.33	120.36	117.18
3	C	201	B12	C7B-C8B-C9B	2.20	125.11	122.47
3	C	201	B12	C15-C16-N24	-2.16	119.34	122.42
3	A	201	B12	C8-C9-C10	-2.14	118.76	123.33
3	C	201	B12	C19-N24-C16	2.13	109.62	107.29
3	A	201	B12	C12-C11-N23	-2.13	108.89	111.83
4	D	501	8ZB	C50-C51-N47	2.12	108.12	105.81
3	C	201	B12	C8-C9-C10	-2.12	118.83	123.33
5	B	502	TAR	O41-C4-C3	2.03	118.95	113.31
3	C	201	B12	C4B-C5B-C6B	2.00	122.63	119.69

There are no chirality outliers.

All (51) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	201	B12	C38-C37-C7-C6
3	A	201	B12	C38-C37-C7-C36
3	A	201	B12	C38-C37-C7-C8
3	A	201	B12	C1P-C2P-O3-P
3	A	201	B12	C3P-C2P-O3-P
3	C	201	B12	C38-C37-C7-C6
3	C	201	B12	C38-C37-C7-C36
3	C	201	B12	C38-C37-C7-C8
3	C	201	B12	C1P-C2P-O3-P
3	C	201	B12	C3P-C2P-O3-P
3	C	201	B12	C2P-O3-P-O4
3	C	201	B12	C2P-O3-P-O2
3	C	201	B12	C3R-O2-P-O3
5	D	502	TAR	O2-C2-C3-C4
3	A	201	B12	C4-C3-C30-C31
3	C	201	B12	C4-C3-C30-C31
5	D	502	TAR	O2-C2-C3-O3
4	B	501	8ZB	C24-C28-C31-C36
4	D	501	8ZB	C24-C28-C31-C36
5	D	502	TAR	C1-C2-C3-O3
5	B	502	TAR	C1-C2-C3-O3
3	A	201	B12	C8-C41-C42-C43
3	C	201	B12	C8-C41-C42-C43
3	C	201	B12	C2-C3-C30-C31
5	D	502	TAR	C1-C2-C3-C4
5	B	502	TAR	O2-C2-C3-O3
3	A	201	B12	C2R-C1R-N1B-C2B
5	B	502	TAR	C1-C2-C3-C4
5	B	502	TAR	O2-C2-C3-C4
3	C	201	B12	C2R-C1R-N1B-C2B
3	C	201	B12	C18-C60-C61-O63
3	C	201	B12	C18-C60-C61-N62
3	A	201	B12	C2R-C1R-N1B-C8B
3	A	201	B12	O6R-C1R-N1B-C8B
3	C	201	B12	C2R-C1R-N1B-C8B
3	C	201	B12	O6R-C1R-N1B-C8B
3	A	201	B12	C2-C3-C30-C31
3	A	201	B12	C18-C60-C61-O63
3	C	201	B12	O6R-C1R-N1B-C2B
3	C	201	B12	C55-C56-C57-O58
3	A	201	B12	C18-C60-C61-N62
3	A	201	B12	C2P-O3-P-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	C	201	B12	C2P-O3-P-O5
3	C	201	B12	C3R-O2-P-O4
3	A	201	B12	C30-C31-C32-N33
3	A	201	B12	C55-C56-C57-O58
3	A	201	B12	O6R-C1R-N1B-C2B
3	A	201	B12	C55-C56-C57-N59
3	C	201	B12	C55-C56-C57-N59
3	C	201	B12	C30-C31-C32-N33
3	A	201	B12	C30-C31-C32-O34


There are no ring outliers.

3 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	201	B12	6	0
3	A	201	B12	3	0
4	D	501	8ZB	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 8ZB B 501



Ligand B12 C 201





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	137/137 (100%)	-0.46	0 100 100	7, 15, 32, 51	0
1	C	137/137 (100%)	-0.24	3 (2%) 62 65	7, 16, 35, 66	0
2	B	483/483 (100%)	-0.75	1 (0%) 91 92	4, 9, 20, 56	0
2	D	483/483 (100%)	-0.73	1 (0%) 91 92	4, 9, 20, 65	0
All	All	1240/1240 (100%)	-0.65	5 (0%) 88 90	4, 10, 25, 66	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	MET	4.7
2	D	483	GLU	3.1
2	B	483	GLU	2.5
1	C	2	GLU	2.3
1	C	137	GLU	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	8ZB	D	501	20/20	0.97	0.05	4,8,9,11	0
5	TAR	B	502	10/10	0.97	0.05	6,6,8,17	0
4	8ZB	B	501	20/20	0.98	0.05	5,8,10,13	0
3	B12	A	201	91/91	0.98	0.05	3,6,11,14	0
3	B12	C	201	91/91	0.98	0.05	2,6,14,19	0
5	TAR	D	502	10/10	0.98	0.05	5,6,7,15	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 8ZB D 501:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 8ZB B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around B12 A 201:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around B12 C 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.