



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 06:22 AM UTC

PDB ID : 2HAW / pdb_00002haw
Title : Crystal structure of family II Inorganic pyrophosphatase in complex with PNP
Authors : Fabrichniy, I.P.; Lehtio, L.; Oksanen, E.; Goldman, A.
Deposited on : 2006-06-13
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	309	28% 95% • •
1	B	309	28% 93% 6% •

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 5483 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Manganese-dependent inorganic pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	307	Total	C	N	O	S	0	8	0
			2431	1533	403	488	7			
1	B	308	Total	C	N	O	S	0	16	0
			2507	1575	413	511	8			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	4	Total	Mg	0	0
			4	4		
2	B	4	Total	Mg	0	0
			4	4		

- Molecule 3 is FLUORIDE ION (CCD ID: F) (formula: F).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	F	0	0
			1	1		
3	B	1	Total	F	0	0
			1	1		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	1
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	1
			5	4	1		

- Molecule 5 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		
6	B	1	Total	C	O	0	0
			13	8	5		

- Molecule 7 is IMIDODIPHOSPHORIC ACID (CCD ID: 2PN) (formula: $\text{H}_5\text{NO}_6\text{P}_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	N	O	P	0	1
			9	1	6	2		
7	B	1	Total	N	O	P	0	1
			9	1	6	2		

- Molecule 8 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	B	1	Total	C	O	0	0
			16	10	6		

- Molecule 9 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			6	3	3		
9	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	166	Total	O	0	0
			166	166		
10	B	270	Total	O	0	0
			270	270		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Manganese-dependent inorganic pyrophosphatase



- Molecule 1: Manganese-dependent inorganic pyrophosphatase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	60.01Å 115.87Å 147.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.84 – 1.75 19.84 – 1.75	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.84-1.75) 99.9 (19.84-1.75)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.45 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.166 , 0.186 (Not available) , 0.294	Depositor DCC
R_{free} test set	5228 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.021	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5483	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.38% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 1PE, CL, GOL, PG4, SO4, MG, F, 2PN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.73	1/2456 (0.0%)	0.86	0/3324
1	B	0.95	1/2532 (0.0%)	0.94	2/3426 (0.1%)
All	All	0.85	2/4988 (0.0%)	0.90	2/6750 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	51	ASP	CG-OD1	7.49	1.39	1.25
1	B	300	VAL	CA-CB	5.70	1.57	1.54

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	191	LEU	N-CA-C	6.86	124.31	114.16
1	B	227	VAL	N-CA-C	-5.50	107.12	111.86

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2431	0	2472	6	0
1	B	2507	0	2530	7	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	4	0	0	0	0
2	B	4	0	0	0	0
3	A	1	0	0	1	0
3	B	1	0	0	0	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
5	A	1	0	0	0	0
6	A	13	0	18	1	0
6	B	13	0	18	0	0
7	A	9	0	1	0	0
7	B	9	0	1	0	0
8	B	16	0	22	1	0
9	B	18	0	24	1	0
10	A	166	0	0	0	0
10	B	270	0	0	0	0
All	All	5483	0	5086	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:28:ASN:HD21	1:A:34:ALA:H	1.30	0.79
1:A:101:ALA:HA	1:B:104[B]:GLU:HG2	1.69	0.73
1:B:66:ASN:HD22	1:B:66:ASN:H	1.39	0.71
1:B:66:ASN:H	1:B:66:ASN:ND2	2.03	0.54
1:A:79:ARG:NH2	1:A:102[B]:ASN:OD1	2.42	0.53
1:B:191:LEU:C	1:B:193:LYS:H	2.18	0.51
8:B:1001:1PE:H251	8:B:1001:1PE:H122	1.94	0.49
1:A:15:ASP:HB3	3:A:3001:F:F	2.06	0.45
1:B:96:ASP:O	1:B:113:ALA:HA	2.16	0.45
1:A:133:LYS:CD	1:A:133:LYS:H	2.28	0.45
6:A:6002:PG4:H42	6:A:6002:PG4:H62	1.52	0.44
1:B:186:LYS:HB3	1:B:186:LYS:HE2	1.73	0.43
1:B:57[B]:SER:HB3	9:B:5003:GOL:H2	2.01	0.42
1:A:42:VAL:HG13	1:A:46:THR:HB	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	313/309 (101%)	305 (97%)	8 (3%)	0	100	100
1	B	322/309 (104%)	314 (98%)	8 (2%)	0	100	100
All	All	635/618 (103%)	619 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	122/262 (47%)	116 (95%)	6 (5%)	22	6
1	B	262/262 (100%)	246 (94%)	16 (6%)	17	3
All	All	384/524 (73%)	362 (94%)	22 (6%)	22	4

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	133	LYS
1	A	219[A]	VAL
1	A	219[B]	VAL
1	A	227	VAL
1	A	230	ARG
1	A	263	LEU
1	B	66	ASN
1	B	84	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	178[A]	GLU
1	B	178[B]	GLU
1	B	186	LYS
1	B	191	LEU
1	B	202[A]	LEU
1	B	202[B]	LEU
1	B	225[A]	GLU
1	B	225[B]	GLU
1	B	258	LEU
1	B	280	VAL
1	B	283	GLU
1	B	288	LEU
1	B	293	VAL
1	B	307	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	ASN
1	A	269	ASN
1	B	33	ASN
1	B	66	ASN
1	B	90	GLN
1	B	102	ASN
1	B	124	ASN
1	B	130	ASN
1	B	131	ASN
1	B	284	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 23 ligands modelled in this entry, 11 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
9	GOL	B	5002	-	5,5,5	0.39	0	5,5,5	0.80	0
6	PG4	A	6002	-	12,12,12	0.47	0	11,11,11	0.35	0
4	SO4	A	4003[B]	2	4,4,4	0.25	0	6,6,6	0.25	0
4	SO4	A	4002	-	4,4,4	0.25	0	6,6,6	0.39	0
4	SO4	B	4001	-	4,4,4	0.26	0	6,6,6	0.86	0
7	2PN	A	2001[A]	2	8,8,8	5.13	6 (75%)	8,13,13	3.07	2 (25%)
8	1PE	B	1001	-	15,15,15	0.54	0	14,14,14	1.04	1 (7%)
6	PG4	B	1002	-	12,12,12	0.63	0	11,11,11	0.48	0
4	SO4	B	4004[B]	2	4,4,4	0.27	0	6,6,6	0.07	0
9	GOL	B	5003	-	5,5,5	0.44	0	5,5,5	0.85	0
9	GOL	B	5001	-	5,5,5	0.59	0	5,5,5	1.34	1 (20%)
7	2PN	B	2002[A]	2	8,8,8	4.47	5 (62%)	8,13,13	2.29	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	GOL	B	5002	-	-	3/4/4/4	-
6	PG4	A	6002	-	-	6/10/10/10	-
7	2PN	A	2001[A]	2	-	2/2/6/6	-
8	1PE	B	1001	-	-	6/13/13/13	-
6	PG4	B	1002	-	-	2/10/10/10	-
9	GOL	B	5003	-	-	2/4/4/4	-
9	GOL	B	5001	-	-	2/4/4/4	-
7	2PN	B	2002[A]	2	-	2/2/6/6	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	B	2002[A]	2PN	P1-O1	9.43	1.60	1.46
7	A	2001[A]	2PN	P2-O4	8.37	1.58	1.46
7	A	2001[A]	2PN	P1-N1	7.74	1.83	1.63
7	A	2001[A]	2PN	P2-N1	-5.96	1.47	1.63
7	A	2001[A]	2PN	P2-O5	-4.75	1.44	1.56
7	B	2002[A]	2PN	P2-O6	-4.63	1.44	1.56
7	B	2002[A]	2PN	P2-N1	4.38	1.74	1.63
7	B	2002[A]	2PN	P2-O4	4.22	1.52	1.46
7	A	2001[A]	2PN	P1-O1	4.05	1.52	1.46
7	B	2002[A]	2PN	P1-O2	2.38	1.63	1.56
7	A	2001[A]	2PN	P1-O2	2.28	1.62	1.56

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	2001[A]	2PN	O4-P2-N1	-6.87	101.65	111.77
7	B	2002[A]	2PN	O4-P2-N1	-5.06	104.32	111.77
7	A	2001[A]	2PN	O2-P1-O1	-4.97	100.99	113.45
7	B	2002[A]	2PN	O3-P1-O1	-2.46	107.29	113.45
9	B	5001	GOL	O1-C1-C2	-2.21	100.44	110.38
8	B	1001	1PE	OH4-C24-C14	2.16	120.21	110.35

There are no chirality outliers.

All (25) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	2001[A]	2PN	P2-N1-P1-O1
7	A	2001[A]	2PN	P1-N1-P2-O4
7	B	2002[A]	2PN	P2-N1-P1-O1
7	B	2002[A]	2PN	P1-N1-P2-O4
9	B	5001	GOL	C1-C2-C3-O3
9	B	5001	GOL	O2-C2-C3-O3
9	B	5003	GOL	C1-C2-C3-O3
9	B	5003	GOL	O2-C2-C3-O3
6	A	6002	PG4	C6-C5-O3-C4
8	B	1001	1PE	C14-C24-OH4-C13
8	B	1001	1PE	OH5-C14-C24-OH4
9	B	5002	GOL	O1-C1-C2-C3
6	B	1002	PG4	O1-C1-C2-O2
9	B	5002	GOL	O1-C1-C2-O2
6	A	6002	PG4	O2-C3-C4-O3
6	A	6002	PG4	O4-C7-C8-O5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
6	B	1002	PG4	O2-C3-C4-O3
6	A	6002	PG4	C5-C6-O4-C7
8	B	1001	1PE	C23-C13-OH4-C24
8	B	1001	1PE	OH6-C15-C25-OH5
6	A	6002	PG4	O1-C1-C2-O2
6	A	6002	PG4	C8-C7-O4-C6
8	B	1001	1PE	C24-C14-OH5-C25
9	B	5002	GOL	C1-C2-C3-O3
8	B	1001	1PE	OH2-C12-C22-OH3

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	6002	PG4	1	0
8	B	1001	1PE	1	0
9	B	5003	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	307/309 (99%)	1.53	86 (28%)  	16, 41, 57, 70	8 (2%)
1	B	308/309 (99%)	1.55	86 (27%)  	17, 39, 62, 80	16 (5%)
All	All	615/618 (99%)	1.54	172 (27%)  	16, 40, 60, 80	24 (3%)

All (172) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	258	LEU	6.5
1	B	192	SER	6.0
1	B	308	ALA	5.7
1	A	308	ALA	5.5
1	B	257	ILE	5.3
1	B	188	GLY	4.7
1	B	306	ALA	4.5
1	A	202[A]	LEU	4.4
1	B	191	LEU	4.4
1	B	222	VAL	4.3
1	A	209	LEU	4.2
1	B	226[A]	ASP	4.2
1	B	260	ASN	4.0
1	A	305	ASP	3.9
1	B	31	GLY	3.9
1	B	202[A]	LEU	3.9
1	B	107[A]	GLU	3.8
1	A	32	PHE	3.8
1	A	109	LEU	3.8
1	A	108	PRO	3.7
1	B	227	VAL	3.7
1	B	221	THR	3.7
1	A	257	ILE	3.6
1	A	137	GLU	3.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	106	ALA	3.5
1	A	86	ILE	3.5
1	B	288	LEU	3.5
1	A	174	GLY	3.5
1	A	261	ASP	3.4
1	B	256	ASP	3.4
1	B	30	LEU	3.4
1	A	291	GLY	3.4
1	B	291	GLY	3.4
1	A	222	VAL	3.4
1	B	223	ASP	3.3
1	B	225[A]	GLU	3.3
1	B	132	VAL	3.3
1	B	224	ILE	3.2
1	B	189	ALA	3.2
1	B	190	ASP	3.2
1	B	229	LYS	3.2
1	A	258	LEU	3.1
1	B	290	LYS	3.1
1	A	226	ASP	3.1
1	A	182	LEU	3.1
1	A	102[A]	ASN	3.1
1	A	260	ASN	3.1
1	B	259	GLU	3.1
1	B	72	ILE	3.1
1	A	177	ALA	3.1
1	A	242	VAL	3.0
1	A	72	ILE	3.0
1	A	100	ILE	3.0
1	A	241	VAL	3.0
1	A	52	TYR	3.0
1	B	261	ASP	3.0
1	A	304	THR	3.0
1	B	282	LEU	3.0
1	B	34	ALA	2.9
1	A	247	LEU	2.9
1	B	32	PHE	2.9
1	B	24	ALA	2.9
1	A	191	LEU	2.9
1	A	188	GLY	2.9
1	B	281	THR	2.9
1	B	26	LEU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	186	LYS	2.9
1	A	105	THR	2.8
1	B	106	ALA	2.8
1	B	108	PRO	2.8
1	B	89	VAL	2.8
1	A	272	ALA	2.8
1	A	111	TYR	2.8
1	A	70	GLY	2.8
1	B	1	MET	2.7
1	A	210	GLY	2.7
1	A	2	GLU	2.7
1	A	89	VAL	2.7
1	A	221	THR	2.7
1	B	232	ALA	2.7
1	B	109	LEU	2.7
1	B	193	LYS	2.6
1	B	68	VAL	2.6
1	A	91	VAL	2.6
1	B	110	TYR	2.6
1	A	207	PHE	2.6
1	A	279	ASN	2.5
1	B	52	TYR	2.5
1	A	94	VAL	2.5
1	B	4	ILE	2.5
1	A	189	ALA	2.5
1	B	277	ALA	2.5
1	A	51	ASP	2.5
1	A	4	ILE	2.5
1	A	31	GLY	2.5
1	B	182	LEU	2.5
1	A	71	VAL	2.5
1	A	268	GLY	2.5
1	B	255	THR	2.5
1	A	282	LEU	2.4
1	A	178	GLU	2.4
1	B	64	ALA	2.4
1	B	6	ILE	2.4
1	B	91	VAL	2.4
1	A	103	PHE	2.4
1	B	287	ALA	2.4
1	B	130	ASN	2.4
1	A	50	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	151	LEU	2.4
1	B	283	GLU	2.4
1	A	48	TYR	2.4
1	B	280	VAL	2.4
1	B	33	ASN	2.3
1	A	256	ASP	2.3
1	B	209	LEU	2.3
1	A	180	TYR	2.3
1	B	220	ASN	2.3
1	B	269	ASN	2.3
1	B	194	LYS	2.3
1	B	289	LEU	2.3
1	B	21	ILE	2.3
1	A	136	LYS	2.3
1	B	22	ALA	2.3
1	B	131	ASN	2.3
1	B	111	TYR	2.3
1	B	196	VAL	2.3
1	A	173	ALA	2.2
1	A	33	ASN	2.2
1	B	265	LEU	2.2
1	A	3	LYS	2.2
1	B	167[A]	LYS	2.2
1	A	176	ASP	2.2
1	A	110	TYR	2.2
1	B	134	ILE	2.2
1	A	225	GLU	2.2
1	A	65	ALA	2.2
1	B	305	ASP	2.2
1	A	288	LEU	2.2
1	A	171	GLU	2.2
1	A	95	ILE	2.2
1	A	134	ILE	2.2
1	B	95	ILE	2.2
1	A	227	VAL	2.2
1	B	211	SER	2.2
1	A	243	ALA	2.2
1	B	65	ALA	2.2
1	A	10[A]	GLN	2.2
1	A	60	LEU	2.2
1	A	63	THR	2.2
1	A	192	SER	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	187	ALA	2.1
1	B	5	LEU	2.1
1	B	28	ASN	2.1
1	B	279[A]	ASN	2.1
1	A	211	SER	2.1
1	A	214	VAL	2.1
1	A	300	VAL	2.1
1	A	306	ALA	2.1
1	B	20	ALA	2.1
1	B	71	VAL	2.1
1	A	152	LEU	2.1
1	B	36	PRO	2.1
1	A	101	ALA	2.1
1	A	64	ALA	2.0
1	A	183	ASN	2.0
1	B	61	VAL	2.0
1	A	283	GLU	2.0
1	A	278	PHE	2.0
1	B	138	ILE	2.0
1	A	61	VAL	2.0
1	A	273	LYS	2.0
1	B	133[A]	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
6	PG4	B	1002	13/13	0.74	0.20	48,56,81,82	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	4002	5/5	0.75	0.14	84,85,87,88	0
9	GOL	B	5003	6/6	0.75	0.17	53,62,68,69	0
9	GOL	B	5002	6/6	0.76	0.20	72,77,77,79	0
6	PG4	A	6002	13/13	0.78	0.18	64,72,82,82	0
8	1PE	B	1001	16/16	0.84	0.16	44,54,66,69	0
9	GOL	B	5001	6/6	0.85	0.15	42,54,60,66	0
5	CL	A	6001	1/1	0.86	0.14	86,86,86,86	0
4	SO4	B	4001	5/5	0.90	0.12	51,55,60,63	0
4	SO4	B	4004[B]	5/5	0.92	0.17	31,31,32,32	5
2	MG	A	1004	1/1	0.92	0.40	42,42,42,42	1
4	SO4	A	4003[B]	5/5	0.93	0.16	28,29,30,30	5
2	MG	B	1008	1/1	0.96	0.19	33,33,33,33	1
2	MG	A	1001	1/1	0.96	0.12	32,32,32,32	0
7	2PN	A	2001[A]	9/9	0.97	0.10	29,32,34,34	9
7	2PN	B	2002[A]	9/9	0.97	0.10	23,28,29,31	9
2	MG	B	1007	1/1	0.97	0.18	32,32,32,32	1
2	MG	B	1005	1/1	0.98	0.12	27,27,27,27	0
3	F	A	3001	1/1	0.98	0.17	35,35,35,35	0
2	MG	A	1002	1/1	0.98	0.13	33,33,33,33	0
2	MG	A	1003	1/1	0.99	0.12	35,35,35,35	1
3	F	B	3002	1/1	0.99	0.16	27,27,27,27	0
2	MG	B	1006	1/1	0.99	0.12	27,27,27,27	0

6.5 Other polymers ⓘ

There are no such residues in this entry.