



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:14 AM UTC

PDB ID : 4HA8 / pdb_00004ha8
Title : Structure of Staphylococcus aureus biotin protein ligase in complex with biotin acetylene
Authors : Yap, M.; Wilce, M.; Polyak, S; Soares da Costa, T.; Tieu, W.
Deposited on : 2012-09-25
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	 72% 24% ..

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2774 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Biotin-[acetyl-CoA-carboxylase] ligase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	322	Total	C	N	O	S	0	0	0
			2613	1658	450	499	6			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-4	HIS	-	expression tag	UNP E5R5T0
A	-3	HIS	-	expression tag	UNP E5R5T0
A	-2	HIS	-	expression tag	UNP E5R5T0
A	-1	HIS	-	expression tag	UNP E5R5T0
A	0	HIS	-	expression tag	UNP E5R5T0
A	1	HIS	-	expression tag	UNP E5R5T0
A	272	ASN	ILE	conflict	UNP E5R5T0

- Molecule 2 is (3aS,4S,6aR)-4-(hex-5-yn-1-yl)tetrahydro-1H-thieno[3,4-d]imidazol-2(3H)-one (CCD ID: BC4) (formula: C₁₁H₁₆N₂OS).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			15	11	2	1	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	146	Total	O	0	0
			146	146		

i

- Molecule 1: Biotin-[acetyl-CoA-carboxylase] ligase



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 ₂ 2 ₁ 2	Depositor
Cell constants a, b, c, α , β , γ	93.02Å 93.02Å 129.94Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.81 – 2.65 19.81 – 2.65	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.81-2.65) 99.4 (19.81-2.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.59 (at 2.67Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.197 , 0.236 0.190 , 0.228	Depositor DCC
R_{free} test set	856 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	52.0	Xtriage
Anisotropy	0.029	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	2774	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BC4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.19	1/2663 (0.0%)	1.15	6/3591 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	97	ALA	CA-CB	-5.21	1.45	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	GLN	N-CA-C	-6.17	101.64	110.59
1	A	83	ASP	N-CA-C	5.95	117.76	111.28
1	A	320	ASP	N-CA-CB	5.62	119.80	110.47
1	A	44	LYS	N-CA-C	-5.45	105.42	111.36
1	A	188	VAL	N-CA-C	-5.14	107.41	113.42
1	A	89	ASP	N-CA-C	-5.03	105.45	112.45

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2613	0	2574	57	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	15	0	16	2	0
3	A	146	0	0	5	0
All	All	2774	0	2590	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:6:GLN:HE21	1:A:70:GLY:H	1.23	0.87
1:A:6:GLN:HE22	1:A:68:TYR:HB3	1.54	0.72
1:A:20:ILE:HD13	1:A:25:ILE:HD11	1.76	0.68
1:A:12:LEU:CD1	1:A:20:ILE:HD11	2.25	0.66
1:A:6:GLN:NE2	1:A:70:GLY:H	1.95	0.64
1:A:308:ASP:OD1	3:A:538:HOH:O	2.16	0.62
1:A:6:GLN:HE21	1:A:70:GLY:N	1.96	0.62
1:A:81:LEU:HD22	1:A:253:GLU:HG3	1.82	0.62
1:A:105:ASN:HD22	1:A:107:SER:H	1.47	0.60
1:A:12:LEU:HD12	1:A:20:ILE:HD11	1.88	0.55
1:A:176:LYS:HD3	1:A:182:TYR:CE1	2.42	0.55
1:A:313:SER:OG	3:A:614:HOH:O	2.19	0.53
1:A:174:LYS:CB	1:A:276:ASN:HB3	2.40	0.51
1:A:20:ILE:CD1	1:A:25:ILE:HD11	2.40	0.51
1:A:317:ILE:HG22	3:A:529:HOH:O	2.10	0.51
1:A:69:GLN:HE22	1:A:87:VAL:H	1.58	0.51
1:A:220:PHE:O	1:A:221:ASP:C	2.54	0.50
1:A:220:PHE:O	1:A:221:ASP:O	2.31	0.48
1:A:75:TYR:N	1:A:75:TYR:HD1	2.12	0.47
1:A:239:LEU:HG	1:A:240:ASP:N	2.26	0.47
1:A:63:LEU:CD1	1:A:71:ILE:HD13	2.44	0.47
1:A:212:ASN:HB3	1:A:215:GLN:OE1	2.14	0.47
1:A:10:GLN:NE2	1:A:74:GLN:OE1	2.34	0.47
1:A:134:GLY:HA3	1:A:136:TRP:CH2	2.50	0.47
1:A:173:VAL:HG22	1:A:183:ILE:HG12	1.98	0.46
1:A:75:TYR:N	1:A:75:TYR:CD1	2.81	0.46
1:A:180:ASP:OD2	1:A:187:LYS:HE2	2.15	0.46
1:A:190:GLY:HA3	2:A:401:BC4:H2	1.98	0.46
1:A:157:ILE:HD12	1:A:157:ILE:HA	1.70	0.45
1:A:214:THR:HA	1:A:232:GLN:HB2	1.98	0.45
1:A:19:TYR:CD1	1:A:56:LYS:HE2	2.51	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:177:TRP:HA	1:A:178:PRO:HA	1.68	0.44
1:A:165:ILE:HG12	1:A:231:ILE:HD11	2.00	0.44
1:A:105:ASN:HD22	1:A:107:SER:N	2.14	0.44
1:A:220:PHE:C	1:A:221:ASP:O	2.60	0.44
1:A:90:SER:CB	1:A:115:GLU:HG2	2.48	0.43
1:A:189:CYS:HA	1:A:210:GLY:O	2.19	0.43
2:A:401:BC4:H15	2:A:401:BC4:H9	1.67	0.43
1:A:95:GLN:NE2	1:A:138:SER:OG	2.50	0.43
1:A:249:LEU:HD23	1:A:249:LEU:HA	1.77	0.43
1:A:181:ILE:HB	1:A:189:CYS:HB3	2.00	0.43
1:A:166:GLN:C	1:A:168:PHE:N	2.76	0.43
1:A:244:PHE:O	1:A:245:LEU:C	2.61	0.43
1:A:12:LEU:HD13	1:A:20:ILE:HD11	2.01	0.43
1:A:76:THR:HG21	1:A:85:SER:HB3	2.01	0.42
1:A:134:GLY:HA3	1:A:136:TRP:CZ2	2.54	0.42
1:A:61:GLN:HG2	3:A:646:HOH:O	2.18	0.42
1:A:276:ASN:HD22	1:A:276:ASN:H	1.67	0.42
1:A:174:LYS:HB3	1:A:276:ASN:HB3	2.01	0.41
1:A:176:LYS:HB2	1:A:276:ASN:HD21	1.86	0.41
1:A:147:PHE:C	1:A:147:PHE:CD1	2.98	0.41
1:A:9:LEU:HA	1:A:9:LEU:HD23	1.85	0.41
1:A:68:TYR:O	1:A:69:GLN:C	2.64	0.40
1:A:90:SER:HB2	1:A:115:GLU:HG2	2.03	0.40
1:A:319:ALA:HB1	3:A:543:HOH:O	2.22	0.40
1:A:88:TYR:O	1:A:113:SER:HA	2.22	0.40
1:A:166:GLN:C	1:A:168:PHE:H	2.29	0.40
1:A:159:LEU:HD21	1:A:271:TYR:HA	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	320/328 (98%)	304 (95%)	15 (5%)	1 (0%)	36	52

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	221	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/294 (98%)	270 (94%)	18 (6%)	16	29

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	12	LEU
1	A	14	LYS
1	A	20	ILE
1	A	33	ARG
1	A	34	THR
1	A	45	LEU
1	A	49	LYS
1	A	115	GLU
1	A	151	SER
1	A	169	SER
1	A	218	GLU
1	A	221	ASP
1	A	222	GLU
1	A	239	LEU
1	A	272	ASN
1	A	276	ASN
1	A	288	ASP
1	A	313	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	GLN
1	A	42	GLN
1	A	69	GLN
1	A	95	GLN
1	A	105	ASN
1	A	124	ASN
1	A	144	ASN
1	A	167	HIS
1	A	212	ASN
1	A	276	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	BC4	A	401	-	16,16,16	0.91	0	21,21,21	2.28	8 (38%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BC4	A	401	-	-	4/5/27/27	0/2/2/2

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	BC4	CAH-CAN-NAI	-5.20	106.48	113.18
2	A	401	BC4	CAO-CAM-SAK	-4.29	100.17	105.03
2	A	401	BC4	NAI-CAL-NAJ	3.37	112.74	108.85
2	A	401	BC4	CAM-CAO-NAJ	-2.90	110.27	113.34
2	A	401	BC4	CAN-CAH-SAK	-2.85	102.13	106.06
2	A	401	BC4	CAO-NAJ-CAL	-2.76	109.13	112.56
2	A	401	BC4	OAB-CAL-NAI	-2.40	122.44	125.89
2	A	401	BC4	CAH-SAK-CAM	2.21	94.57	89.98

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	BC4	CAF-CAG-CAM-SAK
2	A	401	BC4	CAF-CAG-CAM-CAO
2	A	401	BC4	CAC-CAD-CAE-CAF
2	A	401	BC4	CAD-CAE-CAF-CAG

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	401	BC4	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	322/328 (98%)	-0.53	0 100 100	25, 44, 62, 89	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	BC4	A	401	15/15	0.97	0.08	32,34,50,54	0

6.5 Other polymers [i](#)

There are no such residues in this entry.