



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 01:42 AM UTC

PDB ID : 6HAE / pdb_00006hae
Title : Crystal structure of [Fe]-hydrogenase (Hmd) from Methanococcus aeolicus in complex with FeGP cofactor and methenyl-tetrahydromethanopterin (close form B)
Authors : Huang, G.; Wagner, T.; Wodrich, M.D.; Ataka, K.; Bill, E.; Ermler, U.; Hu, X.; Shima, S.
Deposited on : 2018-08-07
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	342	96% 5%
1	K	342	95% 5%

2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 6029 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 5,10-methenyltetrahydromethanopterin hydrogenase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	342	Total	C	N	O	S	0	3	0
			2581	1635	426	496	24			
1	K	342	Total	C	N	O	S	0	5	0
			2592	1644	427	497	24			

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	K	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 1-{4-[(6S,6aR,7R)-3-amino-6,7-dimethyl-1-oxo-1,2,5,6,6a,7-hexahydro-8H-imidazo[1,5-f]pteridin-10-ium-8-yl]phenyl}-1-deoxy-5-O-{5-O-[(S)-{[(1S)-1,3-dicarboxypropyl]oxy}(hydroxy)phosphoryl]-alpha-D-ribofuranosyl}-D-ribitol (CCD ID: E4M) (formula: C₃₁H₄₄N₆O₁₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	S	0	0
			3	1	1	1		
5	A	1	Total	C	N	S	0	0
			3	1	1	1		
5	A	1	Total	C	N	S	0	0
			3	1	1	1		
5	A	1	Total	C	N	S	0	0
			3	1	1	1		
5	A	1	Total	C	N	S	0	0
			3	1	1	1		
5	A	1	Total	C	N	S	0	0
			3	1	1	1		
5	K	1	Total	C	N	S	0	0
			3	1	1	1		
5	K	1	Total	C	N	S	0	0
			3	1	1	1		
5	K	1	Total	C	N	S	0	0
			3	1	1	1		
5	K	1	Total	C	N	S	0	0
			3	1	1	1		
5	K	1	Total	C	N	S	0	0
			3	1	1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	K	1	Total	C	N	S	0	0
			3	1	1	1		


- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		
6	K	1	Total	Cl	0	0
			1	1		

- Molecule 7 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	K	0	0
			2	2		
7	K	2	Total	K	0	0
			2	2		

- Molecule 8 is iron-guanylyl pyridinol cofactor (CCD ID: FE9) (formula: C₂₁H₂₃FeN₆O₁₃PS).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
8	A	1	Total	C	Fe	N	O	P	0	0
			41	21	1	6	12	1		
8	K	1	Total	C	Fe	N	O	P	0	0
			41	21	1	6	12	1		

- Molecule 9 is NONAETHYLENE GLYCOL (CCD ID: 2PE) (formula: C₁₈H₃₈O₁₀).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
9	K	1	Total	C	O	0	0
			9	6	3		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	288	Total	O	0	0
			288	288		
10	K	304	Total	O	0	0
			304	304		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 5,10-methenyltetrahydromethanopterin hydrogenase

Chain A:  96%



- Molecule 1: 5,10-methenyltetrahydromethanopterin hydrogenase

Chain K:  95% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	80.02Å 156.48Å 53.65Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.56 – 1.85 44.56 – 1.85	Depositor EDS
% Data completeness (in resolution range)	97.1 (44.56-1.85) 97.1 (44.56-1.85)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.58 (at 1.86Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.161 , 0.192 0.159 , 0.191	Depositor DCC
R_{free} test set	2804 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	24.7	Xtrriage
Anisotropy	0.250	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6029	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 63.09 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0200e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, NA, GOL, E4M, SCN, 2PE, K, FE9

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	1/2631 (0.0%)	0.95	3/3562 (0.1%)
1	K	0.83	0/2645	0.96	3/3582 (0.1%)
All	All	0.83	1/5276 (0.0%)	0.96	6/7144 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	46	MET	SD-CE	-5.28	1.66	1.79

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	40	THR	N-CA-C	-6.50	102.15	110.53
1	K	280	ASP	CA-CB-CG	6.50	119.10	112.60
1	K	256	VAL	N-CA-C	-6.26	106.43	111.81
1	K	40	THR	N-CA-C	-5.69	103.19	110.53
1	A	280	ASP	CA-CB-CG	5.65	118.25	112.60
1	A	256	VAL	N-CA-C	-5.14	106.79	111.67

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2581	0	2613	7	0
1	K	2592	0	2631	9	0
2	A	6	0	8	0	0
2	K	6	0	8	0	0
3	A	54	0	41	0	0
3	K	54	0	41	4	0
4	A	1	0	0	0	0
4	K	1	0	0	0	0
5	A	24	0	0	0	0
5	K	21	0	0	0	0
6	A	1	0	0	0	0
6	K	1	0	0	0	0
7	A	2	0	0	0	0
7	K	2	0	0	0	0
8	A	41	0	20	1	0
8	K	41	0	20	1	0
9	K	9	0	10	0	0
10	A	288	0	0	1	0
10	K	304	0	0	0	0
All	All	6029	0	5392	18	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:K:402:E4M:H38	3:K:402:E4M:C5J	1.91	1.00
3:K:402:E4M:H38	3:K:402:E4M:H5J	1.38	1.00
1:A:260:VAL:HG22	1:K:260:VAL:HG22	1.70	0.71
10:A:720:HOH:O	1:K:340:HIS:HE1	1.87	0.56
3:K:402:E4M:C5J	3:K:402:E4M:C27	2.76	0.55
1:K:139:ILE:HD11	1:K:165:ALA:HB3	1.90	0.54
1:A:5:ILE:HG21	1:A:43:THR:HB	1.90	0.52
3:K:402:E4M:H38	3:K:402:E4M:H5JA	1.90	0.49
1:A:6:LEU:HD11	1:A:139:ILE:HD11	1.97	0.46
1:K:43:THR:HG23	1:K:148:TRP:CD2	2.50	0.46
1:A:175:ALA:HB1	8:A:415:FE9:O18	2.16	0.46
1:K:213:TYR:HD2	1:K:241:MET:HE3	1.82	0.45
1:K:176:CYS:SG	8:K:415:FE9:C7	3.03	0.45
1:K:5:ILE:HG21	1:K:43:THR:HB	2.00	0.44
1:A:213:TYR:HD2	1:A:241:MET:HE3	1.82	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:163:VAL:HG22	1:K:192:ARG:HG3	2.03	0.41
1:A:43:THR:HG23	1:A:148:TRP:CD2	2.56	0.41
1:A:252:MET:HE3	1:K:285:MET:HE1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	343/342 (100%)	332 (97%)	11 (3%)	0	100	100
1	K	345/342 (101%)	332 (96%)	13 (4%)	0	100	100
All	All	688/684 (101%)	664 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	280/277 (101%)	280 (100%)	0	100	100
1	K	282/277 (102%)	282 (100%)	0	100	100
All	All	562/554 (101%)	562 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 30 ligands modelled in this entry, 8 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SCN	K	404	-	1,2,2	0.22	0	0,1,1	-	-
5	SCN	A	406	-	1,2,2	0.51	0	0,1,1	-	-
2	GOL	A	401	-	5,5,5	0.13	0	5,5,5	0.13	0
3	E4M	A	402	-	54,58,58	1.22	3 (5%)	64,86,86	1.37	7 (10%)
5	SCN	K	408	-	1,2,2	0.48	0	0,1,1	-	-
5	SCN	K	410	-	1,2,2	0.08	0	0,1,1	-	-
5	SCN	A	410	-	1,2,2	0.02	0	0,1,1	-	-
3	E4M	K	402	-	54,58,58	1.21	4 (7%)	64,86,86	1.34	7 (10%)
5	SCN	K	405	-	1,2,2	0.76	0	0,1,1	-	-
5	SCN	A	404	-	1,2,2	0.07	0	0,1,1	-	-
5	SCN	K	409	-	1,2,2	0.39	0	0,1,1	-	-
9	2PE	K	414	-	8,8,27	0.47	0	7,7,26	0.37	0
8	FE9	K	415	1	37,45,47	1.46	4 (10%)	49,71,80	1.93	15 (30%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SCN	A	408	-	1,2,2	0.36	0	0,1,1	-	-
8	FE9	A	415	1	37,45,47	1.48	4 (10%)	49,71,80	1.88	13 (26%)
2	GOL	K	401	-	5,5,5	0.26	0	5,5,5	0.32	0
5	SCN	A	409	-	1,2,2	0.25	0	0,1,1	-	-
5	SCN	K	406	-	1,2,2	0.13	0	0,1,1	-	-
5	SCN	A	407	-	1,2,2	0.50	0	0,1,1	-	-
5	SCN	A	411	-	1,2,2	0.55	0	0,1,1	-	-
5	SCN	K	407	-	1,2,2	0.26	0	0,1,1	-	-
5	SCN	A	405	-	1,2,2	0.37	0	0,1,1	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	E4M	K	402	-	-	7/41/85/85	0/5/5/5
2	GOL	A	401	-	-	2/4/4/4	-
8	FE9	A	415	1	-	2/13/53/65	0/5/5/5
3	E4M	A	402	-	-	6/41/85/85	0/5/5/5
2	GOL	K	401	-	-	2/4/4/4	-
9	2PE	K	414	-	-	4/6/6/25	-
8	FE9	K	415	1	-	3/13/53/65	0/5/5/5

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	402	E4M	C10-N5	-5.51	1.27	1.33
8	A	415	FE9	C4-C3	4.73	1.47	1.36
8	K	415	FE9	C4-C3	4.70	1.47	1.36
3	K	402	E4M	C10-N5	-4.40	1.28	1.33
8	K	415	FE9	C6-C5	3.34	1.47	1.36
3	K	402	E4M	C10-N10	-3.06	1.28	1.35
8	A	415	FE9	C5A-C4A	3.01	1.47	1.38
3	A	402	E4M	C10-N10	-2.95	1.28	1.35
8	K	415	FE9	C5A-C4A	2.86	1.46	1.38
8	A	415	FE9	C6-C5	2.83	1.46	1.36
3	A	402	E4M	OX5-C1J	2.59	1.44	1.40
8	A	415	FE9	C2-N1	2.59	1.40	1.37
3	K	402	E4M	OX5-C1J	2.41	1.44	1.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	K	402	E4M	C4-N3	-2.25	1.34	1.38
8	K	415	FE9	C4A-N9A	-2.21	1.32	1.38

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	415	FE9	C5A-C4A-N3A	-5.44	119.73	128.39
8	A	415	FE9	C2A-N3A-C4A	4.96	120.84	112.30
3	A	402	E4M	C2-N1-C8A	4.86	121.94	113.36
8	K	415	FE9	C5A-C4A-N3A	-4.73	120.86	128.39
8	K	415	FE9	C2A-N3A-C4A	4.73	120.44	112.30
8	A	415	FE9	N9A-C4A-N3A	4.50	134.96	125.95
3	K	402	E4M	O4J-C1J-OX5	-4.45	107.07	111.95
3	A	402	E4M	OH4-C4-C4A	-4.15	118.75	128.01
3	K	402	E4M	C2-N1-C8A	4.13	120.64	113.36
8	K	415	FE9	N9A-C4A-N3A	3.84	133.63	125.95
8	K	415	FE9	C6A-C5A-N7A	3.75	137.11	130.29
3	A	402	E4M	C4A-C4-N3	3.48	119.03	111.84
3	A	402	E4M	O4J-C1J-OX5	-3.32	108.31	111.95
8	A	415	FE9	C6A-C5A-N7A	3.31	136.31	130.29
3	K	402	E4M	OH4-C4-C4A	-3.13	121.02	128.01
8	K	415	FE9	O6A-C6A-C5A	-3.06	118.44	126.53
3	K	402	E4M	C7M-C7-N8	-2.98	106.59	109.44
3	K	402	E4M	C4A-C4-N3	2.93	117.90	111.84
8	K	415	FE9	C5-C4-C3	-2.93	119.19	123.89
8	A	415	FE9	O2P-P1-O3P	2.74	113.82	104.94
8	A	415	FE9	O6A-C6A-C5A	-2.68	119.45	126.53
3	A	402	E4M	C9M-C9-N10	2.65	114.35	111.46
3	A	402	E4M	O16-C31-C27	2.64	119.58	112.71
3	K	402	E4M	O16-C31-C27	2.39	118.92	112.71
8	K	415	FE9	O2-C2-N1	-2.34	119.12	124.11
8	K	415	FE9	C8A-N9A-C4A	2.27	110.27	106.03
8	K	415	FE9	C5A-C6A-N1A	2.25	118.98	113.25
8	A	415	FE9	C8A-N9A-C4A	2.21	110.17	106.03
3	K	402	E4M	O15-C31-C27	-2.21	118.07	122.85
8	K	415	FE9	O2P-P1-O3P	2.21	112.08	104.94
3	A	402	E4M	O15-C31-C27	-2.21	118.09	122.85
8	K	415	FE9	C4A-C5A-N7A	-2.18	107.22	110.67
8	A	415	FE9	O18-C8-C7	2.12	122.31	119.17
8	K	415	FE9	C6A-C5A-C4A	-2.12	115.63	118.83
8	A	415	FE9	C4'-O4'-C1'	-2.12	104.78	109.47
8	A	415	FE9	C5A-C6A-N1A	2.10	118.59	113.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	A	415	FE9	C4A-C5A-N7A	-2.09	107.35	110.67
8	K	415	FE9	N9A-C8A-N7A	-2.06	109.59	113.40
8	K	415	FE9	C4'-O4'-C1'	-2.04	104.95	109.47
8	K	415	FE9	C3M-C3-C2	2.04	121.75	116.18
8	A	415	FE9	N9A-C8A-N7A	-2.02	109.66	113.40
8	A	415	FE9	C5-C6-N1	-2.00	121.47	126.01

There are no chirality outliers.

All (26) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	K	401	GOL	O2-C2-C3-O3
3	A	402	E4M	O2A-C27-C31-O16
3	K	402	E4M	C27-O2A-PA-O5J
3	K	402	E4M	C5J-O5J-PA-O3A
3	K	402	E4M	O2A-C27-C28-C29
3	K	402	E4M	C31-C27-C28-C29
3	K	402	E4M	O2A-C27-C31-O16
9	K	414	2PE	O19-C20-C21-O22
2	A	401	GOL	C1-C2-C3-O3
2	K	401	GOL	C1-C2-C3-O3
3	A	402	E4M	C3J-C4J-C5J-O5J
9	K	414	2PE	C24-C23-O22-C21
9	K	414	2PE	O16-C17-C18-O19
3	K	402	E4M	O2A-C27-C31-O15
8	K	415	FE9	C4-O3P-P1-O1P
8	K	415	FE9	O4'-C4'-C5'-O5'
2	A	401	GOL	O2-C2-C3-O3
3	A	402	E4M	O4J-C4J-C5J-O5J
8	A	415	FE9	C4-O3P-P1-O1P
8	A	415	FE9	O4'-C4'-C5'-O5'
3	A	402	E4M	C28-C29-C30-O14
3	A	402	E4M	C28-C29-C30-O13
9	K	414	2PE	C21-C20-O19-C18
3	K	402	E4M	C27-O2A-PA-O1A
3	A	402	E4M	O2A-C27-C31-O15
8	K	415	FE9	C3'-C4'-C5'-O5'




There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	K	402	E4M	4	0
8	K	415	FE9	1	0
8	A	415	FE9	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	342/342 (100%)	-0.42	1 (0%) 90 92	14, 24, 46, 67	3 (0%)
1	K	342/342 (100%)	-0.43	1 (0%) 90 92	11, 24, 46, 68	5 (1%)
All	All	684/684 (100%)	-0.42	2 (0%) 90 92	11, 24, 46, 68	8 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	342	LYS	2.7
1	A	342	LYS	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SCN	K	408	3/3	0.73	0.20	70,70,74,78	0
5	SCN	A	405	3/3	0.74	0.18	55,55,59,67	0
2	GOL	K	401	6/6	0.77	0.17	44,54,60,61	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	2PE	K	414	9/28	0.78	0.15	39,57,65,66	0
5	SCN	K	409	3/3	0.80	0.16	72,72,73,78	0
5	SCN	A	411	3/3	0.81	0.23	40,40,41,49	3
5	SCN	K	404	3/3	0.82	0.14	67,67,69,75	0
5	SCN	K	405	3/3	0.83	0.25	56,56,57,62	0
5	SCN	A	406	3/3	0.84	0.16	55,55,61,70	0
5	SCN	A	407	3/3	0.85	0.15	78,78,80,83	0
5	SCN	A	409	3/3	0.86	0.15	77,77,77,78	0
5	SCN	K	406	3/3	0.88	0.16	59,59,60,62	0
5	SCN	A	410	3/3	0.89	0.13	64,64,66,66	0
2	GOL	A	401	6/6	0.89	0.11	29,52,58,63	0
5	SCN	K	407	3/3	0.90	0.17	46,46,47,48	0
5	SCN	A	404	3/3	0.92	0.15	45,45,46,48	0
7	K	K	413	1/1	0.93	0.22	58,58,58,58	0
5	SCN	A	408	3/3	0.93	0.13	46,46,54,59	0
5	SCN	K	410	3/3	0.94	0.10	44,44,51,51	0
4	NA	K	403	1/1	0.95	0.09	28,28,28,28	0
3	E4M	K	402	54/54	0.95	0.09	16,25,100,104	0
3	E4M	A	402	54/54	0.96	0.09	16,27,114,115	0
7	K	K	412	1/1	0.97	0.14	41,41,41,41	0
6	CL	A	412	1/1	0.97	0.07	35,35,35,35	0
8	FE9	A	415	41/43	0.97	0.06	20,24,31,38	0
8	FE9	K	415	41/43	0.97	0.06	17,23,30,33	0
7	K	A	413	1/1	0.97	0.18	44,44,44,44	0
7	K	A	414	1/1	0.98	0.23	54,54,54,54	0
6	CL	K	411	1/1	0.98	0.04	32,32,32,32	0
4	NA	A	403	1/1	0.98	0.08	27,27,27,27	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 2PE K 414:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around E4M K 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around E4M A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around FE9 A 415:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FE9 K 415:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.