



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 12, 2026 – 04:52 PM UTC

PDB ID : 6HBD / pdb_00006hbd
Title : Crystal structure of MSMEG_1712 from Mycobacterium smegmatis in complex with Beta-D-Galactofuranose
Authors : Li, M.; Mueller, C.; Einsle, O.; Jessen-Trefzer, C.
Deposited on : 2018-08-10
Resolution : 2.44 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.44 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2340 (2.46-2.42)
Clashscore	190562	2400 (2.46-2.42)
Ramachandran outliers	187476	2379 (2.46-2.42)
Sidechain outliers	187428	2379 (2.46-2.42)
RSRZ outliers	180081	2340 (2.46-2.42)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	320	8% 83% 11% • 5%
1	B	320	30% 80% 13% • 5%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 9099 atoms, of which 4457 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ABC transporter periplasmic-binding protein YtfQ.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	305	Total	C	H	N	O	S	12	0	0
			4519	1429	2232	399	453	6			
1	B	305	Total	C	H	N	O	S	30	0	0
			4485	1428	2201	399	451	6			

There are 28 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP A0QT50
A	308	LYS	-	expression tag	UNP A0QT50
A	309	LEU	-	expression tag	UNP A0QT50
A	310	ALA	-	expression tag	UNP A0QT50
A	311	ALA	-	expression tag	UNP A0QT50
A	312	ALA	-	expression tag	UNP A0QT50
A	313	LEU	-	expression tag	UNP A0QT50
A	314	GLU	-	expression tag	UNP A0QT50
A	315	HIS	-	expression tag	UNP A0QT50
A	316	HIS	-	expression tag	UNP A0QT50
A	317	HIS	-	expression tag	UNP A0QT50
A	318	HIS	-	expression tag	UNP A0QT50
A	319	HIS	-	expression tag	UNP A0QT50
A	320	HIS	-	expression tag	UNP A0QT50
B	1	MET	-	initiating methionine	UNP A0QT50
B	308	LYS	-	expression tag	UNP A0QT50
B	309	LEU	-	expression tag	UNP A0QT50
B	310	ALA	-	expression tag	UNP A0QT50
B	311	ALA	-	expression tag	UNP A0QT50
B	312	ALA	-	expression tag	UNP A0QT50
B	313	LEU	-	expression tag	UNP A0QT50
B	314	GLU	-	expression tag	UNP A0QT50
B	315	HIS	-	expression tag	UNP A0QT50
B	316	HIS	-	expression tag	UNP A0QT50
B	317	HIS	-	expression tag	UNP A0QT50

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	318	HIS	-	expression tag	UNP A0QT50
B	319	HIS	-	expression tag	UNP A0QT50
B	320	HIS	-	expression tag	UNP A0QT50

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	9	Total	Zn	0	0
			9	9		
2	B	6	Total	Zn	0	0
			6	6		

- Molecule 3 is beta-D-galactofuranose (CCD ID: GZL) (formula: C₆H₁₂O₆) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			24	6	12	6		
3	B	1	Total	C	H	O	0	0
			24	6	12	6		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	20	Total	O	0	0
			20	20		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	12	Total	O	0	0
			12	12		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ABC transporter periplasmic-binding protein YtfQ



- Molecule 1: ABC transporter periplasmic-binding protein YtfQ



4 Data and refinement statistics

Property	Value	Source
Space group	I 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	118.30Å 118.30Å 231.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	105.31 – 2.44 105.31 – 2.44	Depositor EDS
% Data completeness (in resolution range)	58.3 (105.31-2.44) 58.3 (105.31-2.44)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 2.45Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.227 , 0.266 0.241 , 0.279	Depositor DCC
R_{free} test set	872 reflections (4.87%)	wwPDB-VP
Wilson B-factor (Å ²)	70.8	Xtriage
Anisotropy	0.069	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 91.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9099	wwPDB-VP
Average B, all atoms (Å ²)	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, GZL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.91	1/2328 (0.0%)	1.24	8/3157 (0.3%)
1	B	0.88	0/2325	1.26	7/3153 (0.2%)
All	All	0.90	1/4653 (0.0%)	1.25	15/6310 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	247	MET	SD-CE	-9.63	1.55	1.79

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	214	ASP	CA-CB-CG	9.09	121.69	112.60
1	B	214	ASP	CA-CB-CG	8.97	121.57	112.60
1	B	241	ASP	CA-C-N	-7.40	111.84	123.24
1	B	241	ASP	C-N-CA	-7.40	111.84	123.24
1	B	56	ASN	CA-CB-CG	6.81	119.41	112.60
1	A	90	VAL	N-CA-CB	6.12	118.86	110.54
1	A	105	ASP	CA-CB-CG	5.56	118.16	112.60
1	A	187	PHE	CA-CB-CG	5.48	119.28	113.80
1	A	241	ASP	CA-C-O	-5.42	112.76	120.51
1	A	158	ASP	CA-CB-CG	5.33	117.93	112.60
1	A	294	ASP	N-CA-C	-5.26	101.71	109.86
1	B	241	ASP	CA-C-O	-5.14	113.16	120.51
1	B	294	ASP	N-CA-C	-5.05	102.14	109.31
1	B	294	ASP	CA-CB-CG	5.04	117.64	112.60
1	A	213	ASN	CA-CB-CG	5.01	117.61	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2287	2232	2237	7	0
1	B	2284	2201	2235	9	0
2	A	9	0	0	0	0
2	B	6	0	0	0	0
3	A	12	12	9	0	0
3	B	12	12	9	0	0
4	A	20	0	0	0	0
4	B	12	0	0	0	0
All	All	4642	4457	4490	16	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (16) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:70:ILE:HD11	1:B:94:THR:HG22	1.65	0.76
1:B:220:ALA:O	1:B:224:ILE:HG23	2.03	0.58
1:A:181:ALA:HB1	1:A:199:PHE:CE1	2.40	0.56
1:B:179:ILE:HG23	1:B:179:ILE:O	2.11	0.51
1:B:25:VAL:HG12	1:B:26:GLY:H	1.77	0.49
1:A:149:VAL:CG1	1:A:199:PHE:CD2	2.96	0.49
1:B:237:ILE:HG23	1:B:255:PHE:HA	1.95	0.48
1:A:36:THR:HG23	1:A:52:PHE:CZ	2.50	0.47
1:B:36:THR:HG23	1:B:52:PHE:CZ	2.50	0.47
1:B:74:VAL:HG23	1:B:99:ILE:HD11	1.98	0.45
1:A:90:VAL:O	1:A:94:THR:HG23	2.16	0.45
1:A:241:ASP:O	1:A:242:ALA:HB3	2.16	0.44
1:A:136:ALA:HA	1:A:177:LEU:HD11	1.99	0.44
1:B:136:ALA:HA	1:B:177:LEU:HD11	2.00	0.44
1:A:250:LEU:HD22	1:A:258:ILE:HG13	2.02	0.42
1:B:250:LEU:HD22	1:B:258:ILE:HG13	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	303/320 (95%)	281 (93%)	18 (6%)	4 (1%)	9	9
1	B	303/320 (95%)	283 (93%)	12 (4%)	8 (3%)	4	2
All	All	606/640 (95%)	564 (93%)	30 (5%)	12 (2%)	6	4

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	ASP
1	A	303	PRO
1	B	56	ASN
1	B	241	ASP
1	B	303	PRO
1	B	204	PRO
1	B	110	THR
1	A	204	PRO
1	A	81	PRO
1	B	25	VAL
1	B	81	PRO
1	B	266	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	233/242 (96%)	213 (91%)	20 (9%)	10	11

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	232/242 (96%)	207 (89%)	25 (11%)	6	5
All	All	465/484 (96%)	420 (90%)	45 (10%)	8	8

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	LEU
1	A	56	ASN
1	A	84	ARG
1	A	90	VAL
1	A	95	LYS
1	A	104	THR
1	A	115	VAL
1	A	120	ILE
1	A	126	GLU
1	A	171	ILE
1	A	183	GLN
1	A	208	VAL
1	A	209	VAL
1	A	213	ASN
1	A	229	LYS
1	A	238	VAL
1	A	247	MET
1	A	287	VAL
1	A	305	ARG
1	A	313	LEU
1	B	20	LEU
1	B	28	GLU
1	B	32	ARG
1	B	72	GLN
1	B	110	THR
1	B	116	TYR
1	B	120	ILE
1	B	143	THR
1	B	150	GLN
1	B	183	GLN
1	B	193	LYS
1	B	205	GLN
1	B	206	ILE
1	B	209	VAL
1	B	213	ASN
1	B	224	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	225	GLU
1	B	230	LYS
1	B	235	ILE
1	B	237	ILE
1	B	238	VAL
1	B	276	VAL
1	B	285	ARG
1	B	305	ARG
1	B	313	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	56	ASN
1	B	92	GLN
1	B	96	ASN
1	B	194	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 15 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GZL	A	410	-	12,12,12	0.32	0	16,17,17	0.83	0
3	GZL	B	407	-	12,12,12	0.23	0	16,17,17	0.86	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GZL	A	410	-	-	0/6/22/22	0/1/1/1
3	GZL	B	407	-	-	1/6/22/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	407	GZL	C1-C2-C3	-2.40	99.33	102.29

There are no chirality outliers.


All (1) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	407	GZL	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	305/320 (95%)	0.76	27 (8%) 15 12	20, 65, 89, 100	1 (0%)
1	B	305/320 (95%)	1.55	95 (31%) 1 1	26, 89, 116, 126	3 (0%)
All	All	610/640 (95%)	1.16	122 (20%) 3 2	20, 74, 111, 126	4 (0%)

All (122) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	112	ASP	5.5
1	A	116	TYR	4.7
1	B	148	ILE	4.4
1	B	111	GLN	4.3
1	B	302	LEU	4.2
1	B	102	ILE	4.1
1	B	63	ILE	4.0
1	B	226	ALA	3.9
1	B	113	THR	3.9
1	A	82	VAL	3.8
1	B	235	ILE	3.8
1	B	79	PHE	3.7
1	B	147	ASN	3.7
1	B	91	LEU	3.6
1	A	108	VAL	3.6
1	B	237	ILE	3.5
1	B	151	LEU	3.5
1	B	278	ALA	3.5
1	B	46	ALA	3.5
1	A	109	ASP	3.4
1	B	149	VAL	3.4
1	B	77	ILE	3.4
1	A	282	VAL	3.3
1	A	171	ILE	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	110	THR	3.3
1	B	107	ALA	3.2
1	B	144	GLY	3.2
1	B	200	LEU	3.2
1	B	146	VAL	3.2
1	B	153	GLY	3.2
1	A	90	VAL	3.1
1	B	20	LEU	3.1
1	B	185	GLY	3.1
1	A	63	ILE	3.1
1	A	144	GLY	3.1
1	B	246	GLY	3.1
1	A	120	ILE	3.1
1	B	209	VAL	3.1
1	B	139	TYR	3.0
1	B	103	LEU	3.0
1	A	119	PHE	3.0
1	B	187	PHE	3.0
1	B	210	PHE	3.0
1	B	120	ILE	2.9
1	A	107	ALA	2.9
1	B	269	LEU	2.9
1	B	179	ILE	2.9
1	B	36	THR	2.9
1	B	108	VAL	2.9
1	B	143	THR	2.9
1	B	115	VAL	2.9
1	B	198	ALA	2.8
1	B	142	ALA	2.8
1	B	180	VAL	2.8
1	B	73	GLY	2.7
1	B	16	GLY	2.7
1	A	142	ALA	2.7
1	B	129	ARG	2.7
1	B	206	ILE	2.7
1	B	208	VAL	2.6
1	B	173	LYS	2.6
1	B	220	ALA	2.6
1	B	231	PRO	2.6
1	B	116	TYR	2.6
1	B	82	VAL	2.6
1	B	66	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	135	VAL	2.5
1	B	150	GLN	2.5
1	A	174	ASN	2.5
1	B	211	ALA	2.5
1	B	87	TRP	2.5
1	B	223	ALA	2.5
1	B	118	THR	2.5
1	A	145	PRO	2.5
1	B	170	GLY	2.5
1	A	84	ARG	2.5
1	B	239	ALA	2.5
1	B	86	GLY	2.4
1	A	286	VAL	2.4
1	B	307	TYR	2.4
1	B	221	MET	2.4
1	B	57	GLY	2.4
1	A	170	GLY	2.4
1	B	218	LEU	2.4
1	B	17	ALA	2.4
1	B	224	ILE	2.3
1	B	234	ASP	2.3
1	B	175	PRO	2.3
1	B	214	ASP	2.3
1	A	115	VAL	2.3
1	B	101	VAL	2.3
1	B	277	ALA	2.3
1	A	70	ILE	2.3
1	B	240	VAL	2.3
1	B	157	ALA	2.2
1	B	232	GLY	2.2
1	A	16	GLY	2.2
1	A	86	GLY	2.2
1	B	227	ALA	2.2
1	B	292	ALA	2.2
1	B	85	THR	2.2
1	B	228	GLY	2.2
1	B	189	ARG	2.2
1	B	90	VAL	2.1
1	A	17	ALA	2.1
1	B	59	GLN	2.1
1	B	242	ALA	2.1
1	B	276	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	145	PRO	2.1
1	A	103	LEU	2.1
1	A	177	LEU	2.1
1	B	264	LEU	2.1
1	B	52	PHE	2.1
1	B	272	LEU	2.1
1	B	225	GLU	2.1
1	B	247	MET	2.1
1	A	79	PHE	2.0
1	B	259	VAL	2.0
1	A	102	ILE	2.0
1	B	203	THR	2.0
1	B	257	TYR	2.0
1	B	169	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	ZN	B	401	1/1	0.87	0.18	148,148,148,148	0
2	ZN	A	409	1/1	0.88	0.09	160,160,160,160	0
2	ZN	A	408	1/1	0.92	0.20	110,110,110,110	0
3	GZL	A	410	12/12	0.94	0.07	46,47,50,51	0
3	GZL	B	407	12/12	0.94	0.08	54,60,66,69	0
2	ZN	B	406	1/1	0.97	0.11	63,63,63,63	0
2	ZN	A	404	1/1	0.98	0.04	87,87,87,87	0
2	ZN	A	405	1/1	0.98	0.11	67,67,67,67	0
2	ZN	A	407	1/1	0.98	0.08	59,59,59,59	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	B	402	1/1	0.99	0.07	52,52,52,52	0
2	ZN	B	403	1/1	0.99	0.09	57,57,57,57	0
2	ZN	B	404	1/1	0.99	0.08	65,65,65,65	0
2	ZN	B	405	1/1	0.99	0.12	65,65,65,65	0
2	ZN	A	403	1/1	0.99	0.08	93,93,93,93	0
2	ZN	A	406	1/1	0.99	0.05	69,69,69,69	0
2	ZN	A	401	1/1	0.99	0.09	48,48,48,48	0
2	ZN	A	402	1/1	1.00	0.08	44,44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around GZL A 410:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around GZL B 407:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.