



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 09:33 AM UTC

PDB ID : 6HBV / pdb_00006hbv
Title : Structure of PLP internal aldimine form of Sphingopyxis sp. MTA144 FumI protein
Authors : Campopiano, D.J.; Serpico, A.; Marles-Wright, J.
Deposited on : 2018-08-13
Resolution : 1.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.48.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	2328 (1.66-1.66)
Clashscore	180529	2515 (1.66-1.66)
Ramachandran outliers	177936	2475 (1.66-1.66)
Sidechain outliers	177891	2475 (1.66-1.66)
RSRZ outliers	164620	2328 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	85% 8% 6%
1	B	440	85% 8% 7%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7424 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aminopentol aminotransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	414	Total	C	N	O	S	0	3	0
			3224	2038	572	594	20			
1	B	411	Total	C	N	O	S	0	3	0
			3193	2017	566	591	19			

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	423	ALA	-	expression tag	UNP D2D3B2
A	424	SER	-	expression tag	UNP D2D3B2
A	425	SER	-	expression tag	UNP D2D3B2
A	426	VAL	-	expression tag	UNP D2D3B2
A	427	ASP	-	expression tag	UNP D2D3B2
A	428	LYS	-	expression tag	UNP D2D3B2
A	429	LEU	-	expression tag	UNP D2D3B2
A	430	ALA	-	expression tag	UNP D2D3B2
A	431	ALA	-	expression tag	UNP D2D3B2
A	432	ALA	-	expression tag	UNP D2D3B2
A	433	LEU	-	expression tag	UNP D2D3B2
A	434	GLU	-	expression tag	UNP D2D3B2
A	435	HIS	-	expression tag	UNP D2D3B2
A	436	HIS	-	expression tag	UNP D2D3B2
A	437	HIS	-	expression tag	UNP D2D3B2
A	438	HIS	-	expression tag	UNP D2D3B2
A	439	HIS	-	expression tag	UNP D2D3B2
A	440	HIS	-	expression tag	UNP D2D3B2
B	423	ALA	-	expression tag	UNP D2D3B2
B	424	SER	-	expression tag	UNP D2D3B2
B	425	SER	-	expression tag	UNP D2D3B2
B	426	VAL	-	expression tag	UNP D2D3B2
B	427	ASP	-	expression tag	UNP D2D3B2
B	428	LYS	-	expression tag	UNP D2D3B2
B	429	LEU	-	expression tag	UNP D2D3B2

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	430	ALA	-	expression tag	UNP D2D3B2
B	431	ALA	-	expression tag	UNP D2D3B2
B	432	ALA	-	expression tag	UNP D2D3B2
B	433	LEU	-	expression tag	UNP D2D3B2
B	434	GLU	-	expression tag	UNP D2D3B2
B	435	HIS	-	expression tag	UNP D2D3B2
B	436	HIS	-	expression tag	UNP D2D3B2
B	437	HIS	-	expression tag	UNP D2D3B2
B	438	HIS	-	expression tag	UNP D2D3B2
B	439	HIS	-	expression tag	UNP D2D3B2
B	440	HIS	-	expression tag	UNP D2D3B2

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	1
			15	8	1	5	1		
2	B	1	Total	C	N	O	P	0	1
			15	8	1	5	1		

- Molecule 3 is 4'-DEOXY-4'-AMINOPYRIDOXAL-5'-PHOSPHATE (CCD ID: PMP) (formula: $C_8H_{13}N_2O_5P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	1
			16	8	2	5	1		
3	B	1	Total	C	N	O	P	0	1
			16	8	2	5	1		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Mg	0	0
			1	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	467	Total	O	0	0
			467	467		
5	B	477	Total	O	0	0
			477	477		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Aminopentol aminotransferase



• Molecule 1: Aminopentol aminotransferase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.25Å 103.90Å 65.33Å 90.00° 109.61° 90.00°	Depositor
Resolution (Å)	46.91 – 1.65 46.91 – 1.65	Depositor EDS
% Data completeness (in resolution range)	99.5 (46.91-1.65) 99.8 (46.91-1.65)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.01 (at 1.64Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.143 , 0.174 0.156 , 0.184	Depositor DCC
R_{free} test set	4764 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	16.9	Xtriage
Anisotropy	0.933	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 33.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7424	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PMP, PLP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	3/3313 (0.1%)	1.01	9/4500 (0.2%)
1	B	0.94	0/3284	1.01	5/4461 (0.1%)
All	All	0.93	3/6597 (0.0%)	1.01	14/8961 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	4
All	All	0	8

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	138	ILE	C-O	-5.40	1.18	1.24
1	A	76	ALA	N-CA	5.19	1.52	1.46
1	A	17	VAL	N-CA	5.18	1.51	1.46

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	12	GLU	CB-CG-CD	6.80	124.16	112.60
1	A	241	ASP	CA-CB-CG	6.74	119.34	112.60
1	A	79	ASP	CA-CB-CG	6.11	118.71	112.60
1	B	120	ASP	CA-CB-CG	5.70	118.30	112.60
1	B	241	ASP	CA-CB-CG	5.60	118.20	112.60
1	A	199	HIS	CA-CB-CG	5.52	119.32	113.80

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	PHE	CA-CB-CG	5.48	119.28	113.80
1	B	197	PHE	CA-CB-CG	5.40	119.20	113.80
1	B	390	THR	CA-CB-OG1	-5.34	101.59	109.60
1	B	114	PHE	CA-CB-CG	-5.29	108.51	113.80
1	A	197	PHE	CA-CB-CG	5.26	119.06	113.80
1	A	293	VAL	N-CA-CB	5.21	114.52	110.45
1	A	290	PHE	CA-CB-CG	-5.17	108.64	113.80
1	A	89	THR	CA-CB-OG1	-5.12	101.92	109.60

There are no chirality outliers.

All (8) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	214	ARG	Sidechain
1	A	356	ARG	Sidechain
1	A	398	ARG	Sidechain
1	A	6	ARG	Sidechain
1	B	107	ARG	Sidechain
1	B	280	ARG	Sidechain
1	B	398	ARG	Sidechain
1	B	405	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3224	0	3113	17	0
1	B	3193	0	3081	19	0
2	A	15	0	6	1	0
2	B	15	0	6	1	0
3	A	16	0	10	0	0
3	B	16	0	11	2	0
4	A	1	0	0	0	0
5	A	467	0	0	6	0
5	B	477	0	0	4	0
All	All	7424	0	6227	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:11:ARG:NH1	5:A:603:HOH:O	2.31	0.63
1:B:185:HIS:HE1	5:B:975:HOH:O	1.83	0.61
1:B:329:SER:HB2	1:B:334:LEU:O	2.01	0.61
1:A:82:ARG:NH1	5:A:602:HOH:O	2.28	0.60
1:B:233:ARG:HB2	2:B:501[A]:PLP:O3	2.05	0.57
1:A:233:ARG:HB2	2:A:501[A]:PLP:O3	2.05	0.56
1:A:246:HIS:HB2	5:A:732:HOH:O	2.05	0.55
1:B:40:ARG:NH1	1:B:50:ALA:O	2.41	0.54
1:A:413:HIS:HB3	1:A:416:LEU:HB2	1.91	0.53
3:B:502[B]:PMP:O3	3:B:502[B]:PMP:N4A	2.42	0.52
1:B:142:LYS:HE3	5:B:989:HOH:O	2.10	0.51
1:B:246:HIS:HB3	5:B:826:HOH:O	2.10	0.51
1:A:210:LEU:HD21	1:A:214:ARG:CZ	2.42	0.50
1:A:67:LEU:HD12	1:A:74:ILE:HD11	1.93	0.49
1:A:140:CYS:O	1:A:169:TYR:HA	2.14	0.48
1:A:89:THR:HA	1:B:61:ALA:HA	1.97	0.47
1:A:59:MET:SD	1:A:373:HIS:HB2	2.55	0.46
1:A:111:TRP:CZ2	1:A:271:GLY:HA3	2.51	0.46
1:A:398:ARG:CZ	5:A:634:HOH:O	2.63	0.46
1:B:229:VAL:HG23	1:B:251:PRO:HB3	1.98	0.46
1:A:61:ALA:HA	1:B:89:THR:HA	1.98	0.45
1:A:107:ARG:HG3	1:A:241:ASP:OD2	2.16	0.45
1:B:34:GLU:HG3	5:B:698:HOH:O	2.16	0.45
1:B:63:GLY:N	1:B:64:PRO:CD	2.80	0.44
1:B:98:LEU:HG	1:B:114:PHE:CZ	2.52	0.44
1:B:140:CYS:O	1:B:169:TYR:HA	2.18	0.43
1:B:59:MET:SD	1:B:373:HIS:HB2	2.59	0.42
1:B:128:LEU:C	1:B:128:LEU:HD23	2.44	0.42
1:A:16:ARG:NE	5:A:608:HOH:O	2.39	0.41
1:A:398:ARG:NH2	5:A:634:HOH:O	2.53	0.41
1:B:310:TYR:CG	1:B:311:LEU:N	2.88	0.41
1:B:66:LEU:HB2	1:B:259:CYS:HB2	2.02	0.41
1:A:310:TYR:CG	1:A:311:LEU:N	2.89	0.41
1:B:230:ASP:OD2	3:B:502[B]:PMP:N1	2.55	0.40
1:B:344:MET:HE3	1:B:379:PHE:CE2	2.55	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	415/440 (94%)	399 (96%)	14 (3%)	2 (0%)	25	11
1	B	412/440 (94%)	399 (97%)	11 (3%)	2 (0%)	25	11
All	All	827/880 (94%)	798 (96%)	25 (3%)	4 (0%)	44	11

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	258[A]	LYS
1	A	258[B]	LYS
1	B	258[A]	LYS
1	B	258[B]	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	325/340 (96%)	322 (99%)	3 (1%)	75	63
1	B	322/340 (95%)	320 (99%)	2 (1%)	84	75
All	All	647/680 (95%)	642 (99%)	5 (1%)	79	68

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	GLU
1	A	231	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	416	LEU
1	B	34	GLU
1	B	231	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	185	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PMP	B	502[B]	-	16,16,16	2.30	3 (18%)	22,23,23	1.38	1 (4%)
2	PLP	A	501[A]	1	15,15,16	2.71	3 (20%)	21,22,23	1.57	5 (23%)
3	PMP	A	502[B]	-	16,16,16	2.55	3 (18%)	22,23,23	1.20	3 (13%)
2	PLP	B	501[A]	1	15,15,16	2.58	3 (20%)	21,22,23	1.68	6 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PMP	B	502[B]	-	-	2/8/8/8	0/1/1/1
2	PLP	A	501[A]	1	-	0/6/6/8	0/1/1/1
3	PMP	A	502[B]	-	-	2/8/8/8	0/1/1/1
2	PLP	B	501[A]	1	-	0/6/6/8	0/1/1/1

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	501[A]	PLP	C5-C4	7.36	1.48	1.40
2	A	501[A]	PLP	C5-C4	6.86	1.48	1.40
2	A	501[A]	PLP	C3-C2	6.47	1.47	1.41
3	A	502[B]	PMP	C3-C2	5.89	1.47	1.41
3	B	502[B]	PMP	C5-C4	5.55	1.48	1.40
3	B	502[B]	PMP	C3-C2	5.50	1.46	1.41
3	A	502[B]	PMP	C3-C4	5.48	1.48	1.40
3	A	502[B]	PMP	C5-C4	5.39	1.48	1.40
2	B	501[A]	PLP	C3-C4	4.63	1.48	1.40
2	B	501[A]	PLP	C3-C2	4.01	1.45	1.41
3	B	502[B]	PMP	C3-C4	3.89	1.45	1.40
2	A	501[A]	PLP	C3-C4	3.88	1.47	1.40

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501[A]	PLP	C4A-C4-C5	3.71	124.76	120.94
2	B	501[A]	PLP	C3-C4-C5	-3.18	114.78	118.59
2	B	501[A]	PLP	O3-C3-C4	3.08	126.13	118.10
2	A	501[A]	PLP	C6-C5-C4	2.84	120.42	118.10
3	B	502[B]	PMP	O3-C3-C2	2.72	123.23	117.58
2	A	501[A]	PLP	C3-C4-C5	-2.53	115.55	118.59
2	A	501[A]	PLP	O3-C3-C2	2.51	122.78	117.58
2	B	501[A]	PLP	C6-N1-C2	2.50	123.74	119.20
2	B	501[A]	PLP	C6-C5-C4	2.48	120.13	118.10
3	A	502[B]	PMP	O3-C3-C2	2.20	122.15	117.58
2	A	501[A]	PLP	O4P-C5A-C5	2.13	113.34	109.36
2	B	501[A]	PLP	C5A-C5-C6	-2.12	115.90	119.36
3	A	502[B]	PMP	C6-N1-C2	2.10	123.00	119.20
3	A	502[B]	PMP	C3-C4-C5	-2.05	116.87	118.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501[A]	PLP	O4P-C5A-C5	2.04	113.18	109.36

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	502[B]	PMP	C3-C4-C4A-N4A
3	A	502[B]	PMP	C5-C4-C4A-N4A
3	B	502[B]	PMP	C3-C4-C4A-N4A
3	B	502[B]	PMP	C5-C4-C4A-N4A

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	502[B]	PMP	2	0
2	A	501[A]	PLP	1	0
2	B	501[A]	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	414/440 (94%)	-0.44	3 (0%) 84 87	9, 18, 34, 55	3 (0%)
1	B	411/440 (93%)	-0.46	1 (0%) 92 94	9, 18, 34, 51	3 (0%)
All	All	825/880 (93%)	-0.45	4 (0%) 87 90	9, 18, 34, 55	6 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	408	ALA	2.7
1	B	246	HIS	2.1
1	A	416	LEU	2.1
1	A	417	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PLP	B	501[A]	15/16	0.98	0.04	15,16,17,17	15

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PMP	A	502[B]	16/16	0.98	0.04	12,13,14,14	16
3	PMP	B	502[B]	16/16	0.98	0.05	12,13,14,17	16
4	MG	A	503	1/1	0.98	0.06	22,22,22,22	0
2	PLP	A	501[A]	15/16	0.99	0.04	12,15,16,17	15

6.5 Other polymers [i](#)

There are no such residues in this entry.