



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:12 PM UTC

PDB ID : 6HBZ / pdb_00006hbz
Title : Bdellovibrio bacteriovorus DgcB Full-length
Authors : Lovering, A.L.; Meek, R.W.
Deposited on : 2018-08-13
Resolution : 1.79 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	279	12% 92% • • •
1	B	279	7% 95% • •

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4744 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GGDEF domain protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	269	Total	C	N	O	S	0	3	0
			2099	1336	360	400	3			
1	B	270	Total	C	N	O	S	0	2	0
			2093	1329	361	400	3			

- Molecule 2 is 9,9'-[(2R,3R,3aS,5S,7aR,9R,10R,10aS,12S,14aR)-3,5,10,12-tetrahydroxy-5,12-dioxidoctahydro-2H,7H-difuro[3,2-d:3',2'-j][1,3,7,9,2,8]tetraoxadiphosphacyclododecine-2,9-diyl]bis(2-amino-1,9-dihydro-6H-purin-6-one) (CCD ID: C2E) (formula: C₂₀H₂₄N₁₀O₁₄P₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Cl 2 2	0	0
3	B	3	Total Cl 3 3	0	0

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	269	Total O 269 269	0	0
5	B	170	Total O 170 170	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: GGDEF domain protein



- Molecule 1: GGDEF domain protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	176.78Å 176.78Å 112.12Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	153.09 – 1.79 153.09 – 1.79	Depositor EDS
% Data completeness (in resolution range)	100.0 (153.09-1.79) 100.0 (153.09-1.79)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 1.78Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.179 , 0.215 0.187 , 0.220	Depositor DCC
R_{free} test set	5021 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.148	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4744	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL, EDO, C2E

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.74	0/2141	0.85	2/2895 (0.1%)
1	B	0.69	0/2134	0.79	0/2883
All	All	0.72	0/4275	0.82	2/5778 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	296	TYR	CB-CA-C	5.64	121.50	110.67
1	A	106	ILE	CB-CA-C	5.17	116.72	110.78

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2099	0	2125	8	0
1	B	2093	0	2125	4	0
2	A	92	0	42	0	0
3	A	2	0	0	0	0
3	B	3	0	0	0	0
4	B	16	0	24	0	1
5	A	269	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	170	0	0	1	1
All	All	4744	0	4316	12	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (12) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:81:PHE:O	5:B:501:HOH:O	1.79	0.98
1:A:298:SER:HB2	1:A:307:THR:HG23	1.77	0.66
1:B:75:SER:H	1:B:78:HIS:HD2	1.43	0.66
1:A:298:SER:CB	1:A:307:THR:HG23	2.36	0.55
1:A:69:ILE:HB	1:A:74[B]:LEU:CD1	2.39	0.53
1:B:78:HIS:HE1	1:B:98:LYS:O	1.90	0.53
1:A:215:LYS:NZ	5:A:504:HOH:O	2.41	0.52
1:B:229:GLU:HG2	1:B:229:GLU:O	2.12	0.47
1:A:124:ASN:ND2	5:A:512:HOH:O	2.49	0.45
1:A:295:LEU:HD23	1:A:295:LEU:C	2.43	0.44
1:A:308:ILE:O	1:A:309:VAL:HB	2.18	0.43
1:A:244:GLY:HA3	1:A:308:ILE:HD11	2.03	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:609:HOH:O	5:B:609:HOH:O[9_555]	0.53	1.67
4:B:405:EDO:O2	4:B:405:EDO:O2[9_555]	1.96	0.24

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/279 (96%)	262 (98%)	5 (2%)	1 (0%)	30	19
1	B	268/279 (96%)	261 (97%)	7 (3%)	0	100	100
All	All	536/558 (96%)	523 (98%)	12 (2%)	1 (0%)	43	31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	145	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/234 (99%)	229 (99%)	2 (1%)	70	67
1	B	230/234 (98%)	228 (99%)	2 (1%)	70	67
All	All	461/468 (98%)	457 (99%)	4 (1%)	70	67

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	298	SER
1	A	307	THR
1	B	220	ASN
1	B	229	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	ASN
1	A	148	GLN
1	A	193	ASN
1	B	78	HIS
1	B	188	HIS
1	B	193	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	220	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	C2E	A	401	-	52,52,52	3.56	26 (50%)	78,82,82	1.89	24 (30%)
4	EDO	B	406	-	3,3,3	0.43	0	2,2,2	0.52	0
4	EDO	B	407	-	3,3,3	0.39	0	2,2,2	0.31	0
4	EDO	B	405	-	3,3,3	0.77	0	2,2,2	0.27	0
4	EDO	B	404	-	3,3,3	0.25	0	2,2,2	0.51	0
2	C2E	A	402	-	52,52,52	3.37	25 (48%)	78,82,82	1.87	18 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	C2E	A	401	-	-	0/30/62/62	0/6/7/7
4	EDO	B	406	-	-	0/1/1/1	-
4	EDO	B	407	-	-	1/1/1/1	-
4	EDO	B	405	-	-	1/1/1/1	-
4	EDO	B	404	-	-	0/1/1/1	-
2	C2E	A	402	-	-	0/30/62/62	0/6/7/7

All (51) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	C2E	C2'-C3'	-12.70	1.25	1.53
2	A	402	C2E	C2'-C3'	-11.45	1.28	1.53
2	A	401	C2E	C3A-C4A	-9.68	1.27	1.52
2	A	402	C2E	C3A-C4A	-9.41	1.28	1.52
2	A	401	C2E	O4A-C4A	7.29	1.61	1.45
2	A	402	C2E	O4A-C4A	7.19	1.61	1.45
2	A	402	C2E	C4-N3	5.94	1.47	1.34
2	A	401	C2E	C4-N3	5.77	1.47	1.34
2	A	402	C2E	C41-N31	5.32	1.46	1.34
2	A	401	C2E	O4A-C1A	-4.82	1.30	1.42
2	A	401	C2E	C2-N3	4.76	1.44	1.33
2	A	401	C2E	C41-N91	-4.55	1.26	1.38
2	A	401	C2E	C41-N31	4.53	1.44	1.34
2	A	401	C2E	C5'-C4'	-4.49	1.38	1.51
2	A	402	C2E	C5'-C4'	-4.49	1.38	1.51
2	A	402	C2E	O4A-C1A	-4.28	1.32	1.42
2	A	401	C2E	C2'-C1'	4.26	1.66	1.53
2	A	402	C2E	C2-N2	4.24	1.44	1.34
2	A	402	C2E	C21-N31	3.91	1.42	1.33
2	A	402	C2E	C2-N3	3.91	1.42	1.33
2	A	401	C2E	C21-N31	3.89	1.42	1.33
2	A	402	C2E	C2'-C1'	3.87	1.65	1.53
2	A	401	C2E	C51-N71	-3.80	1.31	1.39
2	A	401	C2E	C2-N2	3.73	1.42	1.34
2	A	401	C2E	C5-N7	-3.68	1.31	1.39
2	A	401	C2E	C21-N21	3.62	1.42	1.34
2	A	402	C2E	C21-N21	3.60	1.42	1.34
2	A	402	C2E	C41-N91	-3.59	1.28	1.38
2	A	402	C2E	C5-N7	-3.49	1.32	1.39
2	A	401	C2E	O4'-C1'	-3.26	1.34	1.42
2	A	401	C2E	C3'-C4'	3.13	1.61	1.52
2	A	402	C2E	C3'-C4'	2.94	1.60	1.52
2	A	401	C2E	O3A-C3A	2.93	1.54	1.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	402	C2E	C51-N71	-2.77	1.33	1.39
2	A	401	C2E	O4'-C4'	2.77	1.51	1.45
2	A	402	C2E	P11-O3'	2.76	1.67	1.59
2	A	401	C2E	P11-O3'	2.74	1.67	1.59
2	A	402	C2E	O61-C61	-2.72	1.18	1.23
2	A	402	C2E	P1-O3A	2.67	1.67	1.59
2	A	401	C2E	O3'-C3'	2.67	1.53	1.44
2	A	401	C2E	P1-O3A	2.61	1.67	1.59
2	A	402	C2E	C2-N1	2.61	1.44	1.37
2	A	402	C2E	O3A-C3A	2.49	1.52	1.44
2	A	402	C2E	O3'-C3'	2.39	1.52	1.44
2	A	401	C2E	C2-N1	2.37	1.43	1.37
2	A	401	C2E	O6-C6	-2.37	1.19	1.23
2	A	402	C2E	C51-C61	2.34	1.53	1.44
2	A	402	C2E	O4'-C4'	2.31	1.50	1.45
2	A	401	C2E	C1'-N9	-2.23	1.41	1.47
2	A	401	C2E	C51-C61	2.15	1.52	1.44
2	A	402	C2E	O4'-C1'	-2.13	1.37	1.42

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	C2E	C5-C4-N3	-5.61	119.47	128.39
2	A	402	C2E	C5-C4-N3	-4.90	120.60	128.39
2	A	401	C2E	N9-C4-N3	4.30	134.55	125.95
2	A	402	C2E	C1A-N91-C81	-4.27	114.59	126.73
2	A	402	C2E	C1'-N9-C4	-4.23	113.99	126.49
2	A	401	C2E	N91-C81-N71	-4.19	105.63	113.40
2	A	401	C2E	C1A-N91-C81	-4.10	115.07	126.73
2	A	402	C2E	N9-C4-N3	3.89	133.73	125.95
2	A	402	C2E	C1'-N9-C8	3.67	137.17	126.73
2	A	401	C2E	C1'-N9-C4	-3.64	115.74	126.49
2	A	402	C2E	N91-C81-N71	-3.59	106.74	113.40
2	A	402	C2E	O3A-P1-O1P	-3.43	98.84	109.81
2	A	401	C2E	O3A-P1-O1P	-3.38	99.01	109.81
2	A	401	C2E	C2-N3-C4	3.29	117.97	112.30
2	A	401	C2E	C1'-N9-C8	3.29	136.08	126.73
2	A	401	C2E	C1A-N91-C41	3.15	135.79	126.49
2	A	402	C2E	C1A-N91-C41	3.11	135.67	126.49
2	A	402	C2E	C2-N3-C4	3.07	117.59	112.30
2	A	402	C2E	O4A-C1A-N91	3.00	115.16	108.36
2	A	402	C2E	C61-C51-N71	2.97	135.69	130.29

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	C2E	C2-N1-C6	-2.90	119.85	125.11
2	A	402	C2E	C21-N31-C41	2.51	116.63	112.30
2	A	401	C2E	C61-C51-N71	2.47	134.79	130.29
2	A	401	C2E	C6-C5-C4	2.45	122.51	118.83
2	A	401	C2E	C51-C41-N91	2.42	109.99	105.66
2	A	401	C2E	O21-P11-O3'	2.40	116.48	106.70
2	A	401	C2E	C21-N31-C41	2.40	116.43	112.30
2	A	402	C2E	N9-C8-N7	-2.35	109.04	113.40
2	A	401	C2E	C4A-O4A-C1A	-2.32	104.34	109.47
2	A	401	C2E	C81-N71-C51	2.23	108.23	104.26
2	A	401	C2E	C41-C51-N71	-2.22	107.16	110.67
2	A	401	C2E	N9-C8-N7	-2.21	109.30	113.40
2	A	402	C2E	C6-C5-C4	2.19	122.12	118.83
2	A	402	C2E	O21-P11-O3'	2.17	115.51	106.70
2	A	402	C2E	C4A-O4A-C1A	-2.10	104.83	109.47
2	A	402	C2E	C51-C41-N31	-2.09	125.06	128.39
2	A	401	C2E	O3'-P11-O11	-2.09	103.12	109.81
2	A	401	C2E	N21-C21-N11	2.08	121.15	116.76
2	A	401	C2E	C3'-C2'-C1'	2.05	104.40	99.89
2	A	401	C2E	O2P-P1-O3A	2.03	114.97	106.70
2	A	402	C2E	C51-C61-N11	2.01	118.38	113.25
2	A	401	C2E	O3'-C3'-C4'	-2.01	102.94	110.03

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	405	EDO	O1-C1-C2-O2
4	B	407	EDO	O1-C1-C2-O2


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	405	EDO	0	1

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be

highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	269/279 (96%)	0.54	34 (12%) 8 6	11, 29, 69, 113	3 (1%)
1	B	270/279 (96%)	0.32	19 (7%) 22 21	14, 32, 60, 80	3 (1%)
All	All	539/558 (96%)	0.43	53 (9%) 13 11	11, 31, 65, 113	6 (1%)

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	309	VAL	8.8
1	A	152	LEU	8.0
1	A	144	TYR	7.9
1	A	155	ALA	7.5
1	A	153	THR	6.6
1	A	151	ALA	6.6
1	A	154	GLY	6.6
1	A	147	ALA	6.4
1	B	134	SER	6.0
1	B	264[A]	LYS	5.6
1	B	32	VAL	5.2
1	A	150	ASP	5.2
1	B	144	TYR	4.5
1	B	310	ALA	4.3
1	A	149	LYS	4.2
1	A	32	VAL	3.9
1	A	156	HIS	3.9
1	A	306	THR	3.8
1	A	146	ARG	3.7
1	A	134	SER	3.6
1	A	145	ASP	3.6
1	A	308	ILE	3.3
1	A	296	TYR	3.2
1	B	145	ASP	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	300	GLN	3.0
1	A	265	LYS	3.0
1	A	303	ARG	2.9
1	B	263	GLY	2.8
1	B	85	GLY	2.8
1	A	158	LYS	2.8
1	B	53	THR	2.8
1	A	43	PRO	2.7
1	A	157	SER	2.7
1	A	148	GLN	2.7
1	A	304	ASN	2.7
1	A	106	ILE	2.7
1	A	132	LYS	2.6
1	A	305	ARG	2.6
1	A	301	GLY	2.5
1	B	54	ALA	2.5
1	A	302	GLY	2.5
1	B	86	SER	2.5
1	B	147	ALA	2.5
1	B	178	PRO	2.3
1	A	307	THR	2.3
1	B	309	VAL	2.3
1	B	132	LYS	2.2
1	A	298	SER	2.2
1	B	283	GLU	2.1
1	B	266	ILE	2.1
1	B	133	GLY	2.0
1	B	226	TYR	2.0
1	A	299	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	EDO	B	405	4/4	0.85	0.25	76,80,89,89	0
4	EDO	B	406	4/4	0.90	0.14	45,48,55,60	0
4	EDO	B	407	4/4	0.94	0.13	49,51,53,58	0
4	EDO	B	404	4/4	0.95	0.17	39,43,46,48	0
3	CL	B	401	1/1	0.96	0.09	54,54,54,54	0
3	CL	B	402	1/1	0.96	0.15	58,58,58,58	0
2	C2E	A	401	46/46	0.97	0.06	22,28,33,37	0
2	C2E	A	402	46/46	0.98	0.05	21,26,30,34	0
3	CL	A	404	1/1	0.99	0.03	31,31,31,31	0
3	CL	A	403	1/1	1.00	0.01	22,22,22,22	0
3	CL	B	403	1/1	1.00	0.04	29,29,29,29	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around C2E A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around C2E A 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.