



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 05:50 PM UTC

PDB ID : 4HCC / pdb_00004hcc
Title : The zinc ion bound form of crystal structure of E.coli ExoI-ssDNA complex
Authors : Qiu, R.; Wei, J.; Lou, T.; Liu, M.; Ji, C.; Gong, W.
Deposited on : 2012-09-29
Resolution : 2.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1130 (2.98-2.94)
Clashscore	190562	1157 (2.98-2.94)
Ramachandran outliers	187476	1101 (2.98-2.94)
Sidechain outliers	187428	1101 (2.98-2.94)
RSRZ outliers	180081	1130 (2.98-2.94)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	481	
1	B	481	
2	C	12	
2	D	12	

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8059 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Exodeoxyribonuclease I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	465	Total	C	N	O	S	22	2	0
			3792	2421	665	692	14			
1	B	463	Total	C	N	O	S	14	2	0
			3776	2411	662	689	14			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	472	ASP	GLU	SEE REMARK 999	UNP P04995
A	476	HIS	-	expression tag	UNP P04995
A	477	HIS	-	expression tag	UNP P04995
A	478	HIS	-	expression tag	UNP P04995
A	479	HIS	-	expression tag	UNP P04995
A	480	HIS	-	expression tag	UNP P04995
A	481	HIS	-	expression tag	UNP P04995
B	472	ASP	GLU	SEE REMARK 999	UNP P04995
B	476	HIS	-	expression tag	UNP P04995
B	477	HIS	-	expression tag	UNP P04995
B	478	HIS	-	expression tag	UNP P04995
B	479	HIS	-	expression tag	UNP P04995
B	480	HIS	-	expression tag	UNP P04995
B	481	HIS	-	expression tag	UNP P04995

- Molecule 2 is a DNA chain called DNA (5'-D(*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	10	Total	C	N	O	P	0	0	0
			207	100	50	48	9			
2	D	10	Total	C	N	O	P	0	0	0
			207	100	50	48	9			

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		
4	B	1	Total	Zn	0	0
			1	1		

- Molecule 5 is ISOPROPYL ALCOHOL (CCD ID: IPA) (formula: C₃H₈O).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	3	1		
5	B	1	Total	C	O	0	0
			4	3	1		
5	B	1	Total	C	O	0	0
			4	3	1		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	O	P	0	0
			4	3	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	D	1	Total	O	P	0	0
			4	3	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	8	Total	O	0	0
			8	8		
7	B	11	Total	O	0	0
			11	11		
7	C	1	Total	O	0	0
			1	1		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Exodeoxyribonuclease I



• Molecule 1: Exodeoxyribonuclease I




• Molecule 2: DNA (5'-D(*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*A)-3')



• Molecule 2: DNA (5'-D(*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*A)-3')





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	63.04Å 108.69Å 158.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.96 30.00 – 2.96	Depositor EDS
% Data completeness (in resolution range)	98.5 (30.00-2.96) 98.5 (30.00-2.96)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	8.63 (at 2.95Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.185 , 0.245 0.182 , 0.245	Depositor DCC
R_{free} test set	1188 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	60.0	Xtriage
Anisotropy	0.030	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 22.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8059	wwPDB-VP
Average B, all atoms (Å ²)	62.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, PO4, IPA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/3891	0.80	0/5293
1	B	0.69	0/3874	0.85	0/5270
2	C	0.31	0/235	0.66	0/358
2	D	0.31	0/235	0.70	0/358
All	All	0.66	0/8235	0.82	0/11279

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3792	0	3695	34	0
1	B	3776	0	3679	38	0
2	C	207	0	112	4	0
2	D	207	0	112	4	0
3	A	15	0	0	0	0
3	B	20	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	4	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	8	0	16	0	0
6	C	4	0	0	1	0
6	D	4	0	0	0	0
7	A	8	0	0	0	0
7	B	11	0	0	0	0
7	C	1	0	0	0	0
All	All	8059	0	7622	72	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:348:VAL:HG11	1:B:348:VAL:HG11	1.42	0.98
1:B:209:PHE:O	1:B:212:ARG:HG2	1.77	0.83
1:A:374:ASP:OD1	1:A:377:ARG:NH1	2.10	0.82
2:C:12:DA:H8	2:C:12:DA:H5''	1.45	0.80
1:A:348:VAL:HG11	1:B:348:VAL:CG1	2.20	0.69
1:B:111:VAL:O	1:B:115:ILE:HD13	1.93	0.68
1:B:396:ILE:HD13	1:B:404:GLU:OE2	1.94	0.68
1:B:363:VAL:HA	1:B:366:GLN:HG3	1.78	0.63
1:B:287:LYS:NZ	2:D:8:DA:OP2	2.30	0.63
1:A:64:VAL:HG21	1:A:467:LEU:HD21	1.82	0.60
1:A:13:PHE:O	1:A:33:ALA:HA	2.03	0.59
2:C:12:DA:H5''	2:C:12:DA:C8	2.34	0.59
1:A:175:ILE:HD13	1:A:192:ALA:HB1	1.84	0.59
1:A:249:LEU:HD12	1:A:347:VAL:HG21	1.85	0.58
1:A:22:HIS:CE1	1:A:24:ALA:HB3	2.41	0.56
1:A:184:MET:HE3	1:A:184:MET:HA	1.87	0.55
1:B:199:THR:HG22	1:B:200:ARG:HD3	1.91	0.53
1:B:34:ILE:HG12	1:B:46:PRO:HB3	1.91	0.52
1:B:162:PRO:HD2	1:B:238:ALA:HB1	1.92	0.52
1:B:269:SER:N	1:B:270:PRO:HD2	2.25	0.51
2:C:12:DA:H5'	6:C:101:PO4:O3	2.11	0.51
1:B:248:PRO:HB3	1:B:261:MET:HE2	1.92	0.51
1:A:348:VAL:CG1	1:B:348:VAL:HG11	2.28	0.50
1:B:120:PHE:HD1	1:B:380:MET:HG2	1.75	0.50
1:B:105:VAL:HG23	1:B:136:ASP:HB2	1.94	0.50
1:B:34:ILE:HG13	1:B:184:MET:HE1	1.93	0.50
1:A:214:LYS:HB2	1:A:303:ILE:HD12	1.94	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:287:LYS:HZ3	1:B:287:LYS:HB2	1.77	0.48
1:A:40:PHE:HE2	1:A:198:LYS:HD2	1.78	0.48
1:A:249:LEU:HG	1:A:262[A]:VAL:HG23	1.96	0.48
1:A:374:ASP:HA	1:A:377:ARG:NH1	2.28	0.48
1:B:17:GLU:HG2	1:B:183:ALA:HB2	1.96	0.48
1:B:286:ALA:HA	2:D:8:DA:OP2	2.14	0.48
1:B:137:LEU:O	1:B:141:MET:HG2	2.13	0.48
1:B:64:VAL:HG21	1:B:467:LEU:HD21	1.95	0.48
1:A:377:ARG:O	1:A:378:ALA:C	2.55	0.47
1:B:401:LYS:HD3	1:B:401:LYS:HA	1.77	0.47
2:D:8:DA:H5''	2:D:8:DA:H8	1.80	0.47
1:A:345:GLU:HG3	1:B:349:ALA:CB	2.45	0.46
2:D:4:DA:N3	2:D:4:DA:H2'	2.30	0.46
1:B:22:HIS:CE1	1:B:24:ALA:HB3	2.50	0.45
1:A:391:LEU:N	1:A:392:PRO:HD2	2.32	0.45
1:A:22:HIS:HE1	1:A:24:ALA:HB3	1.81	0.45
1:A:447:LEU:O	1:A:451:VAL:HG23	2.18	0.44
1:A:347:VAL:HA	1:A:350:ILE:HD12	1.99	0.44
1:A:241:GLY:HA3	1:A:323:LEU:HD22	2.00	0.44
1:B:459:GLU:CD	1:B:459:GLU:H	2.26	0.44
1:A:145:TYR:CE2	1:A:162:PRO:HG3	2.53	0.43
1:B:249:LEU:HD22	1:B:249:LEU:HA	1.84	0.43
1:A:345:GLU:HG3	1:B:349:ALA:HB1	1.99	0.43
1:B:150:GLU:O	1:B:201:GLN:NE2	2.51	0.43
1:B:475:VAL:HG23	1:B:476:HIS:CE1	2.54	0.43
1:A:58:LEU:HD21	1:A:471:ALA:HA	2.00	0.43
1:A:388:PRO:HA	1:A:391:LEU:HD12	2.00	0.43
1:B:28:PRO:HG3	1:B:115:ILE:HD11	2.01	0.43
1:B:79:GLU:H	1:B:79:GLU:HG2	1.73	0.43
1:A:34:ILE:HD11	1:A:184:MET:HE2	2.01	0.43
1:B:215:HIS:ND1	2:C:1:DA:H5'	2.33	0.43
1:A:409:ASN:O	1:A:413:ARG:HG3	2.19	0.43
1:B:102:TYR:O	1:B:103:ASN:HB3	2.19	0.42
1:A:383:VAL:HG13	1:A:391:LEU:HD21	2.02	0.42
1:B:249:LEU:HD13	1:B:344:ARG:HG3	2.00	0.42
1:A:398:PHE:HB3	1:A:403:ILE:HD12	2.01	0.42
1:A:261:MET:HG3	1:A:301:VAL:CG2	2.51	0.41
1:B:439:PHE:HD2	1:B:440:LEU:HD23	1.84	0.41
1:A:154:TRP:HA	1:A:155:PRO:HD2	1.98	0.41
1:B:208:LEU:HD22	1:B:308:VAL:HG22	2.02	0.41
1:B:274:LEU:HD12	1:B:278:THR:HB	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:ALA:C	1:A:240:ARG:H	2.30	0.40
1:A:280:ARG:NH2	1:A:353:GLU:HB2	2.36	0.40
1:B:384[A]:LEU:HD12	1:B:384[A]:LEU:HA	1.74	0.40
1:A:40:PHE:CE2	1:A:198:LYS:HD2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	461/481 (96%)	437 (95%)	24 (5%)	0	100	100
1	B	459/481 (95%)	445 (97%)	13 (3%)	1 (0%)	43	66
All	All	920/962 (96%)	882 (96%)	37 (4%)	1 (0%)	48	72

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	276	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	402/413 (97%)	388 (96%)	14 (4%)	32	57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	400/413 (97%)	381 (95%)	19 (5%)	23	48
All	All	802/826 (97%)	769 (96%)	33 (4%)	28	53

All (33) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	36	THR
1	A	137	LEU
1	A	144	CYS
1	A	213	ASN
1	A	216	LYS
1	A	230	VAL
1	A	244	SER
1	A	269	SER
1	A	287	LYS
1	A	331	LEU
1	A	332	ASP
1	A	384[A]	LEU
1	A	384[B]	LEU
1	A	456	ASP
1	B	36	THR
1	B	38	SER
1	B	100	LEU
1	B	137	LEU
1	B	150	GLU
1	B	152	ILE
1	B	156	GLU
1	B	170	THR
1	B	249	LEU
1	B	287	LYS
1	B	353	GLU
1	B	384[A]	LEU
1	B	384[B]	LEU
1	B	394	LEU
1	B	396	ILE
1	B	421	TYR
1	B	428	LEU
1	B	445	ASP
1	B	451	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	HIS
1	A	89	HIS
1	A	114	ASN
1	A	157	ASN
1	A	242	ASN
1	B	8	GLN
1	B	14	HIS
1	B	30	GLN
1	B	130	HIS
1	B	225	GLN
1	B	242	ASN
1	B	293	ASN
1	B	302	HIS
1	B	313	ASN
1	B	476	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 14 ligands modelled in this entry, 2 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	B	504	-	4,4,4	0.43	0	6,6,6	0.16	0
3	SO4	A	502	-	4,4,4	0.46	0	6,6,6	0.09	0
5	IPA	A	505	-	3,3,3	0.56	0	3,3,3	0.24	0
5	IPA	B	506	-	3,3,3	0.56	0	3,3,3	0.10	0
3	SO4	A	501	-	4,4,4	0.45	0	6,6,6	0.09	0
5	IPA	B	507	-	3,3,3	0.56	0	3,3,3	0.19	0
6	PO4	C	101	4,2	0,3,4	-	-	0,3,6	-	-
6	PO4	D	101	4,2	0,3,4	-	-	0,3,6	-	-
3	SO4	B	502	-	4,4,4	0.43	0	6,6,6	0.19	0
3	SO4	B	503	-	4,4,4	0.45	0	6,6,6	0.14	0
3	SO4	A	503	-	4,4,4	0.47	0	6,6,6	0.12	0
3	SO4	B	501	-	4,4,4	0.42	0	6,6,6	0.24	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	C	101	PO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	465/481 (96%)	-0.48	1 (0%) 91 90	25, 60, 94, 118	9 (1%)
1	B	463/481 (96%)	-0.51	1 (0%) 91 90	25, 56, 96, 126	8 (1%)
2	C	10/12 (83%)	-0.26	0 100 100	64, 91, 100, 113	0
2	D	10/12 (83%)	-0.46	0 100 100	68, 73, 99, 102	0
All	All	948/986 (96%)	-0.49	2 (0%) 91 90	25, 58, 96, 126	17 (1%)

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	421	TYR	2.5
1	A	356	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	502	5/5	0.75	0.09	116,117,122,123	0
3	SO4	B	503	5/5	0.76	0.13	105,110,118,120	0
5	IPA	A	505	4/4	0.83	0.29	62,64,68,74	0
3	SO4	A	503	5/5	0.84	0.12	115,115,119,119	0
5	IPA	B	507	4/4	0.85	0.19	63,64,66,67	0
3	SO4	B	502	5/5	0.90	0.17	96,98,102,102	0
3	SO4	A	501	5/5	0.92	0.09	100,101,102,106	0
6	PO4	C	101	4/5	0.92	0.08	88,88,93,94	0
3	SO4	B	504	5/5	0.93	0.13	112,112,118,119	0
5	IPA	B	506	4/4	0.94	0.12	52,53,55,56	0
6	PO4	D	101	4/5	0.96	0.06	71,72,73,74	0
3	SO4	B	501	5/5	0.98	0.05	47,51,53,54	0
4	ZN	A	504	1/1	0.99	0.06	83,83,83,83	0
4	ZN	B	505	1/1	1.00	0.04	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.