



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 01:04 AM UTC

PDB ID : 6HCN / pdb_00006hcn
Title : Adenovirus Type 5 Fiber Knob protein at 1.49Å resolution
Authors : Rizkallah, P.J.; Parker, A.L.; Baker, A.T.
Deposited on : 2018-08-15
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	186	9% 87% 12%
1	C	186	16% 87% 12%
2	B	188	15% 90% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	A	602	-	-	X	-
3	EDO	A	603	-	-	X	-
3	EDO	A	604	-	-	X	-
3	EDO	C	602	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Fiber protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	186	Total	C	N	O	S	0	5	0
			1470	929	242	294	5			
1	C	186	Total	C	N	O	S	0	1	0
			1439	912	236	286	5			

- Molecule 2 is a protein called Fiber protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	188	Total	C	N	O	S	0	5	0
			1486	939	245	297	5			

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total	Mg	0	0
			1	1		



- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	161	Total	O	0	0
			161	161		
5	B	101	Total	O	0	0
			101	101		
5	C	147	Total	O	0	0
			147	147		



3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Fiber protein



- Molecule 1: Fiber protein



- Molecule 2: Fiber protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	102.16Å 102.44Å 77.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	61.56 – 1.49 61.56 – 1.49	Depositor EDS
% Data completeness (in resolution range)	99.8 (61.56-1.49) 99.8 (61.56-1.49)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.24 (at 1.49Å)	Xtriage
Refinement program	REFMAC 5.8.0230	Depositor
R, R_{free}	0.211 , 0.233 0.210 , 0.231	Depositor DCC
R_{free} test set	6391 reflections (4.84%)	wwPDB-VP
Wilson B-factor (Å ²)	22.3	Xtriage
Anisotropy	0.755	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.025 for k,h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4825	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.96	1/1504 (0.1%)	1.02	1/2052 (0.0%)
1	C	0.88	1/1473 (0.1%)	0.97	0/2010
2	B	0.93	0/1520	1.00	1/2071 (0.0%)
All	All	0.92	2/4497 (0.0%)	1.00	2/6133 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	C	0	2
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	456	HIS	CE1-NE2	5.13	1.37	1.32
1	C	576	SER	C-O	5.03	1.30	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	540	GLN	CB-CG-CD	6.05	122.88	112.60
2	B	544	ASP	CA-CB-CG	5.29	117.89	112.60

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	543	GLY	Peptide
1	C	481	ARG	Sidechain
1	C	543	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1470	0	1427	23	0
1	C	1439	0	1402	19	0
2	B	1486	0	1446	19	0
3	A	16	0	24	15	0
3	C	4	0	6	6	0
4	C	1	0	0	0	0
5	A	161	0	0	3	0
5	B	101	0	0	1	0
5	C	147	0	0	1	0
All	All	4825	0	4305	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:506[A]:LYS:HD2	2:B:506[A]:LYS:H	1.02	1.07
2:B:506[A]:LYS:HD2	2:B:506[A]:LYS:N	1.74	1.01
2:B:506[A]:LYS:H	2:B:506[A]:LYS:CD	1.62	0.97
1:C:511:THR:HG22	3:C:602:EDO:H11	1.65	0.76
2:B:506[B]:LYS:HD2	2:B:507:SER:N	2.03	0.74
1:C:511:THR:CG2	3:C:602:EDO:H11	2.18	0.73
2:B:453:GLN:HE21	2:B:559:SER:HA	1.54	0.72
1:A:501:LEU:HD22	3:A:601:EDO:H12	1.70	0.71
1:A:489:THR:HG22	5:A:743:HOH:O	1.94	0.66
1:A:470:ASN:HD22	3:A:604:EDO:H12	1.62	0.65
1:A:468:LEU:HD12	3:A:603:EDO:H21	1.79	0.65
1:A:546:THR:CG2	3:A:604:EDO:H22	2.26	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:562:ASN:HD22	1:A:562:ASN:C	2.05	0.64
1:A:414[B]:ASN:HD22	1:A:414[B]:ASN:C	2.04	0.63
1:C:528:LYS:NZ	1:C:566:GLU:OE2	2.28	0.62
2:B:562:ASN:C	2:B:562:ASN:HD22	2.09	0.60
1:A:546:THR:HG21	3:A:604:EDO:H22	1.85	0.59
2:B:453:GLN:NE2	2:B:559:SER:HA	2.18	0.58
1:C:562:ASN:C	1:C:562:ASN:HD22	2.10	0.58
1:C:511:THR:CG2	3:C:602:EDO:C1	2.83	0.56
1:A:524:GLY:O	1:C:540:GLN:OE1	2.24	0.55
2:B:394:LYS:C	2:B:396:ASN:H	2.15	0.55
1:A:515:ASN:OD1	1:A:540:GLN:HG2	2.07	0.55
1:A:470:ASN:HD22	3:A:604:EDO:C1	2.19	0.54
3:A:602:EDO:H21	1:C:513:LYS:NZ	2.23	0.54
2:B:456:HIS:CE1	2:B:555[B]:SER:HG	2.27	0.53
2:B:416[A]:GLU:O	2:B:416[A]:GLU:OE2	2.26	0.53
1:A:526:LYS:HB2	1:C:531:THR:HG21	1.93	0.51
2:B:506[A]:LYS:HG2	5:B:695:HOH:O	2.11	0.50
1:A:414[B]:ASN:C	1:A:414[B]:ASN:ND2	2.69	0.49
1:A:420:LYS:HB2	3:A:602:EDO:H12	1.94	0.49
2:B:501:LEU:HD22	2:B:506[A]:LYS:HG3	1.95	0.49
1:C:396:ASN:CG	1:C:487:GLU:OE2	2.55	0.49
1:C:511:THR:HG23	3:C:602:EDO:H12	1.96	0.48
1:A:546:THR:HG23	3:A:604:EDO:H22	1.96	0.48
3:A:603:EDO:H22	3:A:604:EDO:H11	1.97	0.47
1:A:420:LYS:CB	3:A:602:EDO:H12	2.44	0.47
1:C:565:ASN:ND2	5:C:703:HOH:O	2.44	0.47
1:C:460:ARG:NE	1:C:546:THR:OG1	2.48	0.47
1:A:552[B]:MET:HB3	1:A:552[B]:MET:HE3	1.41	0.46
2:B:395:ASN:O	2:B:395:ASN:ND2	2.49	0.46
1:A:474[B]:ASP:OD1	1:A:475:PRO:HD2	2.16	0.45
3:A:603:EDO:H22	3:A:604:EDO:C1	2.46	0.45
1:A:474[B]:ASP:OD2	5:A:701:HOH:O	2.20	0.44
3:A:602:EDO:H11	3:C:602:EDO:H21	1.99	0.44
1:A:526:LYS:CB	1:C:531:THR:HG21	2.47	0.44
2:B:562:ASN:C	2:B:562:ASN:ND2	2.77	0.43
1:C:511:THR:HG23	3:C:602:EDO:C1	2.49	0.43
2:B:562:ASN:HD22	2:B:564:ILE:H	1.67	0.42
1:A:562:ASN:HD22	1:A:564:ILE:H	1.67	0.42
2:B:531:THR:HG21	1:C:526:LYS:HB2	2.01	0.42
1:C:562:ASN:HD22	1:C:564:ILE:H	1.67	0.42
3:A:601:EDO:H21	5:A:822:HOH:O	2.20	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:426:THR:HG21	1:C:428:CYS:O	2.20	0.41
1:A:396:ASN:OD1	1:A:398:LYS:HD3	2.21	0.41
1:C:479:ASN:HB3	1:C:486:THR:HB	2.03	0.41
3:A:603:EDO:H22	3:A:604:EDO:H22	2.02	0.41
2:B:394:LYS:C	2:B:396:ASN:N	2.78	0.41
2:B:512:ALA:HB2	1:C:523:ASN:HA	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	189/186 (102%)	182 (96%)	7 (4%)	0	100	100
1	C	185/186 (100%)	175 (95%)	9 (5%)	1 (0%)	24	8
2	B	191/188 (102%)	183 (96%)	7 (4%)	1 (0%)	24	8
All	All	565/560 (101%)	540 (96%)	23 (4%)	2 (0%)	30	12

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	541	GLU
1	C	541	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	165/160 (103%)	160 (97%)	5 (3%)	36	9
1	C	161/160 (101%)	156 (97%)	5 (3%)	35	9
2	B	167/162 (103%)	160 (96%)	7 (4%)	26	4
All	All	493/482 (102%)	476 (97%)	17 (3%)	37	7

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	451	THR
1	A	482	ASN
1	A	506[A]	LYS
1	A	506[B]	LYS
1	A	540	GLN
2	B	416[A]	GLU
2	B	416[B]	GLU
2	B	482	ASN
2	B	506[A]	LYS
2	B	506[B]	LYS
2	B	507	SER
2	B	544	ASP
1	C	449	SER
1	C	482	ASN
1	C	540	GLN
1	C	541	GLU
1	C	545	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	431	GLN
1	A	456	HIS
1	A	470	ASN
1	A	482	ASN
1	A	561	HIS
1	A	562	ASN
1	A	565	ASN
2	B	431	GLN
2	B	453	GLN
2	B	469	ASN
2	B	482	ASN
2	B	561	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	562	ASN
2	B	565	ASN
1	C	431	GLN
1	C	469	ASN
1	C	482	ASN
1	C	561	HIS
1	C	562	ASN
1	C	565	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	602	-	3,3,3	0.76	0	2,2,2	1.47	0
3	EDO	C	602	-	3,3,3	0.29	0	2,2,2	0.19	0
3	EDO	A	601	-	3,3,3	0.38	0	2,2,2	0.37	0
3	EDO	A	603	-	3,3,3	0.42	0	2,2,2	0.25	0
3	EDO	A	604	-	3,3,3	0.56	0	2,2,2	0.60	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	602	-	-	1/1/1/1	-
3	EDO	C	602	-	-	0/1/1/1	-
3	EDO	A	601	-	-	0/1/1/1	-
3	EDO	A	603	-	-	0/1/1/1	-
3	EDO	A	604	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	602	EDO	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 20 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	EDO	4	0
3	C	602	EDO	6	0
3	A	601	EDO	2	0
3	A	603	EDO	4	0
3	A	604	EDO	8	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	186/186 (100%)	0.78	17 (9%)	15 15	13, 29, 44, 68	5 (2%)
1	C	186/186 (100%)	1.18	30 (16%)	4 4	13, 32, 48, 78	1 (0%)
2	B	188/188 (100%)	1.04	29 (15%)	5 5	13, 32, 55, 80	5 (2%)
All	All	560/560 (100%)	1.00	76 (13%)	7 7	13, 31, 50, 80	11 (1%)

All (76) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	543	GLY	9.6
1	A	542	THR	9.0
1	A	543	GLY	8.9
1	C	542	THR	7.8
2	B	543	GLY	6.7
2	B	542	THR	6.6
2	B	545	THR	5.9
2	B	544	ASP	5.8
1	A	541	GLU	5.6
1	C	488	GLY	5.5
2	B	546	THR	5.3
1	C	396	ASN	5.0
1	A	397	ASP	4.3
2	B	507	SER	4.2
1	A	396	ASN	4.1
2	B	452	VAL	4.1
1	C	489	THR	3.8
1	C	476	GLU	3.7
1	C	541	GLU	3.7
2	B	396	ASN	3.6
2	B	567	ILE	3.4
2	B	541	GLU	3.3
1	A	507	SER	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	564	ILE	3.1
1	C	397	ASP	3.0
1	C	452	VAL	3.0
2	B	394	LYS	3.0
1	C	428	CYS	2.9
2	B	540	GLN	2.8
2	B	451	THR	2.8
1	C	418	ASP	2.8
1	A	540	GLN	2.8
2	B	458	ILE	2.8
2	B	485	LEU	2.8
2	B	506[A]	LYS	2.7
1	C	456	HIS	2.6
1	C	544	ASP	2.6
1	C	477	TYR	2.6
2	B	547	PRO	2.5
1	C	449	SER	2.5
2	B	395	ASN	2.5
1	C	546	THR	2.5
2	B	539	THR	2.4
1	A	485	LEU	2.4
1	A	488	GLY	2.3
1	C	486	THR	2.3
2	B	397	ASP	2.3
2	B	508	HIS	2.3
1	A	398	LYS	2.3
1	A	467	LEU	2.3
1	C	567	ILE	2.3
1	C	408	SER	2.3
2	B	482	ASN	2.2
1	C	441	VAL	2.2
1	A	451	THR	2.2
2	B	443	GLY	2.2
1	A	453	GLN	2.2
2	B	511	THR	2.2
1	A	489	THR	2.1
1	A	508	HIS	2.1
1	C	495	VAL	2.1
1	C	399	LEU	2.1
2	B	489	THR	2.1
1	C	581	GLU	2.1
2	B	416[A]	GLU	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	564	ILE	2.1
1	C	413	LEU	2.1
1	A	477	TYR	2.1
2	B	456	HIS	2.1
1	A	581	GLU	2.1
1	C	472	PHE	2.0
1	C	448	ILE	2.0
1	C	414	ASN	2.0
2	B	565	ASN	2.0
1	C	451	THR	2.0
1	C	462	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	A	604	4/4	0.78	0.22	38,39,40,40	0
3	EDO	A	602	4/4	0.81	0.20	28,30,33,38	0
3	EDO	A	603	4/4	0.83	0.18	39,39,39,41	0
3	EDO	A	601	4/4	0.88	0.14	32,33,33,34	0
3	EDO	C	602	4/4	0.88	0.14	30,30,31,31	0
4	MG	C	601	1/1	0.98	0.15	40,40,40,40	0

6.5 Other polymers [i](#)

There are no such residues in this entry.