



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 07:47 AM UTC

PDB ID : 3HDB / pdb_00003hdb
Title : Crystal structure of AaHIV, A metalloproteinase from venom of Agkistrodon
Acutus
Authors : Zhu, Z.Q.; Niu, L.W.; Teng, M.K.
Deposited on : 2009-05-07
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7754 (2.34-2.30)
Clashscore	190562	8383 (2.34-2.30)
Ramachandran outliers	187476	8303 (2.34-2.30)
Sidechain outliers	187428	8303 (2.34-2.30)
RSRZ outliers	180081	7760 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	417	
2	L	3	
3	B	2	

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3621 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AAHIV.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	417	Total	C	N	O	S	0	3	0
			3268	2040	541	637	50			

- Molecule 2 is a protein called KNL.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	L	3	Total	C	N	O	0	0	0
			26	16	5	5			

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	B	2	Total	C	N	O	0	0	0
			28	16	2	10			


- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Zn	0	0
			1	1		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	7	Total	Ca	0	0
			7	7		

- Molecule 6 is NONAETHYLENE GLYCOL (CCD ID: 2PE) (formula: $C_{18}H_{38}O_{10}$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		
6	A	1	Total	C	O	0	0
			13	8	5		
6	A	1	Total	C	O	0	0
			13	8	5		
6	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Cl	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	241	Total	O	0	0
			241	241		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: AAHIV



• Molecule 2: KNL



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	111.19Å 122.28Å 94.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.27 – 2.31 38.27 – 2.31	Depositor EDS
% Data completeness (in resolution range)	93.2 (38.27-2.31) 93.4 (38.27-2.31)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.21 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.212 , 0.249 0.216 , 0.221	Depositor DCC
R_{free} test set	1349 reflections (4.69%)	wwPDB-VP
Wilson B-factor (Å ²)	37.9	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 37.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3621	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 2PE, CL, NAG, ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.53	0/3356	0.82	1/4535 (0.0%)
2	L	0.52	0/25	0.97	0/30
All	All	0.53	0/3381	0.82	1/4565 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	516	ALA	N-CA-C	6.12	118.73	111.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3268	0	3034	38	0
2	L	26	0	32	2	0
3	B	28	0	25	1	0
4	A	1	0	0	0	0
5	A	7	0	0	0	0
6	A	49	0	64	1	0
7	A	1	0	0	0	0
8	A	241	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	3621	0	3155	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:232:ASN:HD22	1:A:242:VAL:H	1.35	0.75
1:A:487:ASN:HD21	1:A:500:ASN:H	1.34	0.72
1:A:330:ILE:HG23	2:L:3:LEU:HD23	1.79	0.65
1:A:487:ASN:ND2	1:A:500:ASN:H	1.95	0.64
1:A:372:CYS:SG	3:B:1:NAG:H82	2.38	0.63
1:A:309:HIS:HD2	1:A:311:LYS:H	1.48	0.62
1:A:245:VAL:O	1:A:281:LYS:HE3	2.01	0.61
1:A:285:ASN:HB2	1:A:307:MET:HE1	1.84	0.60
1:A:270:ALA:HB1	1:A:311:LYS:HG2	1.84	0.57
1:A:515:GLY:O	1:A:518:ILE:HG22	2.04	0.56
1:A:309:HIS:CD2	1:A:311:LYS:H	2.23	0.56
1:A:306:SER:HB2	1:A:312:ARG:HB3	1.87	0.55
1:A:514:PHE:CB	1:A:518:ILE:HD13	2.38	0.54
1:A:307:MET:HE3	1:A:313:SER:OG	2.07	0.53
1:A:514:PHE:HB2	1:A:518:ILE:HD13	1.89	0.53
1:A:529:ASN:HB3	1:A:548:CYS:SG	2.51	0.51
1:A:307:MET:HE3	1:A:308:CYS:H	1.75	0.50
1:A:425:ASN:C	1:A:425:ASN:HD22	2.18	0.49
1:A:448:GLU:HB2	1:A:453:ARG:HE	1.77	0.49
1:A:523:ASP:OD2	1:A:554:LYS:NZ	2.33	0.48
1:A:274:LYS:HD2	1:A:311:LYS:HB2	1.95	0.47
1:A:271:GLU:OE2	1:A:311:LYS:NZ	2.45	0.47
1:A:262:VAL:HG21	1:A:298:ILE:HD13	1.96	0.47
1:A:320:TYR:CE2	2:L:1:LYS:HG3	2.50	0.47
1:A:518:ILE:O	1:A:518:ILE:HG23	2.15	0.46
1:A:341:ILE:HD11	1:A:377:CYS:HB2	1.96	0.46
1:A:414:GLU:HB3	1:A:435:LEU:HG	1.98	0.45
1:A:284:ASP:HB3	1:A:307:MET:HE2	1.99	0.45
1:A:582:LYS:HD3	6:A:650:2PE:H21	1.98	0.44
1:A:514:PHE:HB2	1:A:518:ILE:HG21	1.99	0.44
1:A:309:HIS:CD2	1:A:312:ARG:H	2.35	0.43
1:A:227:ALA:HA	1:A:324:ASN:HD21	1.83	0.43
1:A:304:ILE:HA	1:A:337:HIS:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:243:VAL:CG1	1:A:398:ILE:HG12	2.50	0.42
1:A:206:ASP:O	1:A:209:MET:HB2	2.20	0.41
1:A:309:HIS:HD2	1:A:312:ARG:H	1.67	0.41
1:A:306:SER:O	1:A:307:MET:C	2.63	0.41
1:A:392:GLU:HA	1:A:393:PRO:HD3	1.94	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	418/417 (100%)	407 (97%)	11 (3%)	0	100	100
2	L	1/3 (33%)	1 (100%)	0	0	100	100
All	All	419/420 (100%)	408 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/361 (101%)	355 (98%)	9 (2%)	42	59
2	L	3/3 (100%)	2 (67%)	1 (33%)	0	0
All	All	367/364 (101%)	357 (97%)	10 (3%)	39	56

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	214	ASN
1	A	247	LEU
1	A	339	LEU
1	A	403	LEU
1	A	437	SER
1	A	455	SER
1	A	492	LEU
1	A	538	CYS
1	A	577	ASN
2	L	3	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	230	ASN
1	A	232	ASN
1	A	267	ASN
1	A	309	HIS
1	A	324	ASN
1	A	385	ASN
1	A	425	ASN
1	A	487	ASN
1	A	500	ASN
1	A	542	ASN
1	A	568	ASN
1	A	577	ASN
1	A	601	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	NAG	B	1	1,3	14,14,15	1.01	0	17,19,21	1.49	3 (17%)
3	NAG	B	2	3	14,14,15	1.02	0	17,19,21	2.85	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	B	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	B	2	3	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2	NAG	C1-O5-C5	7.16	121.79	112.19
3	B	2	NAG	C4-C3-C2	-6.56	101.41	111.02
3	B	2	NAG	C3-C4-C5	-5.24	100.73	110.23
3	B	1	NAG	C1-O5-C5	3.43	116.78	112.19
3	B	1	NAG	O3-C3-C4	-2.56	104.34	110.38
3	B	2	NAG	C2-N2-C7	-2.41	119.67	122.90
3	B	2	NAG	O5-C1-C2	2.22	114.73	111.29
3	B	1	NAG	C4-C3-C2	2.07	114.05	111.02

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	B	1	NAG	O5-C5-C6-O6
3	B	2	NAG	C8-C7-N2-C2
3	B	2	NAG	O7-C7-N2-C2
3	B	1	NAG	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 9 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	2PE	A	653	-	9,9,27	0.47	0	8,8,26	0.24	0
6	2PE	A	650	-	12,12,27	0.52	0	11,11,26	0.18	0
6	2PE	A	652	-	12,12,27	0.47	0	11,11,26	0.28	0
6	2PE	A	651	-	12,12,27	0.48	0	11,11,26	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	2PE	A	653	-	-	4/7/7/25	-
6	2PE	A	650	-	-	2/10/10/25	-
6	2PE	A	652	-	-	4/10/10/25	-
6	2PE	A	651	-	-	6/10/10/25	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (16) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	652	2PE	O4-C5-C6-O7
6	A	651	2PE	O4-C5-C6-O7
6	A	650	2PE	O1-C2-C3-O4
6	A	651	2PE	O1-C2-C3-O4
6	A	653	2PE	C5-C6-O7-C8
6	A	650	2PE	C5-C6-O7-C8
6	A	653	2PE	C6-C5-O4-C3
6	A	652	2PE	C6-C5-O4-C3
6	A	651	2PE	C8-C9-O10-C11
6	A	651	2PE	C2-C3-O4-C5
6	A	651	2PE	C5-C6-O7-C8
6	A	652	2PE	C12-C11-O10-C9
6	A	653	2PE	C2-C3-O4-C5
6	A	652	2PE	C9-C8-O7-C6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
6	A	653	2PE	O4-C5-C6-O7
6	A	651	2PE	O7-C8-C9-O10


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	650	2PE	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	417/417 (100%)	0.42	23 (5%) 30 33	14, 38, 58, 67	4 (0%)
2	L	3/3 (100%)	1.80	1 (33%) 1 1	60, 60, 60, 60	0
All	All	420/420 (100%)	0.43	24 (5%) 29 32	14, 38, 59, 67	4 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	193	TYR	4.5
1	A	342[A]	HIS	4.2
1	A	518	ILE	3.4
1	A	516	ALA	3.3
1	A	358	GLY	3.2
1	A	449	GLN	3.2
1	A	361	ILE	3.1
1	A	368	PHE	3.0
1	A	353	TYR	3.0
1	A	351	GLY	2.8
1	A	354	PRO	2.7
1	A	320	TYR	2.7
1	A	551[A]	GLU	2.4
2	L	3	LEU	2.4
1	A	438	GLY	2.3
1	A	362	SER	2.3
1	A	373	SER	2.3
1	A	517	ASP	2.3
1	A	359	PRO	2.3
1	A	377	CYS	2.2
1	A	382	MET	2.2
1	A	356	ILE	2.2
1	A	349	TYR	2.1
1	A	369	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAG	B	1	14/15	-	-	64,66,67,69	0
3	NAG	B	2	14/15	-	-	70,71,73,73	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	2PE	A	650	13/28	0.77	0.17	47,48,49,49	0
6	2PE	A	651	13/28	0.82	0.18	57,58,61,61	0
6	2PE	A	653	10/28	0.82	0.15	41,42,42,43	0
5	CA	A	655	1/1	0.84	0.12	79,79,79,79	0
6	2PE	A	652	13/28	0.86	0.20	79,82,84,84	0
7	CL	A	658	1/1	0.87	0.16	68,68,68,68	0
5	CA	A	657	1/1	0.89	0.10	73,73,73,73	0
5	CA	A	654	1/1	0.90	0.10	78,78,78,78	0
5	CA	A	656	1/1	0.95	0.24	49,49,49,49	0
5	CA	A	623	1/1	0.97	0.14	16,16,16,16	0
4	ZN	A	620	1/1	0.97	0.07	24,24,24,24	0
5	CA	A	622	1/1	0.98	0.04	31,31,31,31	0
5	CA	A	621	1/1	0.99	0.03	38,38,38,38	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 2PE A 650:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 2PE A 651:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



**Electron density around 2PE A 653:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 2PE A 652:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.