



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 11:34 AM UTC

PDB ID : 1HEW / pdb_00001hew
Title : REFINEMENT OF AN ENZYME COMPLEX WITH INHIBITOR BOUND AT PARTIAL OCCUPANCY. HEN EGG-WHITE LYSOZYME AND TRI-N-ACETYLCHITOTRIOSE AT 1.75 ANGSTROMS RESOLUTION
Authors : Cheetham, J.C.; Artymiuk, P.J.; Phillips, D.C.
Deposited on : 1992-01-20
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	129	 71% 21% 6% •
2	B	3	 100%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 1147 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HEN EGG WHITE LYSOZYME.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	129	Total	C	N	O	S	0	0	0
			1001	613	193	185	10			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	3	Total	C	N	O	0	0	0
			43	24	3	16			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	103	Total	O	0	0
			103	103		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: HEN EGG WHITE LYSOZYME

Chain A: 



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain B: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.86Å 78.86Å 38.25Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 1.75	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-1.75)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.229 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1147	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.33	5/1021 (0.5%)	1.83	28/1379 (2.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	71	GLY	C-N	13.01	1.51	1.33
1	A	71	GLY	C-O	9.13	1.36	1.23
1	A	72	SER	N-CA	8.55	1.56	1.45
1	A	5	ARG	CZ-NH1	5.23	1.40	1.32
1	A	63	TRP	CA-CB	5.09	1.61	1.53

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	71	GLY	O-C-N	14.92	142.10	122.70
1	A	125	ARG	CA-CB-CG	11.40	136.91	114.10
1	A	74	ASN	CA-CB-CG	9.11	121.71	112.60
1	A	47	THR	N-CA-CB	8.77	123.00	110.12
1	A	63	TRP	N-CA-C	8.12	123.49	113.50
1	A	69	THR	CB-CA-C	8.11	119.48	110.15
1	A	19	ASN	CA-CB-CG	-7.71	104.89	112.60
1	A	125	ARG	N-CA-CB	7.43	121.25	109.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	18	ASP	N-CA-C	-7.43	98.89	109.96
1	A	18	ASP	O-C-N	7.15	132.52	123.01
1	A	53	TYR	N-CA-C	6.71	120.45	109.24
1	A	66	ASP	CA-CB-CG	6.70	119.30	112.60
1	A	4	GLY	N-CA-C	-6.52	102.65	113.02
1	A	119	ASP	CA-CB-CG	-6.29	106.31	112.60
1	A	68	ARG	CA-CB-CG	6.18	126.46	114.10
1	A	103	ASN	CB-CA-C	5.97	120.81	110.72
1	A	92	VAL	CB-CA-C	-5.90	104.18	112.14
1	A	125	ARG	CB-CA-C	-5.82	101.78	110.16
1	A	47	THR	CA-CB-CG2	-5.68	100.84	110.50
1	A	62	TRP	N-CA-C	5.66	120.57	113.43
1	A	16	GLY	N-CA-C	5.41	123.41	115.08
1	A	61	ARG	NE-CZ-NH2	5.41	124.06	119.20
1	A	74	ASN	CB-CA-C	5.39	119.78	111.85
1	A	21	ARG	NE-CZ-NH2	5.33	123.99	119.20
1	A	73	ARG	N-CA-C	-5.26	101.58	109.95
1	A	113	ASN	N-CA-C	5.17	119.64	113.23
1	A	71	GLY	CA-C-O	-5.13	111.65	120.57
1	A	103	ASN	CA-CB-CG	5.04	117.64	112.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	128	ARG	Sidechain
1	A	45	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1001	0	959	8	3
2	B	43	0	39	0	0
3	A	103	0	0	0	2
All	All	1147	0	998	8	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:64:CYS:HA	1:A:74:ASN:ND2	2.03	0.74
1:A:64:CYS:HA	1:A:74:ASN:HD22	1.57	0.67
1:A:128:ARG:HH11	1:A:128:ARG:HB3	1.64	0.61
1:A:122:ALA:O	1:A:125:ARG:HG3	2.04	0.57
1:A:78:ILE:HG13	1:A:79:PRO:HD2	1.93	0.50
1:A:19:ASN:OD1	1:A:24:SER:HA	2.16	0.45
1:A:13:LYS:HD3	1:A:129:LEU:HD22	1.99	0.43
1:A:63:TRP:CE2	1:A:98:ILE:HG12	2.53	0.43

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:253:HOH:O	3:A:283:HOH:O[3_554]	1.92	0.28
1:A:13:LYS:O	1:A:128:ARG:NH1[8_555]	2.09	0.11
1:A:16:GLY:CA	1:A:128:ARG:NH2[8_555]	2.10	0.10
1:A:45:ARG:NH2	3:A:245:HOH:O[7_556]	2.14	0.06

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	127/129 (98%)	120 (94%)	6 (5%)	1 (1%)	16 5

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	70	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	105/105 (100%)	97 (92%)	8 (8%)	12 2

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	LEU
1	A	25	LEU
1	A	45	ARG
1	A	55	ILE
1	A	61	ARG
1	A	68	ARG
1	A	73	ARG
1	A	125	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	65	ASN
1	A	74	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	B	1	2	15,15,15	2.20	5 (33%)	21,21,21	1.81	5 (23%)
2	NAG	B	2	2	14,14,15	3.16	7 (50%)	17,19,21	2.23	3 (17%)
2	NAG	B	3	2	14,14,15	1.75	2 (14%)	17,19,21	1.36	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2	-	0/6/26/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	NAG	B	3	2	-	2/6/23/26	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	2	NAG	C1-C2	9.13	1.64	1.52
2	B	1	NAG	O4-C4	5.91	1.57	1.43
2	B	3	NAG	C2-N2	4.65	1.54	1.46
2	B	2	NAG	C4-C5	3.77	1.61	1.53
2	B	1	NAG	O7-C7	-3.70	1.15	1.23
2	B	2	NAG	O7-C7	-3.47	1.15	1.23
2	B	3	NAG	O7-C7	-3.41	1.15	1.23
2	B	2	NAG	O5-C1	2.71	1.48	1.43
2	B	2	NAG	O5-C5	2.67	1.48	1.43
2	B	1	NAG	O5-C1	2.32	1.48	1.42
2	B	2	NAG	O6-C6	2.19	1.51	1.42
2	B	1	NAG	C1-C2	2.08	1.55	1.52
2	B	2	NAG	C4-C3	2.04	1.57	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	NAG	O6-C6	2.00	1.50	1.42

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2	NAG	C1-O5-C5	-6.06	104.06	112.19
2	B	2	NAG	C4-C3-C2	-5.02	103.66	111.02
2	B	3	NAG	C1-C2-N2	-3.84	104.38	110.43
2	B	1	NAG	C3-C4-C5	3.78	117.08	110.23
2	B	1	NAG	C1-O5-C5	-3.60	106.68	113.65
2	B	1	NAG	O5-C1-C2	-3.56	105.94	109.52
2	B	2	NAG	O5-C5-C4	-2.63	104.42	110.83
2	B	3	NAG	C3-C4-C5	2.53	114.83	110.23
2	B	1	NAG	O4-C4-C3	-2.19	105.22	110.38
2	B	1	NAG	O1-C1-C2	-2.01	105.03	109.22

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	3	NAG	O5-C5-C6-O6
2	B	3	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.