



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:59 PM UTC

PDB ID : 4HEX / pdb\_00004hex  
Title : A novel conformation of calmodulin  
Authors : Kumar, V.; Chichili, V.P.R.; Sivaraman, J.  
Deposited on : 2012-10-04  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtrriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	156	 67% 17% 11%
1	B	156	 65% 19% 5% 10%

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 2289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Calmodulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	139	1098	673	177	241	7	0	0	0
1	B	140	1103	676	177	241	9	0	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-6	MET	-	expression tag	UNP P62204
A	-5	HIS	-	expression tag	UNP P62204
A	-4	HIS	-	expression tag	UNP P62204
A	-3	HIS	-	expression tag	UNP P62204
A	-2	HIS	-	expression tag	UNP P62204
A	-1	HIS	-	expression tag	UNP P62204
A	0	HIS	-	expression tag	UNP P62204
B	-6	MET	-	expression tag	UNP P62204
B	-5	HIS	-	expression tag	UNP P62204
B	-4	HIS	-	expression tag	UNP P62204
B	-3	HIS	-	expression tag	UNP P62204
B	-2	HIS	-	expression tag	UNP P62204
B	-1	HIS	-	expression tag	UNP P62204
B	0	HIS	-	expression tag	UNP P62204

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Zn	0	0
			3	3		
2	B	3	Total	Zn	0	0
			3	3		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Ca 2 2	0	0
3	B	2	Total Ca 2 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	41	Total O 41 41	0	0
4	B	37	Total O 37 37	0	0



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.83Å 116.40Å 38.80Å 90.00° 94.83° 90.00°	Depositor
Resolution (Å)	28.58 – 2.00 28.58 – 2.00	Depositor EDS
% Data completeness (in resolution range)	98.3 (28.58-2.00) 98.3 (28.58-2.00)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.76 (at 2.00Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.7.3_928), REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.210 , 0.237 (Not available) , 0.253	Depositor DCC
$R_{free}$ test set	1166 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.5	Xtriage
Anisotropy	0.355	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 51.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.478 for l,-k,h	Xtriage
Reported twinning fraction	0.526 for H, K, L 0.474 for L, -K, H	Depositor
Outliers	0 of 22820 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	2289	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	47.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.78% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.99	1/1110 (0.1%)	1.19	3/1491 (0.2%)
1	B	0.95	0/1115	1.13	0/1497
All	All	0.97	1/2225 (0.0%)	1.16	3/2988 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	106	LEU	C-O	-5.40	1.17	1.24

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	GLN	N-CA-C	7.97	127.78	110.80
1	A	64	ILE	CB-CA-C	-5.85	103.05	111.19
1	A	64	ILE	N-CA-C	5.24	115.13	107.37

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1098	0	1022	21	0
1	B	1103	0	1022	37	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	3	0	0	0	0
2	B	3	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
4	A	41	0	0	4	0
4	B	37	0	0	1	0
All	All	2289	0	2044	58	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:31:LYS:HD2	1:B:31:LYS:H	1.22	1.00
1:B:69:PHE:CZ	1:B:73:MET:HE2	2.14	0.83
1:A:77:MET:HE2	4:A:341:HOH:O	1.78	0.81
1:B:31:LYS:HD2	1:B:31:LYS:N	1.96	0.80
1:B:110:MET:HA	1:B:110:MET:HE3	1.65	0.79
1:B:49:LEU:HA	1:B:52:MET:HE3	1.65	0.79
1:B:103:ALA:HA	1:B:126:ILE:HD11	1.64	0.77
1:A:93:PHE:CZ	1:A:110:MET:HE2	2.21	0.76
1:B:111:THR:HG23	1:B:116:LYS:CD	2.18	0.74
1:B:31:LYS:H	1:B:31:LYS:CD	1.98	0.74
1:B:41:GLY:O	1:B:42:GLN:HB2	1.88	0.74
1:A:93:PHE:CE1	1:A:110:MET:HE2	2.26	0.70
1:B:88:GLU:OE2	4:B:318:HOH:O	2.12	0.67
1:A:33:LEU:HD21	1:A:52:MET:HE2	1.77	0.66
1:B:103:ALA:HB2	1:B:126:ILE:HD13	1.79	0.64
1:B:111:THR:HG23	1:B:116:LYS:HD2	1.80	0.63
1:A:27:THR:CG2	1:A:63:THR:HG23	2.31	0.60
1:A:27:THR:HG21	1:A:63:THR:CG2	2.31	0.60
1:B:23:ASP:OD1	1:B:25:ASP:HB2	2.02	0.59
1:B:110:MET:HA	1:B:110:MET:CE	2.32	0.59
1:B:146:MET:HA	1:B:146:MET:HE2	1.84	0.59
1:A:6:THR:HG21	1:A:11:ALA:HB2	1.83	0.59
1:A:87:ARG:HG2	1:A:139:TYR:CE1	2.38	0.58
1:A:121:GLU:O	1:A:125:MET:HG3	2.05	0.56
1:B:121:GLU:O	1:B:125:MET:HG3	2.07	0.55
1:B:111:THR:HG23	1:B:116:LYS:HD3	1.87	0.54
1:A:27:THR:CG2	1:A:63:THR:CG2	2.86	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:110:MET:HE2	1:B:115:GLU:HB2	1.90	0.54
1:A:59:ASP:OD2	4:A:328:HOH:O	2.19	0.53
1:A:77:MET:CE	4:A:341:HOH:O	2.47	0.53
1:A:77:MET:HE2	4:A:329:HOH:O	2.07	0.53
1:B:41:GLY:O	1:B:42:GLN:CB	2.56	0.52
1:B:94:ASP:HB2	1:B:101:ILE:HD13	1.91	0.52
1:B:103:ALA:HA	1:B:126:ILE:CD1	2.39	0.52
1:B:29:THR:HB	1:B:31:LYS:HD3	1.92	0.51
1:A:74:ALA:HB1	1:A:76:LYS:HE2	1.92	0.51
1:A:64:ILE:CD1	1:A:72:MET:HE2	2.41	0.50
1:B:94:ASP:HB2	1:B:101:ILE:CD1	2.41	0.50
1:B:13:PHE:CE1	1:B:73:MET:HB3	2.47	0.49
1:B:13:PHE:CZ	1:B:73:MET:HB3	2.50	0.46
1:B:49:LEU:HD23	1:B:52:MET:HE1	1.97	0.46
1:B:103:ALA:CB	1:B:126:ILE:HD13	2.47	0.45
1:B:118:THR:OG1	1:B:121:GLU:HG3	2.15	0.45
1:A:16:ALA:O	1:A:20:PHE:HD1	1.99	0.45
1:B:49:LEU:HD23	1:B:52:MET:CE	2.47	0.44
1:A:37:MET:SD	1:A:52:MET:HE1	2.58	0.44
1:B:33:LEU:O	1:B:37:MET:HG2	2.16	0.44
1:B:82:SER:O	1:B:86:ILE:HG13	2.18	0.44
1:A:80:THR:O	1:A:84:GLU:HG3	2.18	0.43
1:B:43:ASN:OD1	1:B:43:ASN:N	2.52	0.43
1:B:25:ASP:HB3	1:B:27:THR:H	1.84	0.42
1:B:69:PHE:HZ	1:B:73:MET:HE2	1.79	0.41
1:A:96:ASP:OD1	1:A:96:ASP:C	2.63	0.41
1:B:96:ASP:OD1	1:B:96:ASP:C	2.62	0.41
1:B:7:GLU:N	1:B:7:GLU:OE1	2.54	0.41
1:A:118:THR:HB	1:A:120:GLU:HG2	2.02	0.40
1:A:109:VAL:O	1:A:113:LEU:HG	2.21	0.40
1:B:127:ARG:NH1	1:B:127:ARG:HA	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/156 (88%)	130 (95%)	5 (4%)	2 (2%)	8	4
1	B	138/156 (88%)	131 (95%)	4 (3%)	3 (2%)	5	2
All	All	275/312 (88%)	261 (95%)	9 (3%)	5 (2%)	6	3

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	B	8	GLU
1	B	25	ASP
1	A	7	GLU
1	B	42	GLN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	119/134 (89%)	110 (92%)	9 (8%)	12	9
1	B	119/134 (89%)	106 (89%)	13 (11%)	6	3
All	All	238/268 (89%)	216 (91%)	22 (9%)	8	5

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	27	THR
1	A	38	ARG
1	A	63	THR
1	A	72	MET
1	A	120	GLU
1	A	125	MET
1	A	126	ILE

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	127	ARG
1	B	8	GLU
1	B	15	GLU
1	B	25	ASP
1	B	27	THR
1	B	31	LYS
1	B	38	ARG
1	B	43	ASN
1	B	50	GLN
1	B	51	ASP
1	B	55	GLU
1	B	86	ILE
1	B	110	MET
1	B	127	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 10 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	139/156 (89%)	-0.86	0 100 100	32, 44, 70, 88	0
1	B	140/156 (89%)	-0.85	0 100 100	31, 46, 67, 84	0
All	All	279/312 (89%)	-0.85	0 100 100	31, 45, 70, 88	0

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	ZN	A	201	1/1	0.99	0.07	53,53,53,53	0
3	CA	A	202	1/1	0.99	0.02	42,42,42,42	0
2	ZN	A	205	1/1	1.00	0.02	29,29,29,29	0
2	ZN	B	201	1/1	1.00	0.05	59,59,59,59	0
2	ZN	B	204	1/1	1.00	0.04	67,67,67,67	0
2	ZN	B	205	1/1	1.00	0.03	30,30,30,30	0
2	ZN	A	203	1/1	1.00	0.05	47,47,47,47	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CA	A	204	1/1	1.00	0.02	45,45,45,45	0
3	CA	B	202	1/1	1.00	0.05	33,33,33,33	0
3	CA	B	203	1/1	1.00	0.01	35,35,35,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.