



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 03:22 PM UTC

PDB ID : 6HGT / pdb_00006hgt
Title : Crystal structure of human KDM4A complexed with co-substrate analog NOG and histone H3 peptide with K9R mutation
Authors : Maw, J.; Le Bihan, Y.V.; van Montfort, R.L.M.
Deposited on : 2018-08-23
Resolution : 2.33 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.33 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	360	
1	B	360	
1	C	360	
1	D	360	
2	E	15	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	15	 20% 20% 60%
2	G	15	 40% 7% 53%
2	H	15	 27% 13% 60%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 11544 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysine-specific demethylase 4A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	350	Total	C	N	O	S	0	2	0
			2758	1796	451	495	16			
1	B	345	Total	C	N	O	S	0	1	0
			2734	1782	445	492	15			
1	C	348	Total	C	N	O	S	0	0	0
			2705	1761	445	484	15			
1	D	342	Total	C	N	O	S	0	2	0
			2663	1728	433	487	15			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP O75164
B	0	SER	-	expression tag	UNP O75164
C	0	SER	-	expression tag	UNP O75164
D	0	SER	-	expression tag	UNP O75164

- Molecule 2 is a protein called Histone H3.3.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	7	Total	C	N	O	0	0	0
			47	25	13	9			
2	F	6	Total	C	N	O	0	0	0
			44	24	12	8			
2	G	7	Total	C	N	O	0	0	1
			42	22	12	8			
2	H	6	Total	C	N	O	0	0	1
			38	21	11	6			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	9	ARG	LYS	engineered mutation	UNP P84243
F	9	ARG	LYS	engineered mutation	UNP P84243
G	9	ARG	LYS	engineered mutation	UNP P84243
H	9	ARG	LYS	engineered mutation	UNP P84243

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		
3	B	2	Total	Zn	0	0
			2	2		
3	C	2	Total	Zn	0	0
			2	2		
3	D	2	Total	Zn	0	0
			2	2		

- Molecule 4 is N-OXALYLGLYCINE (CCD ID: OGA) (formula: C₄H₅NO₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			10	4	1	5		
4	B	1	Total	C	N	O	0	0
			10	4	1	5		
4	C	1	Total	C	N	O	0	0
			10	4	1	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	D	1	Total	C	N	O	0	0
			10	4	1	5		

- Molecule 5 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	120	Total	O	0	0
			120	120		
5	B	133	Total	O	0	0
			133	133		
5	C	105	Total	O	0	0
			105	105		
5	D	95	Total	O	0	0
			95	95		
5	E	1	Total	O	0	0
			1	1		
5	F	6	Total	O	0	0
			6	6		
5	G	3	Total	O	0	0
			3	3		
5	H	2	Total	O	0	0
			2	2		

Chain E:  33% 13% 53%



- Molecule 2: Histone H3.3

Chain F:  20% 20% 60%




- Molecule 2: Histone H3.3

Chain G:  40% 7% 53%



- Molecule 2: Histone H3.3

Chain H:  27% 13% 60%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.44Å 103.41Å 144.52Å 90.00° 99.63° 90.00°	Depositor
Resolution (Å)	43.16 – 2.33 43.16 – 2.33	Depositor EDS
% Data completeness (in resolution range)	99.7 (43.16-2.33) 99.7 (43.16-2.33)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.73 (at 2.34Å)	Xtriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.251 , 0.280 0.258 , 0.285	Depositor DCC
R_{free} test set	3503 reflections (4.85%)	wwPDB-VP
Wilson B-factor (Å ²)	40.1	Xtriage
Anisotropy	0.280	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.073 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11544	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OGA, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	0/2844	1.34	5/3862 (0.1%)
1	B	0.94	0/2820	1.32	4/3831 (0.1%)
1	C	0.91	0/2790	1.31	4/3794 (0.1%)
1	D	0.89	0/2747	1.33	1/3735 (0.0%)
2	E	0.86	0/46	1.46	0/58
2	F	0.88	0/43	1.40	0/55
2	G	0.80	0/41	1.26	0/52
2	H	0.87	0/37	1.28	0/46
All	All	0.91	0/11368	1.32	14/15433 (0.1%)

There are no bond length outliers.

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	185	PHE	CA-CB-CG	6.53	120.33	113.80
1	C	311	ASP	CA-C-N	6.34	130.76	121.31
1	C	311	ASP	C-N-CA	6.34	130.76	121.31
1	A	185	PHE	CA-CB-CG	6.32	120.12	113.80
1	B	185	PHE	CA-CB-CG	6.28	120.08	113.80
1	B	215	HIS	CA-C-N	6.02	126.66	119.98
1	B	215	HIS	C-N-CA	6.02	126.66	119.98
1	D	185	PHE	CA-CB-CG	6.00	119.80	113.80
1	A	193	ASP	CB-CA-C	5.79	119.96	111.73
1	B	60	ASP	CA-CB-CG	5.51	118.11	112.60
1	A	147	GLU	CA-C-N	5.13	129.41	122.07
1	A	147	GLU	C-N-CA	5.13	129.41	122.07
1	A	284	ASN	CA-CB-CG	5.09	117.69	112.60
1	C	284	ASN	CA-CB-CG	5.06	117.66	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2758	0	2539	11	0
1	B	2734	0	2538	15	0
1	C	2705	0	2461	11	0
1	D	2663	0	2374	16	0
2	E	47	0	44	1	0
2	F	44	0	45	3	0
2	G	42	0	37	1	0
2	H	38	0	37	2	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	C	2	0	0	0	0
3	D	2	0	0	0	0
4	A	10	0	3	0	0
4	B	10	0	3	0	0
4	C	10	0	3	0	0
4	D	10	0	3	0	0
5	A	120	0	0	0	0
5	B	133	0	0	4	0
5	C	105	0	0	1	0
5	D	95	0	0	0	0
5	E	1	0	0	0	0
5	F	6	0	0	0	0
5	G	3	0	0	0	0
5	H	2	0	0	0	0
All	All	11544	0	10087	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:322:ARG:CG	1:D:329:TYR:CE2	2.40	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:322:ARG:HG3	1:D:329:TYR:CE2	1.94	1.03
1:D:322:ARG:CG	1:D:329:TYR:HE2	1.78	0.93
1:D:322:ARG:HG2	1:D:329:TYR:CE2	2.05	0.91
1:B:214:GLU:HG2	5:B:569:HOH:O	1.71	0.89
1:D:322:ARG:HG2	1:D:329:TYR:HE2	1.45	0.78
1:A:15:MET:HE2	1:A:17:PHE:HZ	1.63	0.62
1:B:86:ASN:HD22	2:F:12:GLY:C	2.09	0.60
1:B:342:ASP:OD2	1:B:345:LEU:HD13	2.03	0.58
1:C:93:THR:OG1	1:C:96:GLU:HG3	2.04	0.58
1:C:150:ILE:HG23	1:C:289:THR:HG22	1.87	0.57
1:A:82:PHE:HB2	1:A:244:LEU:HB2	1.86	0.56
1:B:82:PHE:HB2	1:B:244:LEU:HB2	1.90	0.54
1:A:294:ARG:HH11	1:A:348:PRO:HB2	1.73	0.54
1:D:70:PRO:HG3	1:D:92:MET:HG2	1.91	0.53
1:C:81:LEU:HD11	1:C:226:PHE:CD1	2.44	0.53
1:A:15:MET:HE2	1:A:17:PHE:CZ	2.43	0.52
1:D:322:ARG:HG2	1:D:329:TYR:CD2	2.43	0.52
1:B:294:ARG:HH11	1:B:348:PRO:HB2	1.74	0.52
1:C:254:GLY:HA2	5:C:506:HOH:O	2.10	0.52
1:C:70:PRO:HG3	1:C:92:MET:HG2	1.92	0.50
1:A:70:PRO:HG3	1:A:92:MET:HG2	1.93	0.50
1:B:20:THR:OG1	1:B:23:GLU:HG3	2.12	0.50
1:A:46:LYS:HG3	1:A:268:MET:SD	2.51	0.50
1:B:70:PRO:HG3	1:B:92:MET:HG2	1.95	0.49
1:C:200:LEU:HD12	1:C:286:ALA:HB2	1.95	0.49
1:D:74:LEU:HD23	1:D:87:ILE:HD12	1.93	0.49
1:C:74:LEU:HD23	1:C:87:ILE:HD12	1.94	0.48
1:C:169:GLU:HB2	2:G:8:ARG:HB2	1.96	0.47
1:D:250:LEU:O	1:D:255:ILE:HG23	2.14	0.47
1:A:74:LEU:HD23	1:A:87:ILE:HD12	1.95	0.47
1:C:81:LEU:HD11	1:C:226:PHE:HD1	1.80	0.47
5:B:504:HOH:O	2:F:11:THR:HG22	2.14	0.46
1:D:86:ASN:HD22	2:H:12:GLY:C	2.24	0.46
1:B:295:ARG:HB2	1:B:347:THR:HA	1.97	0.46
1:B:307:SER:HA	5:B:605:HOH:O	2.16	0.45
1:B:179:GLY:O	1:B:283:PHE:HA	2.15	0.45
1:D:93:THR:OG1	1:D:96:GLU:HG3	2.18	0.44
1:B:214:GLU:CG	5:B:569:HOH:O	2.44	0.43
1:D:169:GLU:HB2	2:H:8:ARG:HB2	2.00	0.43
1:A:150:ILE:HG21	1:A:287:GLU:HG2	2.00	0.43
1:B:154:ARG:HA	1:B:158:ASP:OD2	2.19	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:310:LYS:HA	1:D:312:MET:N	2.33	0.43
1:A:179:GLY:O	1:A:283:PHE:HA	2.20	0.42
1:A:192:MET:HE2	1:A:315:ILE:HB	2.01	0.42
1:D:250:LEU:O	1:D:255:ILE:CG2	2.68	0.42
1:C:327:GLU:H	1:C:327:GLU:HG3	1.50	0.42
1:D:250:LEU:CA	1:D:255:ILE:HG23	2.50	0.42
1:C:179:GLY:O	1:C:283:PHE:HA	2.19	0.42
1:D:250:LEU:HD22	1:D:255:ILE:HG12	2.03	0.41
1:B:169:GLU:HB2	2:F:8:ARG:HB2	2.03	0.41
1:B:192:MET:HG3	1:B:315:ILE:HD12	2.03	0.41
1:B:20:THR:HG1	1:B:23:GLU:HG3	1.85	0.40
1:A:240:HIS:O	2:E:12:GLY:HA3	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/360 (97%)	343 (98%)	7 (2%)	0	100	100
1	B	342/360 (95%)	335 (98%)	7 (2%)	0	100	100
1	C	346/360 (96%)	339 (98%)	7 (2%)	0	100	100
1	D	340/360 (94%)	335 (98%)	5 (2%)	0	100	100
2	E	5/15 (33%)	5 (100%)	0	0	100	100
2	F	4/15 (27%)	4 (100%)	0	0	100	100
2	G	5/15 (33%)	5 (100%)	0	0	100	100
2	H	4/15 (27%)	4 (100%)	0	0	100	100
All	All	1396/1500 (93%)	1370 (98%)	26 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	265/316 (84%)	259 (98%)	6 (2%)	44	56
1	B	272/316 (86%)	259 (95%)	13 (5%)	23	29
1	C	258/316 (82%)	250 (97%)	8 (3%)	35	45
1	D	250/316 (79%)	246 (98%)	4 (2%)	55	67
2	E	4/11 (36%)	3 (75%)	1 (25%)	0	0
2	F	4/11 (36%)	4 (100%)	0	100	100
2	G	4/11 (36%)	4 (100%)	0	100	100
2	H	4/11 (36%)	4 (100%)	0	100	100
All	All	1061/1308 (81%)	1029 (97%)	32 (3%)	36	47

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	65	LEU
1	A	175	TYR
1	A	182	LYS
1	A	270	THR
1	A	289	THR
1	A	318	ASP
1	B	65	LEU
1	B	131	ILE
1	B	164	SER
1	B	175	TYR
1	B	182	LYS
1	B	207	SER
1	B	214	GLU
1	B	221	ARG
1	B	235	GLU
1	B	270	THR
1	B	289	THR
1	B	336	LYS
1	B	345	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	65	LEU
1	C	125	LEU
1	C	175	TYR
1	C	182	LYS
1	C	270	THR
1	C	289	THR
1	C	327	GLU
1	C	347	THR
1	D	150	ILE
1	D	182	LYS
1	D	255	ILE
1	D	289	THR
2	E	8	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (21) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN
1	A	37	GLN
1	A	72	GLN
1	A	84	GLN
1	A	137	ASN
1	A	198	ASN
1	B	26	ASN
1	B	72	GLN
1	B	84	GLN
1	B	86	ASN
1	B	137	ASN
1	B	232	GLN
1	C	26	ASN
1	C	84	GLN
1	C	124	ASN
1	C	137	ASN
1	C	198	ASN
1	C	262	GLN
1	D	26	ASN
1	D	72	GLN
1	D	137	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	OGA	C	403	3	9,9,9	1.88	4 (44%)	8,11,11	1.32	2 (25%)
4	OGA	B	403	3	9,9,9	1.96	4 (44%)	8,11,11	1.29	2 (25%)
4	OGA	D	403	3	9,9,9	1.84	4 (44%)	8,11,11	1.42	2 (25%)
4	OGA	A	403	3	9,9,9	1.38	2 (22%)	8,11,11	1.23	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	OGA	C	403	3	-	3/8/9/9	-
4	OGA	B	403	3	-	3/8/9/9	-
4	OGA	D	403	3	-	5/8/9/9	-
4	OGA	A	403	3	-	3/8/9/9	-

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	403	OGA	O1-C1	3.63	1.31	1.22
4	C	403	OGA	O1-C1	3.41	1.31	1.22
4	D	403	OGA	O1-C1	3.13	1.30	1.22
4	A	403	OGA	O3-C5	-3.12	1.20	1.30
4	B	403	OGA	O3-C5	-2.79	1.21	1.30
4	B	403	OGA	O2-C1	-2.78	1.23	1.30
4	A	403	OGA	O4-C5	2.72	1.31	1.22
4	D	403	OGA	O4-C5	2.69	1.30	1.22
4	C	403	OGA	O2-C1	-2.67	1.23	1.30
4	D	403	OGA	O3-C5	-2.64	1.22	1.30
4	C	403	OGA	O3-C5	-2.58	1.22	1.30
4	D	403	OGA	O2-C1	-2.53	1.23	1.30
4	C	403	OGA	O4-C5	2.51	1.30	1.22
4	B	403	OGA	O4-C5	2.39	1.29	1.22

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	403	OGA	O3-C5-C4	2.97	124.09	112.81
4	C	403	OGA	O3-C5-C4	2.78	123.40	112.81
4	B	403	OGA	O3-C5-C4	2.74	123.24	112.81
4	A	403	OGA	O3-C5-C4	2.59	122.67	112.81
4	D	403	OGA	O4-C5-C4	-2.52	112.21	122.66
4	C	403	OGA	O4-C5-C4	-2.43	112.58	122.66
4	B	403	OGA	O4-C5-C4	-2.34	112.95	122.66
4	A	403	OGA	O4-C5-C4	-2.14	113.76	122.66

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	403	OGA	C1-C2-N1-C4
4	B	403	OGA	C1-C2-N1-C4
4	C	403	OGA	C1-C2-N1-C4
4	D	403	OGA	C1-C2-N1-C4
4	A	403	OGA	O2'-C2-N1-C4
4	B	403	OGA	O2'-C2-N1-C4
4	C	403	OGA	O2'-C2-N1-C4
4	D	403	OGA	O2'-C2-N1-C4
4	D	403	OGA	N1-C4-C5-O4
4	D	403	OGA	N1-C4-C5-O3

Continued on next page...


Continued from previous page...



Mol	Chain	Res	Type	Atoms
4	B	403	OGA	C5-C4-N1-C2
4	C	403	OGA	C5-C4-N1-C2
4	D	403	OGA	C5-C4-N1-C2
4	A	403	OGA	C5-C4-N1-C2


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	350/360 (97%)	-1.28	0 100 100	22, 42, 74, 102	2 (0%)
1	B	345/360 (95%)	-1.36	0 100 100	18, 37, 58, 69	1 (0%)
1	C	348/360 (96%)	-1.30	0 100 100	25, 44, 80, 94	0
1	D	342/360 (95%)	-1.25	0 100 100	21, 45, 74, 100	2 (0%)
2	E	7/15 (46%)	-1.29	0 100 100	48, 54, 61, 70	0
2	F	6/15 (40%)	-1.11	0 100 100	42, 51, 60, 64	0
2	G	7/15 (46%)	-1.20	0 100 100	50, 52, 55, 66	0
2	H	6/15 (40%)	-0.73	0 100 100	46, 50, 53, 64	0
All	All	1411/1500 (94%)	-1.29	0 100 100	18, 42, 72, 102	5 (0%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	OGA	A	403	10/10	0.99	0.04	42,47,49,49	0
4	OGA	B	403	10/10	0.99	0.03	18,25,29,29	10
4	OGA	C	403	10/10	0.99	0.04	25,30,32,32	10
4	OGA	D	403	10/10	0.99	0.04	33,37,42,42	10
3	ZN	C	401	1/1	1.00	0.01	42,42,42,42	0
3	ZN	C	402	1/1	1.00	0.01	50,50,50,50	1
3	ZN	D	401	1/1	1.00	0.01	40,40,40,40	0
3	ZN	D	402	1/1	1.00	0.01	39,39,39,39	1
3	ZN	A	401	1/1	1.00	0.01	39,39,39,39	0
3	ZN	A	402	1/1	1.00	0.01	49,49,49,49	0
3	ZN	B	401	1/1	1.00	0.01	36,36,36,36	0
3	ZN	B	402	1/1	1.00	0.01	34,34,34,34	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around OGA A 403:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around OGA B 403:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around OGA C 403:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.