



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 03:31 AM UTC

PDB ID : 1HHC / pdb_00001hhc
Title : Crystal structure of Decaplanin - space group P21, second form
Authors : Lehmann, C.; Vertessy, L.; Sheldrick, G.M.; Dauter, Z.; Dauter, M.
Deposited on : 2000-12-22
Resolution : 1.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	2021 (1.14-1.10)
Ramachandran outliers	187476	1978 (1.14-1.10)
Sidechain outliers	187428	1974 (1.14-1.10)
RSRZ outliers	180081	1984 (1.14-1.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	7	
1	B	7	
1	C	7	
1	D	7	
2	E	2	
2	F	2	
2	G	2	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	H	2	 100%

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DECAPLANIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	7	Total	C	Cl	N	O	0	0	0
			79	53	1	8	17			
1	B	7	Total	C	Cl	N	O	0	0	0
			76	50	1	8	17			
1	C	7	Total	C	Cl	N	O	0	0	0
			79	53	1	8	17			
1	D	7	Total	C	Cl	N	O	0	0	0
			79	53	1	8	17			

- Molecule 2 is an oligosaccharide called alpha-L-rhamnopyranose-(1-2)-beta-D-glucopyranoside.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	E	2	Total	C	O	0	0	0
			21	12	9			
2	F	2	Total	C	O	0	0	0
			21	12	9			
2	G	2	Total	C	O	0	0	0
			21	12	9			
2	H	2	Total	C	O	0	0	0
			21	12	9			

- Molecule 3 is 4-epi-vancosamine (CCD ID: ERE) (formula: C₇H₁₅NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			10	7	1	2		
3	B	1	Total	C	N	O	0	0
			10	7	1	2		
3	C	1	Total	C	N	O	0	0
			10	7	1	2		
3	D	1	Total	C	N	O	0	0
			10	7	1	2		

- Molecule 4 is CITRIC ACID (CCD ID: CIT) (formula: $C_6H_8O_7$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 13 6 7	0	0
4	C	1	Total C O 3 1 2	0	0
4	D	1	Total C O 13 6 7	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	41	Total O 41 41	0	0
5	B	42	Total O 42 42	0	0
5	C	38	Total O 38 38	0	0
5	D	42	Total O 42 42	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DECAPLANIN

Chain A: 





- Molecule 1: DECAPLANIN

Chain B: 





- Molecule 1: DECAPLANIN

Chain C: 




- Molecule 1: DECAPLANIN

Chain D: 




- Molecule 2: alpha-L-rhamnopyranose-(1-2)-beta-D-glucopyranose

Chain E: 




- Molecule 2: alpha-L-rhamnopyranose-(1-2)-beta-D-glucopyranose

Chain F: 




- Molecule 2: alpha-L-rhamnopyranose-(1-2)-beta-D-glucopyranose

Chain G:  100%

BGC1
RAM2

- Molecule 2: alpha-L-rhamnopyranose-(1-2)-beta-D-glucopyranose

Chain H:  100%

BGC1
RAM2

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	28.95Å 31.91Å 34.73Å 90.00° 109.87° 90.00°	Depositor
Resolution (Å)	18.08 – 1.13 18.08 – 1.13	Depositor EDS
% Data completeness (in resolution range)	96.8 (18.08-1.13) 89.8 (18.08-1.13)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.75 (at 1.13Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.224 , 0.264 0.205 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	5.1	Xtriage
Anisotropy	0.325	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 92.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	629	wwPDB-VP
Average B, all atoms (Å ²)	8.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 18.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OMX, OMZ, ERE, 3FG, MLU, RAM, GHP, CIT, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.77	0/7	1.19	0/8
1	B	1.27	0/7	4.44	2/8 (25.0%)
1	C	0.86	0/7	0.99	0/8
1	D	0.48	0/7	1.72	0/8
All	All	0.89	0/28	2.50	2/32 (6.2%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	3	ASN	OD1-CG-ND2	-8.85	113.75	122.60
1	B	3	ASN	CB-CG-ND2	6.63	126.34	116.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	79	0	45	1	0
1	B	76	0	38	0	0
1	C	79	0	45	0	0
1	D	79	0	46	0	0
2	E	21	0	18	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	F	21	0	18	0	0
2	G	21	0	18	0	0
2	H	21	0	18	0	0
3	A	10	0	13	0	0
3	B	10	0	13	0	0
3	C	10	0	13	0	0
3	D	10	0	13	0	0
4	A	13	0	5	1	0
4	C	3	0	0	0	0
4	D	13	0	5	0	0
5	A	41	0	0	0	0
5	B	42	0	0	0	0
5	C	38	0	0	0	0
5	D	42	0	0	0	0
All	All	629	0	308	1	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (1) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1:MLU:HD12	4:A:1002:CIT:C6	2.49	0.43

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1/7 (14%)	1 (100%)	0	0	100	100
1	B	1/7 (14%)	0	1 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	1/7 (14%)	0	1 (100%)	0	100	100
1	D	1/7 (14%)	1 (100%)	0	0	100	100
All	All	4/28 (14%)	2 (50%)	2 (50%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1/1 (100%)	1 (100%)	0	100	100
1	B	1/1 (100%)	0	1 (100%)	0	0
1	C	1/1 (100%)	1 (100%)	0	100	100
1	D	1/1 (100%)	1 (100%)	0	100	100
All	All	4/4 (100%)	3 (75%)	1 (25%)	0	0

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	3	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	3	ASN
1	B	3	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

24 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	GHP	A	4	2,1	10,11,12	1.33	1 (10%)	11,14,16	1.98	5 (45%)
1	GHP	C	4	2,1	10,11,12	1.20	0	11,14,16	1.02	0
1	3FG	A	7	1	12,13,13	1.29	1 (8%)	14,18,18	1.66	5 (35%)
1	MLU	C	1	1	7,8,9	0.74	0	7,9,11	1.40	2 (28%)
1	3FG	D	7	1	12,13,13	1.40	2 (16%)	14,18,18	1.91	3 (21%)
1	3FG	C	7	1	12,13,13	1.00	1 (8%)	14,18,18	2.01	5 (35%)
1	OMZ	A	2	1	12,14,15	1.10	0	17,19,21	2.37	4 (23%)
1	OMZ	C	2	1	12,14,15	1.24	1 (8%)	17,19,21	1.67	5 (29%)
1	GHP	B	4	2,1	10,11,12	1.39	1 (10%)	11,14,16	1.80	3 (27%)
1	OMX	B	6	3,1	11,13,14	0.97	0	15,17,19	2.51	6 (40%)
1	OMX	D	6	3,1	11,13,14	1.07	2 (18%)	15,17,19	2.66	8 (53%)
1	GHP	C	5	1	10,11,12	1.10	1 (10%)	11,14,16	2.47	4 (36%)
1	OMZ	B	2	1	12,14,15	0.76	0	17,19,21	1.00	0
1	GHP	B	5	1	10,11,12	1.46	1 (10%)	11,14,16	1.72	3 (27%)
1	MLU	B	1	1	4,5,9	1.36	1 (25%)	2,5,11	2.49	1 (50%)
1	GHP	D	4	2,1	10,11,12	1.31	1 (10%)	11,14,16	1.92	3 (27%)
1	3FG	B	7	1	12,13,13	1.19	0	14,18,18	1.38	3 (21%)
1	OMX	C	6	3,1	11,13,14	1.02	1 (9%)	15,17,19	1.28	2 (13%)
1	OMX	A	6	3,1	11,13,14	1.16	1 (9%)	15,17,19	1.76	5 (33%)
1	GHP	A	5	1	10,11,12	0.73	0	11,14,16	1.70	2 (18%)
1	MLU	A	1	1	7,8,9	1.16	1 (14%)	7,9,11	1.90	2 (28%)
1	GHP	D	5	1	10,11,12	1.01	0	11,14,16	2.13	3 (27%)
1	MLU	D	1	1	7,8,9	1.02	1 (14%)	7,9,11	1.87	2 (28%)
1	OMZ	D	2	1	12,14,15	1.43	2 (16%)	17,19,21	1.55	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	GHP	A	4	2,1	-	0/4/6/8	0/1/1/1
1	GHP	C	4	2,1	-	0/4/6/8	0/1/1/1
1	3FG	A	7	1	-	2/8/8/8	0/1/1/1
1	MLU	C	1	1	-	0/5/8/10	-
1	3FG	D	7	1	-	2/8/8/8	0/1/1/1
1	3FG	C	7	1	-	2/8/8/8	0/1/1/1
1	OMZ	A	2	1	-	1/9/10/12	0/1/1/1
1	OMZ	C	2	1	-	1/9/10/12	0/1/1/1
1	GHP	B	4	2,1	-	1/4/6/8	0/1/1/1
1	OMX	B	6	3,1	-	1/9/10/12	0/1/1/1
1	OMX	D	6	3,1	-	1/9/10/12	0/1/1/1
1	GHP	C	5	1	-	0/4/6/8	0/1/1/1
1	OMZ	B	2	1	-	1/9/10/12	0/1/1/1
1	GHP	B	5	1	-	0/4/6/8	0/1/1/1
1	MLU	B	1	1	-	0/2/4/10	-
1	GHP	D	4	2,1	-	2/4/6/8	0/1/1/1
1	3FG	B	7	1	-	2/8/8/8	0/1/1/1
1	OMX	C	6	3,1	-	1/9/10/12	0/1/1/1
1	OMX	A	6	3,1	-	1/9/10/12	0/1/1/1
1	GHP	A	5	1	-	0/4/6/8	0/1/1/1
1	MLU	A	1	1	-	0/5/8/10	-
1	GHP	D	5	1	-	0/4/6/8	0/1/1/1
1	MLU	D	1	1	-	0/5/8/10	-
1	OMZ	D	2	1	-	0/9/10/12	0/1/1/1

All (19) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	4	GHP	C6-C1	-3.72	1.33	1.39
1	B	5	GHP	C1-CA	3.38	1.56	1.52
1	D	4	GHP	C6-C1	-3.27	1.34	1.39
1	D	2	OMZ	CG-CB	-2.97	1.47	1.51
1	D	7	3FG	CB-CA	2.96	1.55	1.52
1	C	2	OMZ	CZ-CE1	-2.82	1.36	1.39
1	C	5	GHP	CA-C	2.69	1.55	1.51
1	C	6	OMX	CE2-CD2	-2.58	1.34	1.38
1	D	1	MLU	CA-N	2.44	1.51	1.47
1	A	6	OMX	CD2-CG	2.35	1.42	1.39
1	D	2	OMZ	OH-CZ	2.34	1.41	1.36
1	A	7	3FG	OXT-C	-2.28	1.23	1.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1	MLU	CA-N	2.27	1.51	1.47
1	D	7	3FG	CZ-CD2	-2.16	1.35	1.39
1	A	4	GHP	C1-CA	2.14	1.54	1.52
1	D	6	OMX	CD1-CG	-2.13	1.35	1.39
1	C	7	3FG	OXT-C	-2.12	1.23	1.30
1	A	1	MLU	CA-N	2.07	1.51	1.47
1	D	6	OMX	CD2-CG	2.05	1.42	1.39

All (79) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	2	OMZ	CD1-CE1-CZ	-7.79	116.38	120.92
1	D	6	OMX	CE1-CD1-CG	6.13	127.30	121.18
1	B	6	OMX	CD1-CE1-CZ	-5.99	113.54	119.88
1	D	5	GHP	C3-C2-C1	5.16	126.33	121.18
1	B	6	OMX	CE1-CD1-CG	5.02	126.19	121.18
1	D	6	OMX	CD1-CE1-CZ	-4.66	114.95	119.88
1	C	5	GHP	C5-C6-C1	-4.48	116.71	121.18
1	D	4	GHP	C3-C2-C1	-4.17	117.02	121.18
1	B	5	GHP	C2-C3-C4	-4.15	115.48	119.88
1	C	7	3FG	CD2-CZ-CD1	4.10	124.32	119.26
1	A	4	GHP	C5-C6-C1	4.06	125.23	121.18
1	A	6	OMX	CG-CB-CA	-4.03	106.28	111.58
1	D	2	OMZ	CG-CB-CA	4.01	116.85	111.58
1	D	7	3FG	CD2-CZ-CD1	3.91	124.09	119.26
1	D	1	MLU	CB-CA-N	-3.89	104.69	110.59
1	A	1	MLU	CB-CA-N	-3.89	104.69	110.59
1	C	5	GHP	C2-C3-C4	-3.86	115.79	119.88
1	C	5	GHP	C3-C2-C1	3.83	125.01	121.18
1	C	7	3FG	CG2-CD2-CZ	-3.72	115.38	120.47
1	C	2	OMZ	CD2-CE2-CZ	-3.57	116.93	120.50
1	A	2	OMZ	CD1-CE1-CL	3.57	124.29	118.45
1	D	6	OMX	CD1-CG-CB	3.40	125.83	120.71
1	B	1	MLU	O-C-CA	-3.33	113.55	123.94
1	D	7	3FG	CG2-CD2-CZ	-3.30	115.94	120.47
1	D	7	3FG	CD1-CG1-CB	-3.30	117.24	120.09
1	D	2	OMZ	O-C-CA	-3.27	116.36	124.77
1	A	5	GHP	C6-C5-C4	3.25	123.31	119.88
1	D	6	OMX	OC-CB-CA	-3.24	100.37	107.49
1	C	5	GHP	C6-C5-C4	3.23	123.30	119.88
1	D	5	GHP	C2-C3-C4	-3.21	116.48	119.88
1	B	4	GHP	C3-C2-C1	2.99	124.17	121.18

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	5	GHP	C3-C2-C1	2.95	124.13	121.18
1	B	4	GHP	C6-C1-C2	-2.94	114.66	118.30
1	A	2	OMZ	CE1-CD1-CG	2.90	124.32	120.28
1	A	6	OMX	OC-CB-CA	-2.82	101.28	107.49
1	C	2	OMZ	CD1-CE1-CZ	2.78	122.55	120.92
1	A	7	3FG	CD2-CG2-CB	2.78	122.48	120.09
1	C	1	MLU	CB-CA-N	-2.77	106.39	110.59
1	C	2	OMZ	O-C-CA	-2.77	117.66	124.77
1	B	6	OMX	OC-CB-CG	2.73	117.15	111.20
1	A	1	MLU	CB-CA-C	2.73	115.20	110.99
1	B	7	3FG	CD1-CG1-CB	-2.69	117.76	120.09
1	A	4	GHP	C6-C1-C2	-2.64	115.03	118.30
1	A	4	GHP	C2-C3-C4	2.62	122.65	119.88
1	D	4	GHP	C5-C6-C1	2.62	123.79	121.18
1	C	6	OMX	OC-CB-CA	-2.60	101.77	107.49
1	B	4	GHP	C5-C6-C1	2.58	123.75	121.18
1	A	6	OMX	OC-CB-CG	2.51	116.65	111.20
1	A	6	OMX	CD1-CG-CB	2.50	124.47	120.71
1	D	1	MLU	CB-CA-C	-2.49	107.16	110.99
1	C	6	OMX	CD2-CG-CB	-2.49	116.96	120.71
1	A	7	3FG	OXT-C-O	2.49	129.72	124.08
1	C	2	OMZ	CE2-CZ-CE1	2.47	120.82	118.49
1	D	4	GHP	C2-C3-C4	2.47	122.49	119.88
1	D	2	OMZ	CE2-CZ-CE1	2.44	120.79	118.49
1	A	2	OMZ	O-C-CA	-2.43	118.51	124.77
1	A	7	3FG	OD2-CD2-CG2	2.43	126.21	119.85
1	B	7	3FG	CG2-CB-CG1	2.36	121.31	118.09
1	B	5	GHP	C3-C2-C1	2.36	123.53	121.18
1	B	7	3FG	OXT-C-O	2.34	129.39	124.08
1	C	2	OMZ	CE1-CD1-CG	-2.34	117.03	120.28
1	D	6	OMX	CG-CB-CA	-2.28	108.59	111.58
1	A	7	3FG	OD2-CD2-CZ	-2.22	114.04	119.85
1	B	6	OMX	CD2-CG-CB	-2.21	117.39	120.71
1	A	4	GHP	C5-C4-C3	-2.18	116.29	119.77
1	C	7	3FG	CD2-CG2-CB	2.14	121.94	120.09
1	B	6	OMX	CE2-CD2-CG	-2.14	119.05	121.18
1	A	4	GHP	C6-C1-CA	2.14	123.96	120.64
1	B	5	GHP	C1-CA-C	-2.13	106.10	112.09
1	D	6	OMX	CD2-CG-CB	-2.12	117.51	120.71
1	C	7	3FG	CG2-CB-CG1	2.11	120.97	118.09
1	A	6	OMX	CD2-CG-CD1	-2.09	115.71	118.30
1	C	1	MLU	CB-CA-C	2.09	114.20	110.99

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	6	OMX	CD2-CG-CD1	-2.06	115.75	118.30
1	C	7	3FG	CB-CA-C	2.06	114.55	110.10
1	A	7	3FG	CZ-CD1-CG1	2.06	123.30	120.47
1	B	6	OMX	CG-CB-CA	-2.06	108.88	111.58
1	D	5	GHP	C6-C1-C2	-2.04	115.77	118.30
1	D	6	OMX	OC-CB-CG	2.00	115.56	111.20

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	6	OMX	O-C-CA-CB
1	B	6	OMX	O-C-CA-CB
1	D	6	OMX	O-C-CA-CB
1	A	7	3FG	C-CA-CB-CG1
1	A	7	3FG	C-CA-CB-CG2
1	B	7	3FG	C-CA-CB-CG1
1	B	7	3FG	C-CA-CB-CG2
1	C	7	3FG	C-CA-CB-CG2
1	D	7	3FG	C-CA-CB-CG1
1	D	7	3FG	C-CA-CB-CG2
1	B	4	GHP	C6-C1-CA-C
1	D	4	GHP	C2-C1-CA-C
1	D	4	GHP	C6-C1-CA-C
1	C	7	3FG	C-CA-CB-CG1
1	C	6	OMX	O-C-CA-CB
1	A	2	OMZ	O-C-CA-CB
1	B	2	OMZ	O-C-CA-CB
1	C	2	OMZ	O-C-CA-CB

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	1	MLU	1	0

5.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul

statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BGC	E	1	2,1	11,11,12	1.76	2 (18%)	15,15,17	1.35	2 (13%)
2	RAM	E	2	2	10,10,11	2.25	2 (20%)	14,14,16	2.06	4 (28%)
2	BGC	F	1	2,1	11,11,12	1.56	2 (18%)	15,15,17	1.81	4 (26%)
2	RAM	F	2	2	10,10,11	3.34	4 (40%)	14,14,16	1.51	2 (14%)
2	BGC	G	1	2,1	11,11,12	1.40	2 (18%)	15,15,17	1.10	1 (6%)
2	RAM	G	2	2	10,10,11	2.63	1 (10%)	14,14,16	1.40	1 (7%)
2	BGC	H	1	2,1	11,11,12	1.42	1 (9%)	15,15,17	1.32	2 (13%)
2	RAM	H	2	2	10,10,11	2.53	1 (10%)	14,14,16	1.82	5 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	E	1	2,1	-	0/2/19/22	0/1/1/1
2	RAM	E	2	2	-	-	0/1/1/1
2	BGC	F	1	2,1	-	0/2/19/22	0/1/1/1
2	RAM	F	2	2	-	-	0/1/1/1
2	BGC	G	1	2,1	-	0/2/19/22	0/1/1/1
2	RAM	G	2	2	-	-	0/1/1/1
2	BGC	H	1	2,1	-	0/2/19/22	0/1/1/1
2	RAM	H	2	2	-	-	0/1/1/1

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	2	RAM	C2-C3	-8.50	1.39	1.52
2	G	2	RAM	C2-C3	-7.68	1.40	1.52
2	H	2	RAM	C2-C3	-7.21	1.41	1.52
2	E	2	RAM	C2-C3	-5.49	1.44	1.52
2	F	2	RAM	O5-C1	4.57	1.51	1.43
2	E	1	BGC	O4-C4	3.48	1.51	1.43
2	F	2	RAM	O5-C5	-3.21	1.37	1.43

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	BGC	O3-C3	3.18	1.50	1.43
2	F	1	BGC	O5-C5	3.08	1.49	1.43
2	H	1	BGC	O4-C4	2.79	1.49	1.43
2	F	1	BGC	O4-C4	2.55	1.49	1.43
2	E	2	RAM	C4-C3	2.45	1.58	1.52
2	G	1	BGC	O5-C5	2.21	1.47	1.43
2	F	2	RAM	O2-C2	2.20	1.47	1.43
2	G	1	BGC	O4-C4	2.02	1.48	1.43

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	2	RAM	C1-C2-C3	4.44	116.11	109.64
2	E	2	RAM	O3-C3-C2	-3.74	102.42	110.05
2	F	2	RAM	C2-C3-C4	3.72	117.40	110.86
2	E	2	RAM	C3-C4-C5	-3.63	104.30	109.81
2	E	2	RAM	O5-C5-C4	3.46	115.77	109.55
2	F	1	BGC	O5-C5-C4	-3.39	102.58	110.83
2	E	2	RAM	O5-C5-C6	-3.38	100.07	107.40
2	H	2	RAM	O3-C3-C2	3.13	116.45	110.05
2	F	1	BGC	C2-C3-C4	-3.08	105.45	110.86
2	F	1	BGC	O2-C2-C3	-3.08	103.78	110.15
2	H	1	BGC	O2-C2-C1	2.97	116.03	109.22
2	F	1	BGC	C1-O5-C5	-2.90	108.31	112.19
2	F	2	RAM	O2-C2-C1	-2.80	102.81	109.22
2	G	1	BGC	O5-C5-C4	-2.59	104.53	110.83
2	E	1	BGC	O3-C3-C4	-2.45	104.59	110.38
2	H	1	BGC	C6-C5-C4	2.40	118.91	113.02
2	H	2	RAM	C6-C5-C4	-2.39	108.71	113.08
2	E	1	BGC	C6-C5-C4	2.29	118.64	113.02
2	H	2	RAM	O4-C4-C3	2.29	115.77	110.38
2	H	2	RAM	O5-C5-C6	-2.14	102.77	107.40
2	H	2	RAM	C3-C4-C5	-2.06	106.68	109.81



There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	CIT	C	1003	-	2,2,12	0.54	0	1,1,17	0.33	0
3	ERE	A	8	1	7,10,11	1.35	1 (14%)	6,15,17	1.55	1 (16%)
4	CIT	D	1001	-	12,12,12	1.32	2 (16%)	17,17,17	1.72	4 (23%)
3	ERE	B	8	1	7,10,11	1.17	0	6,15,17	1.02	1 (16%)
4	CIT	A	1002	-	12,12,12	1.57	3 (25%)	17,17,17	1.77	4 (23%)
3	ERE	D	8	1	7,10,11	1.91	2 (28%)	6,15,17	1.23	0
3	ERE	C	8	1	7,10,11	1.06	0	6,15,17	1.62	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	CIT	D	1001	-	-	2/16/16/16	-
3	ERE	A	8	1	-	-	0/1/1/1
3	ERE	B	8	1	-	-	0/1/1/1
4	CIT	A	1002	-	-	1/16/16/16	-
3	ERE	D	8	1	-	-	0/1/1/1
3	ERE	C	8	1	-	-	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	8	ERE	C2-C3	-3.96	1.49	1.54
4	A	1002	CIT	O4-C5	-2.86	1.21	1.30
3	A	8	ERE	C2-C3	-2.73	1.50	1.54
4	D	1001	CIT	O4-C5	-2.55	1.22	1.30
4	A	1002	CIT	C3-C6	-2.45	1.50	1.53
3	D	8	ERE	O4-C4	2.35	1.47	1.42

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1002	CIT	C4-C3	-2.33	1.51	1.54
4	D	1001	CIT	C2-C3	2.32	1.56	1.54

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1001	CIT	O6-C6-C3	3.50	119.86	113.14
4	A	1002	CIT	O4-C5-C4	3.39	125.08	114.35
3	A	8	ERE	C5A-C5-C4	-3.19	107.30	112.58
4	A	1002	CIT	O4-C5-O3	-2.94	115.76	123.33
4	D	1001	CIT	C2-C3-C6	-2.83	103.77	110.03
4	D	1001	CIT	O4-C5-C4	2.81	123.24	114.35
4	D	1001	CIT	O3-C5-C4	-2.77	115.09	122.95
4	A	1002	CIT	C2-C3-C6	-2.52	104.45	110.03
3	C	8	ERE	O5-C5-C5A	-2.35	102.29	107.40
3	C	8	ERE	O5-C1-C2	2.34	114.56	111.38
3	B	8	ERE	O5-C1-C2	2.34	114.55	111.38
4	A	1002	CIT	C4-C3-C6	2.28	115.08	110.03

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	1001	CIT	C3-C4-C5-O3
4	D	1001	CIT	C3-C4-C5-O4
4	A	1002	CIT	O7-C3-C4-C5

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1002	CIT	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	1/7 (14%)	0.05	0 100 100	6, 6, 6, 6	0
1	B	1/7 (14%)	0.08	0 100 100	6, 6, 6, 6	0
1	C	1/7 (14%)	-0.36	0 100 100	5, 5, 5, 5	0
1	D	1/7 (14%)	-0.11	0 100 100	6, 6, 6, 6	0
All	All	4/28 (14%)	-0.09	0 100 100	5, 6, 6, 6	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	MLU	B	1	6/10	0.83	0.12	9,10,11,12	0
1	MLU	A	1	9/10	0.93	0.08	6,7,9,11	0
1	MLU	D	1	9/10	0.93	0.09	6,7,8,9	0
1	MLU	C	1	9/10	0.94	0.07	4,5,7,8	0
1	OMX	A	6	13/14	0.94	0.07	5,5,6,6	0
1	OMX	B	6	13/14	0.95	0.06	2,3,4,4	0
1	GHP	D	4	11/12	0.95	0.06	3,3,4,4	0
1	GHP	B	5	11/12	0.95	0.06	2,3,4,4	0
1	GHP	C	5	11/12	0.95	0.06	2,3,4,6	0
1	3FG	A	7	13/13	0.95	0.06	7,8,10,11	0
1	GHP	B	4	11/12	0.96	0.06	3,4,4,5	0
1	GHP	C	4	11/12	0.96	0.05	2,2,3,3	0
1	GHP	D	5	11/12	0.96	0.05	2,3,3,6	0
1	GHP	A	4	11/12	0.96	0.06	3,4,4,4	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	3FG	C	7	13/13	0.96	0.05	4,5,7,8	0
1	OMZ	B	2	14/15	0.97	0.06	5,6,8,8	0
1	OMX	C	6	13/14	0.97	0.04	2,3,3,4	0
1	GHP	A	5	11/12	0.97	0.05	4,5,6,7	0
1	3FG	B	7	13/13	0.97	0.04	2,2,3,4	0
1	OMX	D	6	13/14	0.97	0.04	2,3,3,4	0
1	3FG	D	7	13/13	0.97	0.04	3,3,6,6	0
1	OMZ	A	2	14/15	0.98	0.05	4,5,5,6	0
1	OMZ	D	2	14/15	0.98	0.06	4,5,6,7	0
1	OMZ	C	2	14/15	0.99	0.04	3,4,5,6	0

6.3 Carbohydrates




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	RAM	E	2	10/11	0.91	0.10	7,9,12,13	0
2	BGC	F	1	11/12	0.92	0.07	5,7,8,9	0
2	RAM	G	2	10/11	0.93	0.07	5,6,9,9	0
2	RAM	H	2	10/11	0.93	0.07	6,7,9,9	0
2	BGC	H	1	11/12	0.94	0.07	5,5,7,7	0
2	RAM	F	2	10/11	0.94	0.06	6,7,9,10	0
2	BGC	G	1	11/12	0.95	0.06	3,4,4,6	0
2	BGC	E	1	11/12	0.95	0.06	4,5,6,6	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain E:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around Chain F:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around Chain G:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain H:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	CIT	A	1002	13/13	0.86	0.13	2,5,7,7	13
3	ERE	C	8	10/11	0.93	0.07	4,5,5,7	0
3	ERE	A	8	10/11	0.94	0.07	6,7,9,10	0
3	ERE	D	8	10/11	0.95	0.07	3,4,5,7	0
3	ERE	B	8	10/11	0.97	0.05	3,4,5,6	0
4	CIT	C	1003	3/13	0.97	0.05	8,8,8,8	0
4	CIT	D	1001	13/13	0.97	0.04	4,4,5,5	0

6.5 Other polymers [i](#)

There are no such residues in this entry.