



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:47 AM UTC

PDB ID : 3HHK / pdb_00003hhk
Title : HCV NS5b polymerase complex with a substituted benzothiadiazine
Authors : Concha, N.O.; Singh, O.
Deposited on : 2009-05-15
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5551 (1.70-1.70)
Clashscore	190562	5924 (1.70-1.70)
Ramachandran outliers	187476	5846 (1.70-1.70)
Sidechain outliers	187428	5846 (1.70-1.70)
RSRZ outliers	180081	5554 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	563	
1	B	563	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	77Z	A	564	X	-	-	-
2	77Z	B	564	X	-	-	-

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 9302 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HCV NS5 polymerase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	563	Total	C	N	O	S	26	5	0
			4383	2762	771	816	34			
1	B	563	Total	C	N	O	S	8	3	0
			4378	2760	771	814	33			

- Molecule 2 is 2-((3R)-3-[(3S)-1-(3-methylbutyl)-2,4-dioxo-1,2,3,4-tetrahydroquinolin-3-yl]-1,1-dioxido-3,4-dihydro-2H-1,2,4-benzothiadiazin-7-yl}oxy)acetamide (CCD ID: 77Z) (formula: C₂₃H₂₆N₄O₆S).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	228	Total 228	O 228	0	0
3	B	245	Total 245	O 245	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HCV NS5 polymerase



- Molecule 1: HCV NS5 polymerase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.74Å 107.44Å 126.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.92 – 1.70 81.92 – 1.70	Depositor EDS
% Data completeness (in resolution range)	87.5 (81.92-1.70) 87.5 (81.92-1.70)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.80 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.0	Depositor
R, R_{free}	0.229 , 0.250 0.228 , 0.231	Depositor DCC
R_{free} test set	5700 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	15.4	Xtriage
Anisotropy	0.424	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 34.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	9302	wwPDB-VP
Average B, all atoms (Å ²)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 45.67 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.2773e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 77Z

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.42	0/4503	0.77	5/6109 (0.1%)
1	B	0.43	1/4489 (0.0%)	0.78	3/6091 (0.0%)
All	All	0.43	1/8992 (0.0%)	0.77	8/12200 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	150	GLU	CB-CG	8.29	1.77	1.52

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	150	GLU	CB-CG-CD	13.69	135.88	112.60
1	B	151	LYS	CA-CB-CG	-11.47	91.16	114.10
1	B	150	GLU	CA-CB-CG	10.02	134.13	114.10
1	A	148	GLN	OE1-CD-NE2	-9.76	112.84	122.60
1	A	148	GLN	CB-CG-CD	9.45	128.66	112.60
1	A	152	GLY	N-CA-C	-7.37	105.16	115.32
1	A	148	GLN	CG-CD-NE2	6.51	126.16	116.40
1	A	151	LYS	N-CA-C	-5.02	106.74	112.92

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4383	0	4403	28	0
1	B	4378	0	4399	27	0
2	A	34	0	24	1	0
2	B	34	0	24	3	0
3	A	228	0	0	1	0
3	B	245	0	0	3	0
All	All	9302	0	8850	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:510:ARG:HH11	1:A:510:ARG:HG2	1.13	1.06
1:A:292[B]:THR:HG22	1:A:317:GLY:H	1.32	0.91
1:A:517:ARG:HG2	1:A:517:ARG:HH11	1.47	0.78
1:A:5:THR:HG23	1:A:278:ARG:HH12	1.50	0.75
1:B:5:THR:HG23	1:B:278:ARG:HH12	1.52	0.74
1:A:308:LEU:HB3	1:A:311:CYS:SG	2.29	0.72
1:A:510:ARG:HG2	1:A:510:ARG:NH1	1.91	0.66
1:B:291:ASN:HD22	2:B:564:77Z:HN54	1.43	0.65
1:B:119:ILE:CG2	1:B:173:MET:HE1	2.27	0.65
1:B:169:VAL:HG12	1:B:173:MET:HE2	1.80	0.64
1:A:291:ASN:HD22	2:A:564:77Z:HN54	1.49	0.61
1:A:52:VAL:HG12	1:A:223[B]:CYS:SG	2.42	0.59
1:A:292[B]:THR:CG2	1:A:317:GLY:H	2.11	0.57
1:B:170:CYS:HA	1:B:173:MET:HE3	1.88	0.56
1:A:144:VAL:HB	1:A:394:ARG:HG2	1.89	0.55
1:B:169:VAL:HG12	1:B:173:MET:CE	2.40	0.52
1:A:292[B]:THR:HG22	1:A:317:GLY:N	2.13	0.52
1:A:517:ARG:HG2	1:A:517:ARG:NH1	2.18	0.52
1:B:2:MET:N	3:B:740:HOH:O	2.42	0.51
1:B:144:VAL:HB	1:B:394:ARG:HG2	1.93	0.51
1:A:236:GLU:CD	1:A:280:ARG:HH22	2.19	0.51
1:A:510:ARG:NH1	1:A:510:ARG:CG	2.67	0.50
1:B:17:GLU:HB2	1:B:43:ARG:NH2	2.26	0.49
1:A:22:PRO:HG2	1:A:401:ARG:HG3	1.94	0.49
1:B:52:VAL:HG12	1:B:223[A]:CYS:SG	2.53	0.49
1:B:455:GLU:HB3	3:B:654:HOH:O	2.11	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:219:TYR:HB3	1:A:320:LEU:HD23	1.95	0.48
1:B:160:ILE:HA	1:B:282:SER:OG	2.15	0.47
1:B:22:PRO:HG2	1:B:401:ARG:HG3	1.97	0.47
1:B:562:HIS:O	1:B:563:SER:C	2.58	0.46
1:A:83:LEU:HB2	1:A:173:MET:HA	1.97	0.46
1:A:160:ILE:HA	1:A:282:SER:OG	2.16	0.46
1:A:258:GLU:HA	1:A:262:ILE:HD12	1.97	0.46
1:A:31:LEU:HD11	1:A:492:LEU:HD22	1.98	0.46
1:A:501:ARG:NH1	3:A:664:HOH:O	2.49	0.45
1:B:219:TYR:HB3	1:B:320:LEU:HD23	1.98	0.45
1:A:506:SER:O	1:A:510:ARG:HD3	2.16	0.44
1:A:182:LEU:HD12	1:A:243:CYS:SG	2.58	0.44
1:B:82:LEU:HD13	1:B:249:ALA:HB2	1.99	0.43
1:A:561:TYR:CE2	1:A:563:SER:HB2	2.54	0.43
1:A:288:SER:O	1:A:292[B]:THR:HG23	2.18	0.43
1:B:182:LEU:HD12	1:B:243:CYS:SG	2.58	0.43
1:B:17:GLU:HB2	1:B:43:ARG:HH22	1.83	0.42
1:B:348:ALA:N	1:B:349:PRO:HD3	2.33	0.42
1:B:414:MET:HE3	2:B:564:77Z:H44A	2.01	0.42
1:B:155:LYS:HA	1:B:156:PRO:HD3	1.94	0.42
1:B:405:VAL:HG13	1:B:445:CYS:HA	2.02	0.42
1:B:178:VAL:HG23	3:B:734:HOH:O	2.19	0.41
1:B:172:LYS:HE3	1:B:560:ILE:HD13	2.02	0.41
1:B:414:MET:HB2	2:B:564:77Z:H1	2.02	0.41
1:A:160:ILE:HD12	1:A:282:SER:OG	2.20	0.41
1:B:268:ASN:HB3	1:B:274:CYS:SG	2.62	0.40
1:A:268:ASN:HB3	1:A:274:CYS:SG	2.61	0.40
1:B:119:ILE:HG23	1:B:173:MET:HE1	2.03	0.40
1:A:236:GLU:OE2	1:A:280:ARG:NH2	2.42	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	566/563 (100%)	555 (98%)	10 (2%)	1 (0%)	43	28
1	B	564/563 (100%)	550 (98%)	12 (2%)	2 (0%)	30	16
All	All	1130/1126 (100%)	1105 (98%)	22 (2%)	3 (0%)	36	22

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	2	MET
1	A	536	LEU
1	B	536	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	484/479 (101%)	477 (99%)	7 (1%)	59	46
1	B	482/479 (101%)	475 (98%)	7 (2%)	57	43
All	All	966/958 (101%)	952 (99%)	14 (1%)	59	46

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	150	GLU
1	A	159	LEU
1	A	160	ILE
1	A	308	LEU
1	A	510	ARG
1	A	517	ARG
1	A	535	LYS
1	B	131[A]	VAL
1	B	131[B]	VAL
1	B	148	GLN
1	B	150	GLU
1	B	151	LYS
1	B	160	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	308	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	273	ASN
1	A	436	GLN
1	B	95	HIS
1	B	148	GLN
1	B	309	GLN
1	B	374	HIS
1	B	446	GLN
1	B	502	HIS
1	B	544	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	77Z	A	564	-	35,37,37	2.78	5 (14%)	45,55,55	1.77	12 (26%)
2	77Z	B	564	-	35,37,37	2.78	6 (17%)	45,55,55	1.80	12 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	77Z	A	564	-	2/2/8/10	0/10/49/49	0/4/4/4
2	77Z	B	564	-	2/2/8/10	0/10/49/49	0/4/4/4

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	564	77Z	O23-C10	11.79	1.37	1.22
2	A	564	77Z	O23-C10	11.76	1.37	1.22
2	B	564	77Z	C11-N12	-7.37	1.37	1.46
2	A	564	77Z	C11-N12	-7.34	1.37	1.46
2	B	564	77Z	C9-C10	-6.67	1.34	1.51
2	A	564	77Z	C9-C10	-6.65	1.34	1.51
2	A	564	77Z	C8-N7	2.18	1.38	1.36
2	B	564	77Z	C8-N7	2.14	1.38	1.36
2	A	564	77Z	C4-C10	-2.07	1.44	1.48
2	B	564	77Z	C4-C10	-2.05	1.44	1.48
2	B	564	77Z	C4-C3	-2.04	1.37	1.41

All (24) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	564	77Z	O23-C10-C4	-5.09	114.31	122.18
2	A	564	77Z	O23-C10-C4	-5.03	114.40	122.18
2	B	564	77Z	C50-O49-C19	-4.30	109.92	117.63
2	A	564	77Z	C50-O49-C19	-4.03	110.42	117.63
2	A	564	77Z	O21-S15-N16	3.97	111.95	108.00
2	A	564	77Z	O21-S15-C14	-3.96	104.35	109.30
2	B	564	77Z	O21-S15-C14	-3.89	104.45	109.30
2	B	564	77Z	O21-S15-N16	3.88	111.86	108.00
2	B	564	77Z	C4-C10-C9	3.39	121.39	116.18
2	A	564	77Z	C4-C10-C9	3.39	121.39	116.18
2	B	564	77Z	O22-S15-C14	-2.84	105.75	109.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	564	77Z	O22-S15-C14	-2.79	105.82	109.30
2	B	564	77Z	O22-S15-O21	2.79	121.27	118.44
2	A	564	77Z	O22-S15-O21	2.48	120.96	118.44
2	A	564	77Z	C13-C14-S15	-2.47	117.14	119.00
2	B	564	77Z	C20-C14-C13	2.31	123.16	120.85
2	B	564	77Z	C50-C51-N54	-2.28	111.28	117.60
2	B	564	77Z	O55-C51-C50	2.20	123.50	119.11
2	A	564	77Z	C20-C14-C13	2.19	123.04	120.85
2	B	564	77Z	C13-C14-S15	-2.19	117.36	119.00
2	A	564	77Z	C50-C51-N54	-2.15	111.63	117.60
2	B	564	77Z	O23-C10-C9	2.11	124.30	120.38
2	A	564	77Z	O55-C51-C50	2.07	123.24	119.11
2	A	564	77Z	O23-C10-C9	2.06	124.20	120.38

All (4) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	564	77Z	C9
2	A	564	77Z	C11
2	B	564	77Z	C9
2	B	564	77Z	C11


There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	564	77Z	1	0
2	B	564	77Z	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	560/563 (99%)	0.81	65 (11%)	9 9	6, 16, 26, 41	6 (1%)
1	B	563/563 (100%)	0.69	54 (9%)	13 14	10, 15, 25, 40	5 (0%)
All	All	1123/1126 (99%)	0.75	119 (10%)	11 11	6, 15, 26, 41	11 (0%)

All (119) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	542	ALA	10.6
1	B	543	SER	7.8
1	A	545	LEU	7.7
1	B	1	SER	7.6
1	A	542	ALA	7.5
1	A	540	PRO	7.1
1	B	545	LEU	7.0
1	B	563	SER	6.6
1	B	152	GLY	6.5
1	B	535	LYS	6.3
1	B	537	THR	6.0
1	A	539	ILE	6.0
1	B	95	HIS	5.8
1	A	535	LYS	5.8
1	A	541	ALA	5.8
1	B	539	ILE	5.6
1	A	537	THR	5.6
1	B	534	LEU	5.4
1	B	540	PRO	5.2
1	B	548	SER	5.2
1	A	14	CYS	5.1
1	A	543	SER	4.9
1	B	538	PRO	4.7
1	A	153	GLY	4.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	536	LEU	4.6
1	B	404	PRO	4.6
1	A	544	GLN	4.4
1	A	148	GLN	4.3
1	A	536	LEU	4.3
1	A	538	PRO	4.3
1	B	43	ARG	4.2
1	A	402	HIS	4.1
1	B	150	GLU	4.1
1	A	405	VAL	4.1
1	B	405	VAL	4.0
1	A	377	SER	4.0
1	A	16	ALA	3.9
1	A	95	HIS	3.9
1	B	402	HIS	3.8
1	A	311	CYS	3.8
1	B	14	CYS	3.8
1	A	100	LYS	3.7
1	B	541	ALA	3.7
1	A	534	LEU	3.7
1	B	149	PRO	3.6
1	A	548	SER	3.5
1	B	533	LYS	3.5
1	B	532	THR	3.5
1	A	353	PRO	3.4
1	B	544	GLN	3.4
1	A	531	LYS	3.3
1	A	404	PRO	3.3
1	B	110	ASN	3.3
1	A	113	SER	3.3
1	B	466	LEU	3.2
1	A	466	LEU	3.2
1	A	563	SER	3.2
1	A	160	ILE	3.1
1	B	148	GLN	3.1
1	B	546	ASP	3.1
1	A	376	ALA	3.1
1	B	490	ARG	3.0
1	A	532	THR	3.0
1	B	153	GLY	3.0
1	A	530	VAL	3.0
1	A	461	GLN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	513	SER	2.9
1	B	50	LYS	2.8
1	B	100	LYS	2.8
1	A	82	LEU	2.8
1	B	549	GLY	2.8
1	A	533	LYS	2.7
1	A	515	GLY	2.7
1	A	86	GLU	2.6
1	A	128	GLU	2.6
1	A	513	SER	2.6
1	B	94	PRO	2.6
1	B	106	LYS	2.5
1	B	151	LYS	2.5
1	A	465	ARG	2.5
1	A	505	ARG	2.5
1	B	16	ALA	2.5
1	A	546	ASP	2.4
1	A	512	LEU	2.4
1	A	110	ASN	2.4
1	B	355	GLN	2.4
1	A	528	TRP	2.4
1	B	330	GLN	2.3
1	B	401	ARG	2.3
1	A	403	THR	2.3
1	B	128	GLU	2.3
1	A	51	LYS	2.3
1	A	330	GLN	2.2
1	B	5	THR	2.2
1	B	337	ARG	2.2
1	B	469	LEU	2.2
1	A	106	LYS	2.2
1	A	149	PRO	2.2
1	A	550	TRP	2.2
1	A	15	ALA	2.2
1	A	97	ALA	2.2
1	A	352	ASP	2.2
1	A	469	LEU	2.2
1	A	154	ARG	2.2
1	A	81	LYS	2.2
1	A	361	GLU	2.1
1	A	355	GLN	2.1
1	B	403	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	377	SER	2.1
1	A	159	LEU	2.1
1	A	547	LEU	2.1
1	B	82	LEU	2.1
1	B	547	LEU	2.1
1	A	516	GLY	2.0
1	A	437	GLU	2.0
1	A	517	ARG	2.0
1	B	154	ARG	2.0
1	B	347	SER	2.0
1	B	530	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	77Z	A	564	34/34	0.90	0.10	13,14,16,16	0
2	77Z	B	564	34/34	0.91	0.10	13,14,17,17	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 77Z A 564:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 77Z B 564:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.