



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 01:55 AM UTC

PDB ID : 5HI3 / pdb_00005hi3
Title : Binding site elucidation and structure guided design of macrocyclic IL-17A antagonists
Authors : Liu, S.
Deposited on : 2016-01-11
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	
1	B	132	
2	C	237	
2	H	237	
3	D	214	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	L	214	2%86%11%
4	I	16	62%69%12%6%12%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8343 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-17A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	100	Total	C	N	O	S	0	0	0
			806	503	151	146	6			
1	B	88	Total	C	N	O	S	0	0	0
			708	442	132	129	5			

- Molecule 2 is a protein called CAT-2000 FAB heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	220	Total	C	N	O	S	0	0	0
			1622	1017	278	321	6			
2	H	220	Total	C	N	O	S	0	0	0
			1622	1017	278	321	6			


- Molecule 3 is a protein called CAT-2000 FAB light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	213	Total	C	N	O	S	0	1	0
			1618	1015	267	331	5			
3	L	213	Total	C	N	O	S	0	1	0
			1618	1015	267	331	5			

- Molecule 4 is a protein called synthetic IL-17A peptide inhibitor.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	I	14	Total	C	N	O	0	0	0
			120	81	19	20			

- Molecule 5 is N-(4-{2-[(1-[2-(dimethylamino)-2-oxoethyl]cyclopentyl)acetyl]amino}ethyl}phenyl)-2-fluoro-Nalpha-[(1-methyl-1H-pyrazol-5-yl)carbonyl]-L-phenylalaninamide (CCD ID: 63O) (formula: C₃₃H₄₁FN₆O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	B	1	Total	C	F	N	O	0	1
			88	66	2	12	8		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	O	0	0
			3	3		
6	B	7	Total	O	0	0
			7	7		
6	C	31	Total	O	0	0
			31	31		
6	D	22	Total	O	0	0
			22	22		
6	H	40	Total	O	0	0
			40	40		
6	L	38	Total	O	0	0
			38	38		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Interleukin-17A




• Molecule 1: Interleukin-17A




• Molecule 2: CAT-2000 FAB heavy chain





• Molecule 2: CAT-2000 FAB heavy chain








- Molecule 3: CAT-2000 FAB light chain



- Molecule 3: CAT-2000 FAB light chain



- Molecule 4: synthetic IL-17A peptide inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	91.11Å 68.32Å 100.45Å 90.00° 90.82° 90.00°	Depositor
Resolution (Å)	47.80 – 2.15 47.80 – 2.15	Depositor EDS
% Data completeness (in resolution range)	94.5 (47.80-2.15) 94.5 (47.80-2.15)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.24 (at 2.16Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.197 , 0.231 (Not available) , 0.229	Depositor DCC
R_{free} test set	3146 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	33.0	Xtriage
Anisotropy	0.093	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.025 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8343	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 63O

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/824	1.11	1/1120 (0.1%)
1	B	0.73	0/725	1.09	2/986 (0.2%)
2	C	0.91	2/1658 (0.1%)	1.12	3/2256 (0.1%)
2	H	0.87	1/1658 (0.1%)	1.14	2/2256 (0.1%)
3	D	0.81	0/1660	1.15	8/2269 (0.4%)
3	L	0.82	0/1660	1.15	7/2269 (0.3%)
4	I	0.86	0/125	1.40	0/174
All	All	0.83	3/8310 (0.0%)	1.14	23/11330 (0.2%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	82	MET	SD-CE	-7.00	1.62	1.79
2	H	82	MET	SD-CE	-6.78	1.62	1.79
2	C	189	LEU	CA-C	5.04	1.59	1.52

All (23) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	47	VAL	N-CA-C	-7.64	105.72	111.90
2	H	67	PHE	CA-CB-CG	7.54	121.34	113.80
2	C	67	PHE	CA-CB-CG	7.11	120.91	113.80
3	L	92	ASP	CA-CB-CG	7.05	119.65	112.60
3	D	47	VAL	N-CA-C	-6.97	106.25	111.90
3	D	92	ASP	CA-CB-CG	6.68	119.28	112.60
1	A	88	ASN	CA-CB-CG	6.08	118.68	112.60
3	D	182	THR	N-CA-C	-5.80	102.59	110.36
1	B	88	ASN	CA-CB-CG	5.76	118.36	112.60
3	L	164	THR	N-CA-C	-5.72	102.84	109.93
3	L	60	ASP	CA-C-N	5.71	128.50	120.28

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	L	60	ASP	C-N-CA	5.71	128.50	120.28
3	D	60	ASP	CA-C-N	5.67	128.45	120.28
3	D	60	ASP	C-N-CA	5.67	128.45	120.28
3	L	160	VAL	N-CA-CB	5.61	117.72	110.99
3	D	203	VAL	N-CA-CB	5.53	118.83	111.64
3	L	182	THR	N-CA-C	-5.50	103.00	110.36
2	C	151	THR	CB-CA-C	5.47	119.83	109.37
2	C	144	ASP	N-CA-C	5.36	117.72	111.02
1	B	86	HIS	N-CA-C	-5.26	106.84	113.20
3	D	160	VAL	N-CA-CB	5.19	117.22	110.99
2	H	151	THR	CB-CA-C	5.11	119.13	109.37
3	D	164	THR	N-CA-C	-5.02	103.70	109.93

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	806	0	775	3	0
1	B	708	0	680	7	0
2	C	1622	0	1601	14	0
2	H	1622	0	1601	16	0
3	D	1618	0	1555	11	0
3	L	1618	0	1555	14	0
4	I	120	0	115	3	0
5	B	88	0	0	1	0
6	A	3	0	0	0	0
6	B	7	0	0	0	0
6	C	31	0	0	1	0
6	D	22	0	0	0	0
6	H	40	0	0	0	0
6	L	38	0	0	2	0
All	All	8343	0	7882	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:L:117:THR:HB	6:L:330:HOH:O	1.42	1.16
2:C:82:MET:HE2	2:C:82(C):LEU:HD21	1.58	0.86
2:C:11:LEU:HB2	2:C:147:PRO:HG3	1.85	0.58
1:A:92:ILE:HD11	1:A:122:THR:HB	1.84	0.58
1:B:92:ILE:HD11	1:B:122:THR:HB	1.86	0.57
3:L:28:LEU:H	3:L:69:ASN:ND2	2.04	0.56
3:L:28:LEU:H	3:L:69:ASN:HD21	1.53	0.55
3:D:195:GLN:HG2	3:D:204:GLU:HG3	1.90	0.54
3:L:195:GLN:HG2	3:L:204:GLU:HG3	1.90	0.54
2:C:148:GLN:NE2	6:C:301:HOH:O	2.41	0.53
2:H:82:MET:HE2	2:H:82(C):LEU:HD21	1.90	0.53
2:H:150:VAL:HG23	2:H:178:LEU:HD21	1.91	0.53
2:H:181:VAL:HG11	3:L:136:LEU:CD1	2.39	0.52
3:D:47:VAL:HG12	3:D:48:ILE:HG12	1.93	0.51
3:L:47:VAL:HG12	3:L:48:ILE:HG12	1.92	0.51
1:B:112:LEU:HD12	4:I:5:ILE:HG22	1.91	0.51
2:C:151:THR:HG22	2:C:199:ASN:HB3	1.93	0.50
2:H:5:LEU:HD23	2:H:105:GLN:HE22	1.76	0.50
2:C:5:LEU:HD23	2:C:105:GLN:HE22	1.76	0.50
2:H:130:SER:HB2	2:H:137:ALA:HB3	1.94	0.50
2:C:145:TYR:CE2	2:C:150:VAL:HG13	2.47	0.49
3:D:143:GLY:O	3:D:165:PRO:HG2	2.12	0.49
1:B:62:TYR:HB3	1:B:99:LEU:HB3	1.95	0.48
1:A:87:MET:HB3	1:A:125:THR:HG22	1.94	0.48
2:H:181:VAL:CG1	3:L:136:LEU:CD1	2.91	0.48
1:B:87:MET:HB3	1:B:125:THR:HG22	1.95	0.47
3:L:78:LEU:HD21	3:L:106:VAL:HG22	1.95	0.47
1:A:62:TYR:HB3	1:A:99:LEU:HB3	1.96	0.47
2:H:151:THR:HG22	2:H:199:ASN:HB3	1.97	0.46
2:H:144:ASP:HB3	2:H:175:LEU:HD13	1.97	0.46
2:H:181:VAL:HG11	3:L:136:LEU:HD12	1.98	0.46
3:D:28:LEU:H	3:D:69:ASN:HD21	1.63	0.45
2:C:67:PHE:CZ	2:C:82:MET:HE3	2.51	0.44
3:D:36:TYR:CE1	3:D:46:ILE:HG22	2.52	0.44
1:B:86:HIS:HE1	6:L:319:HOH:O	2.00	0.44
1:B:63:PRO:HB3	5:B:400[A]:63O:C33	2.48	0.43
3:D:36:TYR:HE1	3:D:46:ILE:HG22	1.84	0.43
3:D:46:ILE:HD13	3:D:49:PHE:HB3	2.01	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:82:MET:HE2	2:C:82(C):LEU:CD2	2.40	0.42
3:L:4:LEU:HD11	3:L:90:THR:HG22	2.01	0.42
2:H:119:PRO:HB3	2:H:145:TYR:HB3	2.00	0.42
2:C:47:TRP:CG	3:D:96:VAL:HB	2.55	0.42
2:C:95:ASP:OD1	2:C:100(A):THR:HB	2.20	0.42
2:H:47:TRP:CG	3:L:96:VAL:HB	2.54	0.42
2:C:150:VAL:HG12	2:C:200:HIS:CD2	2.55	0.42
4:I:5:ILE:HB	4:I:9:LEU:HD23	2.02	0.42
2:H:34:MET:HB3	2:H:78:LEU:HD22	2.02	0.41
3:D:4:LEU:HD11	3:D:90:THR:HG22	2.01	0.41
2:H:95:ASP:OD1	2:H:100(A):THR:HB	2.20	0.41
3:D:28:LEU:H	3:D:69:ASN:ND2	2.18	0.41
2:C:142:VAL:HG11	2:C:150:VAL:HG21	2.02	0.41
2:C:167:PRO:HG2	3:D:166:SER:OG	2.21	0.41
2:H:148:GLN:HA	2:H:148:GLN:HE21	1.84	0.41
1:B:109:SER:HA	4:I:2:HIS:O	2.21	0.40
2:C:67:PHE:CE2	2:C:82:MET:HE3	2.56	0.40
3:L:46:ILE:HD13	3:L:49:PHE:HB3	2.02	0.40
3:L:36:TYR:CE1	3:L:46:ILE:HG22	2.56	0.40
2:H:167:PRO:HG2	3:L:166:SER:OG	2.21	0.40
2:H:29:PHE:CD2	2:H:76:ASN:HA	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	94/132 (71%)	91 (97%)	3 (3%)	0	100	100
1	B	84/132 (64%)	84 (100%)	0	0	100	100
2	C	218/237 (92%)	215 (99%)	3 (1%)	0	100	100
2	H	218/237 (92%)	214 (98%)	4 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	D	212/214 (99%)	205 (97%)	7 (3%)	0	100	100
3	L	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
4	I	12/16 (75%)	12 (100%)	0	0	100	100
All	All	1050/1182 (89%)	1028 (98%)	22 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	93/124 (75%)	87 (94%)	6 (6%)	15	11
1	B	81/124 (65%)	77 (95%)	4 (5%)	22	19
2	C	182/196 (93%)	171 (94%)	11 (6%)	17	13
2	H	182/196 (93%)	173 (95%)	9 (5%)	22	19
3	D	185/186 (100%)	176 (95%)	9 (5%)	22	19
3	L	185/186 (100%)	176 (95%)	9 (5%)	22	19
4	I	13/14 (93%)	12 (92%)	1 (8%)	12	7
All	All	921/1026 (90%)	872 (95%)	49 (5%)	20	17

All (49) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	SER
1	A	87	MET
1	A	94	GLN
1	A	98	VAL
1	A	112	LEU
1	A	115	ILE
1	B	40	SER
1	B	87	MET
1	B	112	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	115	ILE
2	C	7	SER
2	C	13	GLN
2	C	35	SER
2	C	135	THR
2	C	148	GLN
2	C	150	VAL
2	C	151	THR
2	C	159	LEU
2	C	178	LEU
2	C	189	LEU
2	C	206	LYS
3	D	19	VAL
3	D	46	ILE
3	D	78	LEU
3	D	103	LYS
3	D	156	VAL
3	D	157	LYS
3	D	160	VAL
3	D	167	LYS
3	D	203	VAL
2	H	7	SER
2	H	13	GLN
2	H	35	SER
2	H	128	SER
2	H	148	GLN
2	H	151	THR
2	H	159	LEU
2	H	189	LEU
2	H	206	LYS
4	I	5	ILE
3	L	19	VAL
3	L	28	LEU
3	L	46	ILE
3	L	81	GLU
3	L	103	LYS
3	L	156	VAL
3	L	157	LYS
3	L	160	VAL
3	L	203	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (31) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	25	ASN
1	A	27	ASN
1	A	29	HIS
1	A	36	ASN
1	A	86	HIS
1	B	36	ASN
1	B	73	HIS
1	B	86	HIS
2	C	39	GLN
2	C	82(A)	ASN
2	C	105	GLN
2	C	148	GLN
2	C	171	GLN
2	C	192	GLN
3	D	38	GLN
3	D	53	GLN
3	D	69	ASN
3	D	89	GLN
3	D	127	GLN
3	D	195	GLN
2	H	82(A)	ASN
2	H	105	GLN
2	H	148	GLN
2	H	192	GLN
2	H	199	ASN
3	L	53	GLN
3	L	69	ASN
3	L	89	GLN
3	L	109	GLN
3	L	127	GLN
3	L	195	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	63O	B	400[A]	-	47,47,47	0.52	1 (2%)	61,65,65	0.72	2 (3%)
5	63O	B	400[B]	-	47,47,47	0.46	1 (2%)	61,65,65	0.62	1 (1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	63O	B	400[A]	-	-	3/40/49/49	0/4/4/4
5	63O	B	400[B]	-	-	3/40/49/49	0/4/4/4

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	400[A]	63O	C2-N1	2.54	1.40	1.36
5	B	400[B]	63O	C2-N1	2.16	1.40	1.36

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	400[A]	63O	C25-C30-C31	2.95	121.33	116.39
5	B	400[A]	63O	C2-N1-N2	-2.28	110.87	112.06
5	B	400[B]	63O	C25-C30-C31	2.01	119.75	116.39

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	400[B]	63O	C23-C24-C25-C29
5	B	400[A]	63O	C20-C15-N4-C14
5	B	400[A]	63O	C16-C15-N4-C14
5	B	400[B]	63O	C23-C24-C25-C26
5	B	400[B]	63O	C23-C24-C25-C30
5	B	400[A]	63O	N3-C6-C7-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	400[A]	63O	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	100/132 (75%)	1.62	38 (38%) 1 1	31, 66, 190, 216	0
1	B	88/132 (66%)	1.55	36 (40%) 0 1	28, 55, 132, 173	0
2	C	220/237 (92%)	0.01	7 (3%) 50 54	20, 32, 54, 112	0
2	H	220/237 (92%)	0.12	9 (4%) 41 46	23, 33, 59, 85	0
3	D	213/214 (99%)	0.36	15 (7%) 22 25	20, 43, 72, 103	1 (0%)
3	L	213/214 (99%)	0.04	5 (2%) 61 64	18, 36, 53, 97	1 (0%)
4	I	14/16 (87%)	2.77	10 (71%) 0 0	105, 112, 131, 131	0
All	All	1068/1182 (90%)	0.42	120 (11%) 10 11	18, 37, 101, 216	2 (0%)

All (120) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	24	VAL	8.0
3	D	210	THR	7.4
1	B	37	PRO	7.3
1	A	131	VAL	6.8
4	I	1	ILE	6.4
1	A	22	VAL	6.4
1	B	62	TYR	5.9
1	B	59	PRO	5.3
3	L	210	THR	5.3
1	B	38	LYS	5.2
2	C	215	SER	5.1
1	B	108	ASN	5.0
1	A	26	LEU	5.0
1	B	65	VAL	4.7
1	B	36	ASN	4.6
1	A	23	MET	4.5
1	B	101	ARG	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	110	PHE	4.3
1	A	28	ILE	4.2
1	B	67	TRP	4.2
1	B	129	HIS	4.2
1	B	54	HIS	4.2
2	C	214	LYS	4.1
1	A	102	GLU	4.1
1	A	65	VAL	4.0
1	A	108	ASN	4.0
1	A	130	HIS	4.0
1	A	25	ASN	3.9
3	L	157	LYS	3.9
1	A	109	SER	3.9
1	A	112	LEU	3.7
4	I	3	VAL	3.7
2	H	215	SER	3.7
1	A	29	HIS	3.6
1	A	38	LYS	3.6
1	A	62	TYR	3.6
4	I	10	TRP	3.5
1	A	129	HIS	3.5
1	B	79	ALA	3.5
1	B	99	LEU	3.4
1	B	109	SER	3.4
3	D	111	LYS	3.3
3	D	157	LYS	3.3
4	I	14	ASN	3.3
1	B	55	ARG	3.2
1	A	30	ASN	3.2
4	I	2	HIS	3.2
2	C	191	THR	3.1
3	D	8	HIS	3.1
1	B	60	GLU	3.0
1	B	39	ARG	3.0
1	A	97	LEU	3.0
3	L	94[A]	TYR	3.0
1	B	100	ARG	3.0
1	A	59	PRO	3.0
1	A	101	ARG	2.9
1	A	36	ASN	2.9
2	H	164	HIS	2.9
1	A	110	PHE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	56	ASN	2.8
4	I	12	TRP	2.8
1	A	115	ILE	2.8
1	B	58	ASP	2.8
1	B	97	LEU	2.7
3	D	80	THR	2.7
1	B	111	ARG	2.7
3	D	1	ASN	2.7
1	B	113	GLU	2.7
2	C	190	GLY	2.7
1	A	55	ARG	2.7
1	A	99	LEU	2.7
1	B	114	LYS	2.7
1	A	79	ALA	2.7
1	A	27	ASN	2.6
3	D	46	ILE	2.6
1	B	63	PRO	2.6
1	B	112	LEU	2.6
1	B	64	SER	2.6
3	D	94[A]	TYR	2.6
2	H	129	LYS	2.6
3	D	20	THR	2.6
3	D	127	GLN	2.5
1	B	115	ILE	2.5
4	I	5	ILE	2.5
4	I	13	ILE	2.5
4	I	7	ALA	2.5
2	H	201	LYS	2.5
1	A	37	PRO	2.5
1	A	54	HIS	2.5
1	A	98	VAL	2.5
1	A	119	VAL	2.5
2	H	191	THR	2.5
2	H	193	THR	2.5
3	L	60	ASP	2.4
1	A	96	ILE	2.4
1	B	66	ILE	2.4
2	C	128	SER	2.3
1	A	80	ASP	2.3
1	B	98	VAL	2.3
3	D	60	ASP	2.3
1	B	53	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	113	SER	2.2
3	D	9	SER	2.2
1	A	39	ARG	2.2
3	D	56	SER	2.2
1	B	40	SER	2.2
2	H	128	SER	2.2
1	B	70	LYS	2.2
1	A	67	TRP	2.2
1	A	92	ILE	2.1
3	L	187	LYS	2.1
1	B	80	ASP	2.1
4	I	4	THR	2.1
2	C	212	GLU	2.1
2	C	3	GLN	2.1
1	B	57	GLU	2.0
1	A	81	GLY	2.0
3	D	12	SER	2.0
3	D	190	ARG	2.0
2	H	214	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	63O	B	400[A]	44/44	0.79	0.25	74,85,87,88	44
5	63O	B	400[B]	44/44	0.79	0.25	60,66,75,76	44

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 63O B 400 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 63O B 400 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.