



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 11:08 AM UTC

PDB ID : 5HI4 / pdb_00005hi4
Title : Binding site elucidation and structure guided design of macrocyclic IL-17A antagonists
Authors : Liu, S.
Deposited on : 2016-01-11
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	132	26% 58%9%33%
1	B	132	21% 58%11%31%
2	C	237	3% 86%5%7%
2	H	237	5% 86%6%7%
3	D	214	6% 86%13%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	L	214	2% 88% 10%
4	I	16	75% 81% 6% 12%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 8461 atoms, of which 39 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interleukin-17A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	89	Total	C	N	O	S	0	0	0
			718	449	131	133	5			
1	B	91	Total	C	N	O	S	0	0	0
			731	457	135	134	5			

- Molecule 2 is a protein called CAT-2000 FAB heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	220	Total	C	N	O	S	0	0	0
			1622	1017	278	321	6			
2	H	220	Total	C	N	O	S	0	0	0
			1622	1017	278	321	6			


- Molecule 3 is a protein called CAT-2000 FAB light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	213	Total	C	N	O	S	0	1	0
			1618	1015	267	331	5			
3	L	213	Total	C	N	O	S	0	1	0
			1618	1015	267	331	5			

- Molecule 4 is a protein called synthetic IL-17A peptide inhibitor.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	I	14	Total	C	N	O	0	0	0
			120	81	19	20			

- Molecule 5 is (9'S,17'R)-6'-chloro-N-methyl-9'-{[(1-methyl-1H-pyrazol-5-yl)carbonyl]amino}-10',19'-dioxo-2'-oxa-11',18'-diazaspiro[cyclopentane-1,21'-tetracyclo[20.2.2.2 12,15 .1 3,7]nonacosane]-1'(24'),3'(29'),4',6',12',14',22',25',27'-nonaene-17'-carboxamide (CCD ID: 63P) (formula: C₃₇H₃₉ClN₆O₅).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
5	A	1	Total	C	Cl	H	N	O	39	0
			88	37	1	39	6	5		

• Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	16	Total	O	0	0
			16	16		
6	B	17	Total	O	0	0
			17	17		
6	C	72	Total	O	0	0
			72	72		
6	D	45	Total	O	0	0
			45	45		
6	H	95	Total	O	0	0
			95	95		
6	L	79	Total	O	0	0
			79	79		



i

- Molecule 1: Interleukin-17A









- Molecule 3: CAT-2000 FAB light chain



- Molecule 3: CAT-2000 FAB light chain



- Molecule 4: synthetic IL-17A peptide inhibitor



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	91.24Å 68.19Å 100.12Å 90.00° 91.33° 90.00°	Depositor
Resolution (Å)	50.05 – 1.80 50.05 – 1.80	Depositor EDS
% Data completeness (in resolution range)	85.6 (50.05-1.80) 85.6 (50.05-1.80)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	17.80 (at 1.79Å)	Xtriage
Refinement program	BUSTER 2.11.6	Depositor
R, R_{free}	0.200 , 0.224 0.203 , 0.232	Depositor DCC
R_{free} test set	4869 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	0.056	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.027 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8461	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.36% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: 63P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.70	0/737	1.02	0/1004
1	B	0.72	0/750	1.05	1/1022 (0.1%)
2	C	0.88	1/1658 (0.1%)	1.04	1/2256 (0.0%)
2	H	0.85	1/1658 (0.1%)	1.05	1/2256 (0.0%)
3	D	0.76	0/1660	1.09	6/2269 (0.3%)
3	L	0.80	0/1660	1.10	3/2269 (0.1%)
4	I	0.84	0/125	1.32	0/174
All	All	0.81	2/8248 (0.0%)	1.07	12/11250 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	82	MET	SD-CE	-9.00	1.57	1.79
2	H	82	MET	SD-CE	-6.73	1.62	1.79

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	H	67	PHE	CA-CB-CG	7.02	120.82	113.80
2	C	67	PHE	CA-CB-CG	6.81	120.61	113.80
3	L	92	ASP	CA-CB-CG	6.45	119.05	112.60
3	D	92	ASP	CA-CB-CG	6.12	118.72	112.60
3	D	182	THR	N-CA-C	-5.69	102.74	110.36
3	D	164	THR	N-CA-C	-5.56	102.76	109.83
1	B	88	ASN	CA-CB-CG	5.42	118.02	112.60
3	D	151	ALA	N-CA-C	-5.33	98.55	107.99
3	L	47	VAL	N-CA-C	-5.31	106.62	111.67
3	D	60	ASP	CA-C-N	5.08	128.04	120.31
3	D	60	ASP	C-N-CA	5.08	128.04	120.31
3	L	182	THR	N-CA-C	-5.02	103.63	110.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	718	0	686	7	0
1	B	731	0	700	8	0
2	C	1622	0	1601	8	0
2	H	1622	0	1601	10	0
3	D	1618	0	1555	9	0
3	L	1618	0	1555	12	0
4	I	120	0	115	1	0
5	A	49	39	0	0	0
6	A	16	0	0	1	0
6	B	17	0	0	0	0
6	C	72	0	0	0	0
6	D	45	0	0	1	0
6	H	95	0	0	0	0
6	L	79	0	0	1	0
All	All	8422	39	7813	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:82:MET:HE2	2:C:82(C):LEU:HD21	1.49	0.93
3:L:28:LEU:H	3:L:69:ASN:HD21	1.29	0.80
1:B:125:THR:HG22	1:B:127:ILE:H	1.51	0.76
6:A:4103:HOH:O	1:B:125:THR:HB	1.87	0.74
1:A:125:THR:HG22	1:A:127:ILE:H	1.52	0.74
3:D:28:LEU:H	3:D:69:ASN:HD21	1.43	0.66
3:L:28:LEU:H	3:L:69:ASN:ND2	1.93	0.66
3:D:28:LEU:H	3:D:69:ASN:ND2	1.98	0.62
2:H:13:GLN:HG3	2:H:14:PRO:HD2	1.82	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:181:VAL:HG11	3:L:136:LEU:CD1	2.34	0.58
1:A:92:ILE:HD11	1:A:122:THR:HB	1.87	0.57
1:B:92:ILE:HD11	1:B:122:THR:HB	1.86	0.56
2:H:82:MET:HE2	2:H:82(C):LEU:HD21	1.90	0.54
1:B:76:CYS:HB2	1:B:87:MET:HG3	1.90	0.53
2:C:67:PHE:CZ	2:C:82:MET:HE3	2.44	0.52
3:L:78:LEU:HD21	3:L:106:VAL:HG22	1.91	0.51
2:H:181:VAL:CG1	3:L:136:LEU:CD1	2.89	0.49
3:L:47:VAL:HG12	3:L:48:ILE:HG12	1.95	0.49
2:C:82:MET:HE2	2:C:82(C):LEU:CD2	2.34	0.49
2:C:151:THR:HG22	2:C:199:ASN:HB3	1.95	0.49
1:A:86:HIS:HE1	6:L:337:HOH:O	1.95	0.48
1:A:76:CYS:HB2	1:A:87:MET:HG3	1.94	0.48
2:H:181:VAL:HG11	3:L:136:LEU:HD12	1.95	0.48
1:B:84:ASP:OD1	1:B:86:HIS:HD2	1.96	0.47
3:L:185:GLN:O	3:L:189:HIS:HD2	1.97	0.47
3:D:20:THR:HB	3:D:74:THR:HG22	1.97	0.47
1:B:86:HIS:HE1	6:D:324:HOH:O	1.97	0.47
1:A:84:ASP:OD1	1:A:86:HIS:HD2	1.97	0.46
3:D:47:VAL:HG12	3:D:48:ILE:HG12	1.97	0.46
3:D:185:GLN:O	3:D:189:HIS:HD2	1.99	0.46
2:H:150:VAL:HG23	2:H:178:LEU:HD21	1.97	0.46
3:D:36:TYR:CE1	3:D:46:ILE:HG22	2.53	0.43
3:D:195:GLN:HG2	3:D:204:GLU:HG3	2.01	0.43
2:H:119:PRO:HB3	2:H:145:TYR:HB3	2.01	0.43
1:A:92:ILE:HD12	1:B:92:ILE:HD12	2.01	0.42
1:A:109:SER:HA	4:I:2:HIS:O	2.20	0.42
1:B:36:ASN:HA	1:B:37:PRO:HD3	1.95	0.42
2:H:181:VAL:CG1	3:L:136:LEU:HD13	2.50	0.42
3:L:4:LEU:HD11	3:L:90:THR:HG22	2.02	0.41
2:C:214:LYS:NZ	3:D:123:SER:H	2.18	0.41
3:D:4:LEU:HD11	3:D:90:THR:HG22	2.02	0.41
3:L:195:GLN:HG2	3:L:204:GLU:HG3	2.02	0.41
2:C:127:SER:O	2:C:131:THR:HG23	2.20	0.41
2:H:178:LEU:C	2:H:178:LEU:HD12	2.46	0.41
2:C:145:TYR:CE2	2:C:150:VAL:HG13	2.55	0.40
2:C:67:PHE:CE2	2:C:82:MET:HE3	2.57	0.40
2:H:167:PRO:HG2	3:L:166:SER:OG	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	85/132 (64%)	85 (100%)	0	0	100	100
1	B	87/132 (66%)	87 (100%)	0	0	100	100
2	C	218/237 (92%)	212 (97%)	6 (3%)	0	100	100
2	H	218/237 (92%)	212 (97%)	5 (2%)	1 (0%)	24	14
3	D	212/214 (99%)	206 (97%)	6 (3%)	0	100	100
3	L	212/214 (99%)	207 (98%)	5 (2%)	0	100	100
4	I	12/16 (75%)	12 (100%)	0	0	100	100
All	All	1044/1182 (88%)	1021 (98%)	22 (2%)	1 (0%)	48	34

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	H	132	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/124 (67%)	80 (96%)	3 (4%)	31	18
1	B	84/124 (68%)	81 (96%)	3 (4%)	31	18
2	C	182/196 (93%)	177 (97%)	5 (3%)	39	27
2	H	182/196 (93%)	176 (97%)	6 (3%)	33	21
3	D	185/186 (100%)	172 (93%)	13 (7%)	14	4

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	L	185/186 (100%)	175 (95%)	10 (5%)	20	8
4	I	13/14 (93%)	13 (100%)	0	100	100
All	All	914/1026 (89%)	874 (96%)	40 (4%)	25	13

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	ASP
1	A	112	LEU
1	A	115	ILE
1	B	80	ASP
1	B	98	VAL
1	B	112	LEU
2	C	12	VAL
2	C	35	SER
2	C	148	GLN
2	C	191	THR
2	C	206	LYS
3	D	13	GLU
3	D	19	VAL
3	D	20	THR
3	D	28	LEU
3	D	46	ILE
3	D	56	SER
3	D	80	THR
3	D	81	GLU
3	D	103	LYS
3	D	127	GLN
3	D	157	LYS
3	D	160	VAL
3	D	203	VAL
2	H	13	GLN
2	H	35	SER
2	H	120	SER
2	H	148	GLN
2	H	191	THR
2	H	206	LYS
3	L	19	VAL
3	L	28	LEU
3	L	46	ILE
3	L	56	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	L	69	ASN
3	L	81	GLU
3	L	103	LYS
3	L	156	VAL
3	L	157	LYS
3	L	160	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (27) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	82	ASN
1	A	86	HIS
1	B	36	ASN
1	B	82	ASN
1	B	86	HIS
2	C	3	GLN
2	C	39	GLN
2	C	82(A)	ASN
2	C	98	HIS
2	C	105	GLN
2	C	192	GLN
2	C	204	ASN
3	D	38	GLN
3	D	69	ASN
3	D	89	GLN
3	D	127	GLN
3	D	195	GLN
2	H	3	GLN
2	H	82(A)	ASN
2	H	98	HIS
2	H	105	GLN
2	H	192	GLN
2	H	199	ASN
3	L	69	ASN
3	L	109	GLN
3	L	127	GLN
3	L	195	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	63P	A	4000	-	54,54,54	1.06	3 (5%)	68,77,77	1.19	5 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	63P	A	4000	-	-	2/49/58/58	0/5/6/6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	A	4000	63P	C2-N1	4.22	1.39	1.33
5	A	4000	63P	C33-N4	2.74	1.39	1.34
5	A	4000	63P	C32-N3	2.48	1.39	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	4000	63P	C34-N6-N5	4.35	114.34	112.06

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	4000	63P	C35-C34-N6	-4.24	103.98	106.22
5	A	4000	63P	C30-C26-C27	-3.69	97.77	102.40
5	A	4000	63P	C36-N5-N6	-2.60	103.34	104.41
5	A	4000	63P	C11-C12-N4	-2.52	104.31	111.11

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	4000	63P	C26-C31-C32-O4
5	A	4000	63P	C26-C31-C32-N3

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	89/132 (67%)	1.49	34 (38%) 1 0	23, 45, 118, 135	0
1	B	91/132 (68%)	1.22	28 (30%) 1 1	25, 50, 106, 129	0
2	C	220/237 (92%)	0.05	8 (3%) 46 46	17, 26, 46, 104	0
2	H	220/237 (92%)	0.16	12 (5%) 30 29	18, 27, 48, 74	0
3	D	213/214 (99%)	0.49	13 (6%) 27 25	17, 36, 64, 90	1 (0%)
3	L	213/214 (99%)	0.08	5 (2%) 61 61	14, 28, 44, 90	1 (0%)
4	I	14/16 (87%)	3.59	12 (85%) 0 0	85, 92, 119, 122	0
All	All	1060/1182 (89%)	0.43	112 (10%) 11 9	14, 30, 81, 135	2 (0%)

All (112) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	I	1	ILE	7.0
3	D	210	THR	6.4
1	A	104	PRO	5.7
3	L	210	THR	5.6
4	I	3	VAL	5.5
4	I	10	TRP	5.2
1	A	62	TYR	5.0
1	A	107	PRO	4.9
1	A	103	PRO	4.9
2	C	190	GLY	4.9
1	B	103	PRO	4.8
1	A	67	TRP	4.7
1	B	104	PRO	4.6
3	D	157	LYS	4.6
2	C	215	SER	4.5
1	A	79	ALA	4.5
4	I	2	HIS	4.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	110	PHE	4.3
1	A	59	PRO	4.3
1	B	129	HIS	4.2
4	I	7	ALA	4.0
1	A	60	GLU	4.0
3	D	94[A]	TYR	3.9
2	C	214	LYS	3.9
3	L	157	LYS	3.9
1	A	99	LEU	3.8
4	I	4	THR	3.8
2	H	215	SER	3.8
3	D	46	ILE	3.7
2	C	191	THR	3.7
1	A	101	ARG	3.7
1	A	54	HIS	3.7
2	H	131	THR	3.7
4	I	13	ILE	3.6
4	I	12	TRP	3.6
1	B	110	PHE	3.5
1	A	65	VAL	3.5
1	A	108	ASN	3.5
3	D	111	LYS	3.5
1	A	109	SER	3.4
1	B	62	TYR	3.4
1	B	109	SER	3.3
1	B	38	LYS	3.3
1	B	102	GLU	3.3
1	B	108	ASN	3.3
4	I	5	ILE	3.3
1	A	97	LEU	3.3
1	A	129	HIS	3.2
2	H	190	GLY	3.2
1	A	39	ARG	3.2
1	A	100	ARG	3.1
1	B	101	ARG	3.1
1	A	58	ASP	3.0
1	B	36	ASN	3.0
2	H	129	LYS	2.9
2	H	164	HIS	2.9
2	H	191	THR	2.9
1	B	99	LEU	2.9
4	I	14	ASN	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	55	ARG	2.7
4	I	9	LEU	2.7
1	A	40	SER	2.7
3	L	94[A]	TYR	2.7
3	D	60	ASP	2.7
3	D	80	THR	2.7
2	C	5	LEU	2.7
1	A	102	GLU	2.7
1	A	64	SER	2.6
2	C	130	SER	2.6
3	D	8	HIS	2.6
3	D	127	GLN	2.6
1	B	96	ILE	2.6
1	B	65	VAL	2.6
1	B	98	VAL	2.6
2	C	128	SER	2.6
3	L	40	PRO	2.5
1	A	112	LEU	2.5
3	L	22	SER	2.5
2	H	207	VAL	2.5
1	B	111	ARG	2.5
1	A	66	ILE	2.5
3	D	9	SER	2.5
1	B	119	VAL	2.5
1	A	63	PRO	2.5
1	B	55	ARG	2.4
1	B	100	ARG	2.4
2	H	128	SER	2.4
2	H	130	SER	2.4
2	H	193	THR	2.4
1	A	113	GLU	2.4
3	D	153	SER	2.4
2	H	140	CYS	2.4
1	B	53	LEU	2.3
1	A	98	VAL	2.3
1	A	119	VAL	2.3
1	B	54	HIS	2.3
1	B	37	PRO	2.3
1	B	128	VAL	2.3
1	B	67	TRP	2.2
1	B	79	ALA	2.2
1	B	59	PRO	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	D	12	SER	2.2
1	A	115	ILE	2.2
3	D	20	THR	2.2
1	B	97	LEU	2.2
1	B	112	LEU	2.2
2	C	129	LYS	2.2
1	A	111	ARG	2.1
2	H	206	LYS	2.1
4	I	6	PRO	2.1
1	A	57	GLU	2.1
1	A	114	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	63P	A	4000	49/49	0.77	0.18	62,73,77,77	39

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 63P A 4000:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.