



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 08:57 AM UTC

PDB ID : 2HKR / pdb_00002hkr
Title : Structures of the carbinolamine and schiff-base intermediates in the reductive half-reaction of aromatic amine dehydrogenase (AADH) with p-methoxyphenylethylamine
Authors : Roujeinikova, A.; Leys, D.
Deposited on : 2006-07-05
Resolution : 1.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	122	80% 9% • 10%
1	H	122	3% 87% 13%
2	A	362	% 92% 8% •
2	B	362	92% 7% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	ZHZ	H	1501[A]	-	X	-	-
4	ZHZ	H	1501[B]	-	X	-	-

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 8897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Aromatic amine dehydrogenase, small subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	110	Total	C	N	O	S	0	5	0
			863	530	151	167	15			
1	H	122	Total	C	N	O	S	0	11	0
			987	603	170	199	15			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	109	TRQ	TRP	modified residue	UNP Q0VKG6
H	109	TRQ	TRP	modified residue	UNP Q0VKG6


- Molecule 2 is a protein called Aromatic amine dehydrogenase, large subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	360	Total	C	N	O	S	0	8	0
			2836	1792	492	538	14			
2	B	360	Total	C	N	O	S	0	3	0
			2804	1768	486	536	14			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	433	THR	-	expression tag	UNP Q0VKG7
B	433	THR	-	expression tag	UNP Q0VKG7

- Molecule 3 is 2-(4-METHOXYPHENYL)ETHANAMINE (CCD ID: ZHH) (formula: C₉H₁₃NO).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	D	1	Total	C	N	O	0	0
			11	9	1	1		

- Molecule 4 is 2-(4-METHOXYPHENYL)ACETAMIDE (CCD ID: ZHZ) (formula: $C_9H_{11}NO_2$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	H	1	Total	C	N	O	0	1
			13	10	1	2		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	130	Total 130	O 130	0	1
5	H	134	Total 134	O 134	0	2
5	A	578	Total 578	O 578	0	0
5	B	541	Total 541	O 541	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Aromatic amine dehydrogenase, small subunit

Chain D: 




- Molecule 1: Aromatic amine dehydrogenase, small subunit

Chain H: 



- Molecule 2: Aromatic amine dehydrogenase, large subunit

Chain A: 



- Molecule 2: Aromatic amine dehydrogenase, large subunit

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.85Å 88.81Å 80.04Å 90.00° 90.39° 90.00°	Depositor
Resolution (Å)	15.00 – 1.40 15.00 – 1.40	Depositor EDS
% Data completeness (in resolution range)	100.0 (15.00-1.40) 97.4 (15.00-1.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.76 (at 1.40Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.149 , 0.171 0.150 , 0.172	Depositor DCC
R_{free} test set	9515 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	13.1	Xtriage
Anisotropy	0.104	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.012 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8897	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZHZ, TRQ, ZHH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	D	0.68	0/886	0.68	0/1206
1	H	0.65	0/1005	0.70	0/1372
2	A	0.68	0/2929	0.68	0/3967
2	B	0.69	0/2878	0.70	0/3901
All	All	0.68	0/7698	0.69	0/10446

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	310	PRO	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	863	0	776	10	0
1	H	987	0	856	14	0
2	A	2836	0	2796	28	0
2	B	2804	0	2728	20	0
3	D	11	0	12	2	0
4	H	13	0	6	4	0
5	A	578	0	0	3	0
5	B	541	0	0	0	0
5	D	130	0	0	3	0
5	H	134	0	0	2	0
All	All	8897	0	7174	62	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:104:ARG:HH11	2:A:106:HIS:HE1	1.29	0.78
1:D:95[B]:THR:HG22	5:D:1536:HOH:O	1.85	0.75
1:H:105:SER:HA	1:H:164:ASN:HD21	1.52	0.75
2:B:104:ARG:HH11	2:B:106:HIS:HE1	1.32	0.74
1:H:110[B]:ILE:HD13	1:H:125[B]:SER:HA	1.70	0.73
5:H:1560:HOH:O	2:B:118[B]:MET:SD	2.48	0.72
2:A:124:ASN:HD21	2:A:178:GLY:H	1.36	0.71
2:B:91:TYR:OH	2:B:428:HIS:HD2	1.75	0.69
1:D:105:SER:HA	1:D:164:ASN:HD21	1.58	0.68
2:B:197:GLN:NE2	2:B:227:SER:H	1.95	0.65
2:A:91:TYR:OH	2:A:428:HIS:HD2	1.80	0.65
2:A:197:GLN:NE2	2:A:227:SER:H	1.97	0.62
2:A:335:ARG:H	2:A:386:GLN:HE22	1.46	0.61
2:B:335:ARG:H	2:B:386:GLN:HE22	1.48	0.61
2:A:335:ARG:H	2:A:386:GLN:NE2	2.01	0.58
2:B:197:GLN:HE21	2:B:227:SER:H	1.51	0.58
2:A:271:LYS:NZ	5:A:905:HOH:O	2.35	0.57
2:B:335:ARG:H	2:B:386:GLN:NE2	2.01	0.57
3:D:1502:ZHH:HF12	2:B:179:LEU:HD23	1.86	0.57
1:D:167:SER:HB2	2:B:177:GLN:HE22	1.68	0.57
2:A:197:GLN:HE21	2:A:227:SER:H	1.51	0.56
2:A:91:TYR:OH	2:A:428:HIS:CD2	2.59	0.56
1:H:167:SER:HB2	2:A:177:GLN:HE22	1.72	0.55
2:B:106:HIS:HD2	2:B:418:ALA:O	1.90	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:71:GLU:N	5:D:1588:HOH:O	2.43	0.51
2:A:128:GLN:HE22	2:A:186:ARG:C	2.19	0.51
4:H:1501[B]:ZHZ:HF11	2:A:124:ASN:CG	2.36	0.50
2:A:106:HIS:HD2	2:A:418:ALA:O	1.95	0.50
2:A:248:LEU:HD13	2:A:263:ARG:HD3	1.92	0.50
1:D:146:GLU:CD	2:B:354:LYS:HZ1	2.19	0.50
2:A:133:GLY:O	2:A:428:HIS:HE1	1.96	0.49
2:A:124:ASN:ND2	2:A:178:GLY:H	2.09	0.49
2:B:128:GLN:HE22	2:B:186:ARG:C	2.20	0.49
1:D:109:TRQ:HB2	1:D:160:TRP:NE1	2.29	0.48
2:A:131:ASN:H	2:A:187:GLN:HE22	1.61	0.47
1:H:159:ASN:HD22	2:A:177:GLN:NE2	2.11	0.47
2:A:143:HIS:HD2	5:A:622:HOH:O	1.96	0.47
1:D:78:TRP:CE2	1:D:79[B]:ARG:HG3	2.50	0.47
1:H:173[B]:THR:HG21	2:B:76:LEU:CD1	2.45	0.46
2:B:275:ILE:HA	2:B:290:SER:HA	1.97	0.46
1:H:155:HIS:HE1	5:A:457:HOH:O	1.97	0.46
1:H:146:GLU:CD	2:A:354:LYS:HZ1	2.23	0.46
5:D:1535:HOH:O	2:B:143:HIS:HE1	1.99	0.45
2:A:275:ILE:HA	2:A:290:SER:HA	2.00	0.44
1:H:159:ASN:HB2	4:H:1501[B]:ZHZ:HF13	2.00	0.44
2:B:343:PHE:HB3	2:B:357:ALA:HB2	2.00	0.43
1:H:173[B]:THR:HG21	2:B:76:LEU:HD13	2.00	0.43
5:H:1539:HOH:O	2:A:143:HIS:HE1	2.01	0.43
1:H:108:SER:HB2	1:H:126[B]:TYR:O	2.18	0.43
1:D:173:THR:HG21	2:A:76:LEU:CD1	2.48	0.43
4:H:1501[A]:ZHZ:HF11	2:A:124:ASN:OD1	2.19	0.43
1:D:173:THR:HG21	2:A:76:LEU:HD13	2.01	0.42
2:A:186:ARG:HH22	2:A:426:GLN:NE2	2.17	0.42
1:D:159:ASN:HD22	2:B:177:GLN:NE2	2.18	0.42
4:H:1501[B]:ZHZ:HF11	2:A:124:ASN:ND2	2.36	0.41
2:A:131:ASN:H	2:A:187:GLN:NE2	2.19	0.40
2:B:334:HIS:HA	2:B:386:GLN:HE22	1.86	0.40
3:D:1502:ZHH:HF11	2:B:124:ASN:ND2	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	112/122 (92%)	110 (98%)	2 (2%)	0	100	100
1	H	129/122 (106%)	126 (98%)	3 (2%)	0	100	100
2	A	366/362 (101%)	352 (96%)	13 (4%)	1 (0%)	36	16
2	B	360/362 (99%)	346 (96%)	12 (3%)	2 (1%)	21	5
All	All	967/968 (100%)	934 (97%)	30 (3%)	3 (0%)	36	16

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	216	VAL
2	A	200	SER
2	B	200	SER

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	100/106 (94%)	100 (100%)	0	100	100
1	H	113/106 (107%)	113 (100%)	0	100	100
2	A	312/306 (102%)	310 (99%)	2 (1%)	78	56
2	B	305/306 (100%)	305 (100%)	0	100	100
All	All	830/824 (101%)	828 (100%)	2 (0%)	88	71

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	406[A]	GLN
2	A	406[B]	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (38) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	139	GLN
1	D	143	GLN
1	D	164	ASN
1	H	63	ASN
1	H	139	GLN
1	H	143	GLN
1	H	155	HIS
1	H	164	ASN
2	A	106	HIS
2	A	124	ASN
2	A	128	GLN
2	A	143	HIS
2	A	177	GLN
2	A	180	ASN
2	A	187	GLN
2	A	197	GLN
2	A	231	GLN
2	A	251	ASN
2	A	386	GLN
2	A	424	GLN
2	A	426	GLN
2	A	428	HIS
2	B	106	HIS
2	B	128	GLN
2	B	143	HIS
2	B	177	GLN
2	B	180	ASN
2	B	187	GLN
2	B	197	GLN
2	B	231	GLN
2	B	251	ASN
2	B	322	ASN
2	B	369	GLN
2	B	386	GLN
2	B	406	GLN
2	B	424	GLN
2	B	426	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	428	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

3 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	TRQ	H	109[A]	1	15,16,18	3.20	4 (26%)	13,22,26	1.34	2 (15%)
1	TRQ	H	109[B]	1	15,16,18	3.21	4 (26%)	13,22,26	1.34	2 (15%)
1	TRQ	D	109	1,3	15,16,18	2.85	5 (33%)	13,22,26	1.32	2 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TRQ	H	109[A]	1	-	0/5/16/21	0/2/2/2
1	TRQ	H	109[B]	1	-	0/5/16/21	0/2/2/2
1	TRQ	D	109	1,3	-	0/5/16/21	0/2/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	109[A]	TRQ	O7-CZ2	10.30	1.44	1.23
1	H	109[B]	TRQ	O7-CZ2	10.30	1.44	1.23
1	D	109	TRQ	O7-CZ2	9.26	1.41	1.23
1	H	109[A]	TRQ	CE2-CZ2	-3.61	1.38	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	109[B]	TRQ	CE2-CZ2	-3.61	1.38	1.45
1	H	109[A]	TRQ	CE3-CZ3	3.23	1.41	1.33
1	H	109[B]	TRQ	CE3-CZ3	3.23	1.41	1.33
1	D	109	TRQ	CE2-CZ2	-3.13	1.39	1.45
1	H	109[A]	TRQ	CH2-CZ3	-3.07	1.41	1.49
1	H	109[B]	TRQ	CH2-CZ3	-3.07	1.41	1.49
1	D	109	TRQ	CH2-CZ3	-2.92	1.41	1.49
1	D	109	TRQ	CE3-CZ3	2.41	1.39	1.33
1	D	109	TRQ	CD2-CG	2.20	1.46	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	109[A]	TRQ	CZ3-CH2-CZ2	3.46	121.07	113.35
1	H	109[B]	TRQ	CZ3-CH2-CZ2	3.46	121.07	113.35
1	D	109	TRQ	CZ3-CH2-CZ2	3.30	120.71	113.35
1	D	109	TRQ	O7-CZ2-CE2	-2.21	119.84	122.86
1	H	109[A]	TRQ	O7-CZ2-CE2	-2.04	120.08	122.86
1	H	109[B]	TRQ	O7-CZ2-CE2	-2.04	120.08	122.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	109	TRQ	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ZHH	D	1502	1	11,11,11	1.02	1 (9%)	13,13,13	0.82	0
4	ZHZ	H	1501[B]	-	12,12,12	6.58	5 (41%)	15,15,15	2.68	7 (46%)
4	ZHZ	H	1501[A]	-	12,12,12	6.58	5 (41%)	15,15,15	2.66	7 (46%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ZHH	D	1502	1	-	3/5/5/5	0/1/1/1
4	ZHZ	H	1501[B]	-	-	2/6/6/6	0/1/1/1
4	ZHZ	H	1501[A]	-	-	2/6/6/6	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	1501[A]	ZHZ	CE2-CD2	21.29	1.73	1.38
4	H	1501[B]	ZHZ	CE2-CD2	21.29	1.73	1.38
4	H	1501[A]	ZHZ	O2-CA	5.88	1.42	1.23
4	H	1501[B]	ZHZ	O2-CA	5.88	1.42	1.23
4	H	1501[A]	ZHZ	CA-N2	4.42	1.47	1.32
4	H	1501[B]	ZHZ	CA-N2	4.42	1.47	1.32
3	D	1502	ZHH	O1-CZ	2.92	1.43	1.37
4	H	1501[A]	ZHZ	O1-CZ	2.27	1.42	1.37
4	H	1501[B]	ZHZ	O1-CZ	2.27	1.42	1.37
4	H	1501[A]	ZHZ	CE1-CZ	2.02	1.42	1.38
4	H	1501[B]	ZHZ	CE1-CZ	2.02	1.42	1.38

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1501[A]	ZHZ	O2-CA-N2	-6.33	105.61	122.53
4	H	1501[B]	ZHZ	O2-CA-N2	-6.33	105.61	122.53
4	H	1501[A]	ZHZ	O2-CA-CB	-4.94	110.51	121.03
4	H	1501[B]	ZHZ	O2-CA-CB	-4.94	110.51	121.03
4	H	1501[A]	ZHZ	CD2-CE2-CZ	-3.17	116.11	119.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	1501[B]	ZHZ	CD2-CE2-CZ	-3.17	116.11	119.73
4	H	1501[A]	ZHZ	CD1-CE1-CZ	2.98	123.14	119.73
4	H	1501[B]	ZHZ	CD1-CE1-CZ	2.98	123.14	119.73
4	H	1501[A]	ZHZ	CE2-CD2-CG	-2.87	117.23	121.00
4	H	1501[B]	ZHZ	CE2-CD2-CG	-2.87	117.23	121.00
4	H	1501[A]	ZHZ	CB-CA-N2	-2.66	111.93	116.54
4	H	1501[B]	ZHZ	CB-CA-N2	-2.66	111.93	116.54
4	H	1501[A]	ZHZ	CE1-CD1-CG	2.54	124.33	121.00
4	H	1501[B]	ZHZ	CE1-CD1-CG	2.54	124.33	121.00

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	1502	ZHH	CE2-CZ-O1-CF1
3	D	1502	ZHH	CE1-CZ-O1-CF1
4	H	1501[A]	ZHZ	CE2-CZ-O1-CF1
4	H	1501[A]	ZHZ	CE1-CZ-O1-CF1
4	H	1501[B]	ZHZ	CE2-CZ-O1-CF1
4	H	1501[B]	ZHZ	CE1-CZ-O1-CF1
3	D	1502	ZHH	N2-CA-CB-CG

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1502	ZHH	2	0
4	H	1501[B]	ZHZ	3	0
4	H	1501[A]	ZHZ	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	D	109/122 (89%)	-0.20	0 100 100	9, 14, 25, 34	5 (4%)
1	H	121/122 (99%)	0.06	4 (3%) 49 51	6, 14, 41, 48	10 (8%)
2	A	360/362 (99%)	-0.58	4 (1%) 78 81	5, 11, 23, 33	8 (2%)
2	B	360/362 (99%)	-0.21	0 100 100	5, 15, 27, 41	3 (0%)
All	All	950/968 (98%)	-0.32	8 (0%) 82 85	5, 13, 28, 48	26 (2%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	60	ILE	3.2
2	A	432	GLY	2.9
1	H	180	ALA	2.6
2	A	431	GLY	2.4
2	A	433	THR	2.4
1	H	68	ASN	2.2
1	H	123	LEU	2.2
2	A	75	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	TRQ	D	109	15/17	0.95	0.06	12,13,16,16	0
1	TRQ	H	109[A]	15/17	0.97	0.04	9,12,16,17	2
1	TRQ	H	109[B]	15/17	0.97	0.04	9,12,14,15	2

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	ZHZ	H	1501[A]	12/12	0.85	0.11	18,24,29,32	1
4	ZHZ	H	1501[B]	12/12	0.85	0.11	18,24,29,32	1
3	ZHH	D	1502	11/11	0.87	0.11	19,26,36,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.