



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 05:16 AM UTC

PDB ID : 4HKI / pdb_00004hki
Title : Tankyrase 2 in complex with flavone
Authors : Narwal, M.; Haikarainen, T.; Lehtio, L.
Deposited on : 2012-10-15
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2057 (2.16-2.16)
Clashscore	190562	2159 (2.16-2.16)
Ramachandran outliers	187476	2134 (2.16-2.16)
Sidechain outliers	187428	2133 (2.16-2.16)
RSRZ outliers	180081	2059 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	191	 2% 77% 7% 16%
1	H	191	 2% 79% 6% 15%
2	C	49	 6% 88% 8% .
2	D	49	 4% 90% 6% .

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3660 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tankyrase-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	161	Total	C	N	O	S	0	0	0
			1296	813	240	235	8			
1	H	162	Total	C	N	O	S	0	3	0
			1318	826	243	240	9			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	923	MET	-	expression tag	UNP Q9H2K2
A	924	HIS	-	expression tag	UNP Q9H2K2
A	925	HIS	-	expression tag	UNP Q9H2K2
A	926	HIS	-	expression tag	UNP Q9H2K2
A	927	HIS	-	expression tag	UNP Q9H2K2
A	928	HIS	-	expression tag	UNP Q9H2K2
A	929	HIS	-	expression tag	UNP Q9H2K2
A	930	SER	-	expression tag	UNP Q9H2K2
A	931	SER	-	expression tag	UNP Q9H2K2
A	932	GLY	-	expression tag	UNP Q9H2K2
A	933	VAL	-	expression tag	UNP Q9H2K2
A	934	ASP	-	expression tag	UNP Q9H2K2
A	935	LEU	-	expression tag	UNP Q9H2K2
A	936	GLY	-	expression tag	UNP Q9H2K2
A	937	THR	-	expression tag	UNP Q9H2K2
A	938	GLU	-	expression tag	UNP Q9H2K2
A	939	ASN	-	expression tag	UNP Q9H2K2
A	940	LEU	-	expression tag	UNP Q9H2K2
A	941	TYR	-	expression tag	UNP Q9H2K2
A	942	PHE	-	expression tag	UNP Q9H2K2
A	943	GLN	-	expression tag	UNP Q9H2K2
A	944	SER	-	expression tag	UNP Q9H2K2
A	945	MET	-	expression tag	UNP Q9H2K2
H	923	MET	-	expression tag	UNP Q9H2K2
H	924	HIS	-	expression tag	UNP Q9H2K2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	925	HIS	-	expression tag	UNP Q9H2K2
H	926	HIS	-	expression tag	UNP Q9H2K2
H	927	HIS	-	expression tag	UNP Q9H2K2
H	928	HIS	-	expression tag	UNP Q9H2K2
H	929	HIS	-	expression tag	UNP Q9H2K2
H	930	SER	-	expression tag	UNP Q9H2K2
H	931	SER	-	expression tag	UNP Q9H2K2
H	932	GLY	-	expression tag	UNP Q9H2K2
H	933	VAL	-	expression tag	UNP Q9H2K2
H	934	ASP	-	expression tag	UNP Q9H2K2
H	935	LEU	-	expression tag	UNP Q9H2K2
H	936	GLY	-	expression tag	UNP Q9H2K2
H	937	THR	-	expression tag	UNP Q9H2K2
H	938	GLU	-	expression tag	UNP Q9H2K2
H	939	ASN	-	expression tag	UNP Q9H2K2
H	940	LEU	-	expression tag	UNP Q9H2K2
H	941	TYR	-	expression tag	UNP Q9H2K2
H	942	PHE	-	expression tag	UNP Q9H2K2
H	943	GLN	-	expression tag	UNP Q9H2K2
H	944	SER	-	expression tag	UNP Q9H2K2
H	945	MET	-	expression tag	UNP Q9H2K2

- Molecule 2 is a protein called Tankyrase-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	47	Total	C	N	O	S	0	1	0
			381	243	68	68	2			
2	D	47	Total	C	N	O	S	0	0	0
			373	238	65	68	2			

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Zn	0	0
			1	1		
3	H	1	Total	Zn	0	0
			1	1		

- Molecule 4 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		
4	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is 2-PHENYL-4H-CHROMEN-4-ONE (CCD ID: FLN) (formula: C₁₅H₁₀O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			17	15	2		
5	H	1	Total	C	O	0	0
			17	15	2		

- Molecule 6 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			6	3	3		
6	H	1	Total	C	O	0	0
			6	3	3		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	C	1	Total	C	O	0	0
			7	4	3		
7	H	1	Total	C	O	0	0
			7	4	3		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	98	Total	O	0	0
			98	98		
8	C	18	Total	O	0	0
			18	18		
8	H	76	Total	O	0	0
			76	76		
8	D	18	Total	O	0	0
			18	18		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Tankyrase-2




• Molecule 1: Tankyrase-2



• Molecule 2: Tankyrase-2



• Molecule 2: Tankyrase-2



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	91.40Å 97.95Å 118.76Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.39 – 2.15 44.39 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.2 (44.39-2.15) 99.2 (44.39-2.15)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.60 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.191 , 0.229 0.194 , 0.231	Depositor DCC
R_{free} test set	1457 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	20.8	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 38.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3660	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FLN, PEG, SO4, GOL, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.01	1/1327 (0.1%)	0.91	0/1779
1	H	0.99	0/1358	0.89	0/1819
2	C	1.06	0/396	0.93	0/539
2	D	0.93	0/385	0.87	0/525
All	All	1.00	1/3466 (0.0%)	0.90	0/4662

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1031	HIS	CG-CD2	5.36	1.41	1.35

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1296	0	1241	8	0
1	H	1318	0	1267	11	0
2	C	381	0	370	3	0
2	D	373	0	357	5	0
3	A	1	0	0	0	0
3	H	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	10	0	0	1	0
4	H	10	0	0	1	0
5	A	17	0	10	1	0
5	H	17	0	10	1	0
6	C	6	0	8	0	0
6	H	6	0	8	1	0
7	C	7	0	10	0	0
7	H	7	0	10	0	0
8	A	98	0	0	0	0
8	C	18	0	0	1	0
8	D	18	0	0	0	0
8	H	76	0	0	1	0
All	All	3660	0	3291	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1111:SER:O	2:C:1130:SER:HB3	1.93	0.69
1:A:1111:SER:HB2	8:H:1360:HOH:O	1.97	0.63
1:H:1046:GLU:HA	2:D:1115:MET:HE2	1.82	0.60
2:C:1143[A]:ARG:NH2	2:C:1145:GLU:OE2	2.36	0.58
1:H:990:ASN:ND2	4:H:1203:SO4:O4	2.30	0.57
2:C:1128:ARG:HD3	8:C:1305:HOH:O	2.05	0.55
1:A:1033:SER:O	1:A:1036:VAL:HG23	2.08	0.52
1:A:990:ASN:ND2	4:A:1203:SO4:O4	2.34	0.52
1:H:1046:GLU:CA	2:D:1115:MET:HE2	2.40	0.51
1:H:1112:ALA:O	2:D:1131:VAL:CG2	2.61	0.48
1:H:988:ILE:HG23	6:H:1205:GOL:H11	1.97	0.47
1:H:972:MET:SD	1:H:1096:LEU:HD12	2.55	0.46
1:H:1032:GLY:O	5:H:1204:FLN:H3	2.16	0.45
1:A:1032:GLY:O	5:A:1204:FLN:H3	2.19	0.43
1:H:954:ILE:HD12	1:H:956:ILE:HD11	2.00	0.43
1:A:972:MET:HE3	1:A:992:TYR:CD2	2.54	0.43
1:H:1113:MET:N	1:H:1113:MET:HE2	2.35	0.42
1:A:1045:ASP:OD2	1:A:1047:ARG:NH2	2.46	0.41
1:A:1055:PHE:CE1	1:A:1109:GLN:HG2	2.56	0.41
1:H:1112:ALA:O	2:D:1131:VAL:HG21	2.21	0.41
1:H:1095:GLN:HA	2:D:1153:ILE:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/191 (83%)	157 (99%)	2 (1%)	0	100	100
1	H	163/191 (85%)	160 (98%)	3 (2%)	0	100	100
2	C	46/49 (94%)	42 (91%)	4 (9%)	0	100	100
2	D	45/49 (92%)	43 (96%)	2 (4%)	0	100	100
All	All	413/480 (86%)	402 (97%)	11 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	137/164 (84%)	135 (98%)	2 (2%)	57	64
1	H	141/164 (86%)	141 (100%)	0	100	100
2	C	40/40 (100%)	40 (100%)	0	100	100
2	D	39/40 (98%)	39 (100%)	0	100	100
All	All	357/408 (88%)	355 (99%)	2 (1%)	78	84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1003	LYS
1	A	1037	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1021	HIS
1	A	1095	GLN
1	A	1109	GLN
1	H	993	ASN
1	H	1011	HIS
1	H	1095	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 2 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	GOL	H	1205	-	5,5,5	0.39	0	5,5,5	0.53	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	FLN	H	1204	-	19,19,19	1.43	3 (15%)	26,26,26	1.36	4 (15%)
4	SO4	H	1203	-	4,4,4	0.49	0	6,6,6	0.48	0
7	PEG	H	1206	-	6,6,6	0.48	0	5,5,5	0.25	0
5	FLN	A	1204	-	19,19,19	1.41	3 (15%)	26,26,26	1.44	4 (15%)
7	PEG	C	1202	-	6,6,6	0.38	0	5,5,5	0.58	0
4	SO4	A	1202	-	4,4,4	0.36	0	6,6,6	1.27	1 (16%)
6	GOL	C	1201	-	5,5,5	0.38	0	5,5,5	0.42	0
4	SO4	A	1203	-	4,4,4	0.58	0	6,6,6	0.38	0
4	SO4	H	1202	-	4,4,4	0.40	0	6,6,6	1.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	GOL	H	1205	-	-	2/4/4/4	-
5	FLN	H	1204	-	-	0/4/4/4	0/3/3/3
7	PEG	H	1206	-	-	2/4/4/4	-
5	FLN	A	1204	-	-	0/4/4/4	0/3/3/3
7	PEG	C	1202	-	-	1/4/4/4	-
6	GOL	C	1201	-	-	2/4/4/4	-

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	H	1204	FLN	C4A-C4	-4.63	1.40	1.48
5	A	1204	FLN	C4A-C4	-3.48	1.42	1.48
5	A	1204	FLN	C1'-C2	-3.05	1.41	1.48
5	H	1204	FLN	C1'-C2	-2.34	1.42	1.48
5	A	1204	FLN	C3-C4	-2.28	1.39	1.44
5	H	1204	FLN	C3-C4	-2.17	1.40	1.44

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	A	1204	FLN	O1-C8A-C4A	-3.21	118.65	121.57
5	H	1204	FLN	C2-C3-C4	-2.98	119.07	122.14
5	A	1204	FLN	C8A-O1-C2	2.49	122.70	119.67
5	H	1204	FLN	O1-C8A-C8	2.43	119.48	116.25
5	H	1204	FLN	C5'-C4'-C3'	-2.40	116.58	119.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1202	SO4	O3-S-O2	-2.35	97.25	109.56
5	A	1204	FLN	C2-C3-C4	-2.34	119.73	122.14
5	A	1204	FLN	C6-C7-C8	-2.15	117.58	120.24
5	H	1204	FLN	O1-C8A-C4A	-2.01	119.74	121.57

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	C	1201	GOL	C1-C2-C3-O3
6	C	1201	GOL	O2-C2-C3-O3
7	H	1206	PEG	O1-C1-C2-O2
6	H	1205	GOL	O1-C1-C2-C3
7	H	1206	PEG	O2-C3-C4-O4
7	C	1202	PEG	O2-C3-C4-O4
6	H	1205	GOL	O1-C1-C2-O2

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	1205	GOL	1	0
5	H	1204	FLN	1	0
4	H	1203	SO4	1	0
5	A	1204	FLN	1	0
4	A	1203	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	161/191 (84%)	-0.25	3 (1%) 66 70	10, 18, 29, 52	0
1	H	162/191 (84%)	-0.17	4 (2%) 58 62	12, 19, 35, 55	3 (1%)
2	C	47/49 (95%)	0.06	3 (6%) 25 28	11, 18, 44, 52	1 (2%)
2	D	47/49 (95%)	0.11	2 (4%) 40 45	13, 22, 48, 57	0
All	All	417/480 (86%)	-0.14	12 (2%) 53 57	10, 19, 39, 57	4 (0%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1112	ALA	3.9
1	H	1113	MET	3.9
2	C	1131	VAL	3.8
2	D	1116	ALA	2.6
1	H	1008	ARG	2.5
1	H	1112	ALA	2.3
2	C	1132	ASN	2.3
2	C	1130	SER	2.2
1	H	1037[A]	ASN	2.2
1	A	1037	ASN	2.1
2	D	1161	GLU	2.1
1	A	1111	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	PEG	C	1202	7/7	0.82	0.16	40,43,49,50	0
4	SO4	A	1203	5/5	0.86	0.15	51,52,57,62	0
7	PEG	H	1206	7/7	0.88	0.14	52,54,56,57	0
6	GOL	H	1205	6/6	0.91	0.10	33,35,38,42	0
4	SO4	H	1203	5/5	0.91	0.12	46,48,51,53	0
6	GOL	C	1201	6/6	0.91	0.13	19,21,21,22	6
5	FLN	A	1204	17/17	0.96	0.06	12,13,15,15	0
5	FLN	H	1204	17/17	0.96	0.05	14,15,15,15	0
4	SO4	H	1202	5/5	0.97	0.06	19,20,23,23	0
4	SO4	A	1202	5/5	0.97	0.07	23,24,25,29	0
3	ZN	A	1201	1/1	0.99	0.02	25,25,25,25	0
3	ZN	H	1201	1/1	1.00	0.02	17,17,17,17	0

6.5 Other polymers [i](#)

There are no such residues in this entry.