



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 11:36 PM UTC

PDB ID : 6HKJ / pdb_00006hkj
Title : X-ray structure of human glutamate carboxypeptidase II (GCPII) in complex with a inhibitor RNA 2-19-1
Authors : Motlova, L.; Novakova, Z.; Barinka, C.
Deposited on : 2018-09-06
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	3% 87% 10% ..
2	B	2	50% 50%
2	C	2	100%
3	D	4	100%

2 Entry composition [i](#)


There are 10 unique types of molecules in this entry. The entry contains 6342 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	47	0
			5739	3687	962	1068	22			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		
4	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 5 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Zn	0	0
			2	2		


- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Ca	0	0
			1	1		

- Molecule 7 is CHLORIDE ION (CCD ID: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	2	Total	Cl	0	0
			2	2		

- Molecule 8 is TRIS(HYDROXYETHYL)AMINOMETHANE (CCD ID: TAM) (formula: $C_7H_{17}NO_3$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
8	A	1	Total	C	N	O	0	0
			11	7	1	3		

- Molecule 9 is (2 {S})-2-[[[(2 {S})-6-[(6-fluoranylpyridin-3-yl)amino]-1-oxidanyl-1,6-bis(oxidanylidene)hexan-2-yl]carbamoylamino]pentanedioic acid (CCD ID: G8W) (formula: C₁₇H₂₁FN₄O₈).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total	C	F	N	O	0	0
			30	17	1	4	8		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	395	Total	O	0	0
			395	395		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutamate carboxypeptidase 2




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	102.30Å 131.66Å 160.02Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.30 – 2.09 47.30 – 2.09	Depositor EDS
% Data completeness (in resolution range)	99.4 (47.30-2.09) 99.3 (47.30-2.09)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.10Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.160 , 0.204 0.175 , 0.212	Depositor DCC
R_{free} test set	2100 reflections (3.32%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.612	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 60.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6342	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: G8W, MAN, NAG, CA, TAM, CL, ZN, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.25	21/6062 (0.3%)	1.13	16/8206 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	705	PHE	N-CA	10.45	1.53	1.46
1	A	537	TYR	N-CA	6.49	1.54	1.45
1	A	707	GLY	C-O	6.37	1.31	1.23
1	A	573	HIS	CE1-NE2	6.21	1.38	1.32
1	A	691	ILE	N-CA	5.96	1.53	1.46
1	A	383	PHE	N-CA	5.82	1.53	1.46
1	A	165	SER	N-CA	5.56	1.52	1.45
1	A	79	GLN	N-CA	5.47	1.53	1.46
1	A	567	ASP	N-CA	5.45	1.52	1.46
1	A	76	ASN	C-O	5.44	1.30	1.24
1	A	207	LYS	N-CA	5.43	1.53	1.46
1	A	285	GLU	N-CA	5.35	1.52	1.46
1	A	461	THR	N-CA	5.34	1.52	1.46
1	A	90[A]	PHE	N-CA	5.34	1.52	1.46
1	A	90[B]	PHE	N-CA	5.34	1.52	1.46
1	A	528	LEU	N-CA	5.30	1.53	1.46
1	A	483	SER	C-O	-5.27	1.17	1.24
1	A	371	TYR	N-CA	5.23	1.52	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	441	LEU	C-O	5.16	1.30	1.24
1	A	527	ARG	N-CA	5.16	1.52	1.46
1	A	419	ALA	C-O	5.08	1.29	1.24

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	146	PRO	CA-C-N	-7.74	112.41	120.38
1	A	146	PRO	C-N-CA	-7.74	112.41	120.38
1	A	652	ASP	N-CA-C	5.94	119.09	107.57
1	A	295	HIS	CA-C-N	5.75	127.03	119.84
1	A	295	HIS	C-N-CA	5.75	127.03	119.84
1	A	705	PHE	CA-C-N	5.68	125.35	119.05
1	A	705	PHE	C-N-CA	5.68	125.35	119.05
1	A	240	LYS	CA-C-N	5.67	128.76	120.71
1	A	240	LYS	C-N-CA	5.67	128.76	120.71
1	A	543	THR	CB-CA-C	5.59	121.55	110.42
1	A	148	PRO	CA-C-N	5.56	126.79	119.84
1	A	148	PRO	C-N-CA	5.56	126.79	119.84
1	A	351	GLU	CB-CG-CD	-5.39	103.44	112.60
1	A	652	ASP	CA-C-O	-5.29	115.46	121.39
1	A	593	LEU	CA-C-N	5.10	128.79	121.91
1	A	593	LEU	C-N-CA	5.10	128.79	121.91

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	204	ARG	Sidechain
1	A	320	ARG	Sidechain
1	A	411	ARG	Sidechain
1	A	414	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5739	0	5616	28	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
3	D	50	0	43	0	0
4	A	56	0	52	3	0
5	A	2	0	0	0	0
6	A	1	0	0	0	0
7	A	2	0	0	0	0
8	A	11	0	17	1	0
9	A	30	0	0	1	0
10	A	395	0	0	4	0
All	All	6342	0	5778	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:207:LYS:HD3	1:A:546:PHE:HB3	1.78	0.64
1:A:195:ASN:HD22	4:A:805:NAG:H83	1.65	0.61
1:A:286:ALA:HB3	1:A:289[B]:LEU:HD22	1.82	0.61
1:A:412:PRO:HA	1:A:589[B]:ASN:HD21	1.66	0.58
1:A:526[A]:GLN:OE1	1:A:692:TYR:O	2.22	0.57
1:A:652:ASP:O	1:A:653:PHE:HB3	2.07	0.55
1:A:659[B]:ILE:O	1:A:663[B]:MET:HG3	2.06	0.55
1:A:98:SER:O	1:A:102[B]:GLU:HG3	2.10	0.52
1:A:656[A]:SER:N	10:A:913:HOH:O	2.44	0.50
1:A:246:TRP:CD1	4:A:806:NAG:H83	2.49	0.47
1:A:131:ILE:HD11	1:A:171[B]:GLU:HG3	1.97	0.46
1:A:169:MET:HE3	1:A:347:HIS:NE2	2.31	0.46
1:A:65:LEU:HD23	1:A:401[B]:SER:OG	2.15	0.46
1:A:56:HIS:CD2	1:A:405[B]:LEU:HD22	2.52	0.44
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.53	0.44
1:A:131:ILE:HG22	1:A:137:GLU:HG2	2.00	0.44
1:A:664:MET:O	1:A:668[A]:LEU:HD22	2.18	0.43
1:A:412:PRO:HA	1:A:589[A]:ASN:OD1	2.17	0.43
8:A:817:TAM:C6	8:A:817:TAM:C4	2.97	0.43
1:A:155:SER:O	1:A:156:ASP:HB2	2.18	0.43
9:A:818:G8W:OBC	9:A:818:G8W:CAF	2.67	0.42
1:A:131:ILE:CG2	1:A:137:GLU:HG2	2.49	0.42
1:A:514[B]:LYS:HB3	1:A:514[B]:LYS:HE3	1.89	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:242:TYR:CG	1:A:243:PRO:HA	2.55	0.41
1:A:308:LYS:NZ	10:A:921:HOH:O	2.53	0.41
1:A:514[A]:LYS:HG3	10:A:1249:HOH:O	2.21	0.41
1:A:178:ASN:HB2	1:A:319:TRP:CE3	2.56	0.41
1:A:207:LYS:HE2	1:A:546:PHE:CG	2.55	0.41
1:A:542:GLU:OE1	1:A:542:GLU:CA	2.69	0.41
4:A:803:NAG:H83	10:A:943:HOH:O	2.19	0.41
1:A:579:VAL:HG12	1:A:583[B]:MET:HE2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	736/707 (104%)	715 (97%)	19 (3%)	2 (0%)	36	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	543	THR
1	A	382	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	638/603 (106%)	618 (97%)	20 (3%)	35	39

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	55	LYS
1	A	122	LYS
1	A	187	LYS
1	A	189	GLU
1	A	239	VAL
1	A	303	GLN
1	A	537	TYR
1	A	542	GLU
1	A	545	LYS
1	A	557	GLU
1	A	589[A]	ASN
1	A	589[B]	ASN
1	A	600	TYR
1	A	652	ASP
1	A	656[A]	SER
1	A	656[B]	SER
1	A	663[A]	MET
1	A	663[B]	MET
1	A	668[A]	LEU
1	A	668[B]	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	136	ASN
1	A	212	ASN
1	A	350	ASN
1	A	451	ASN
1	A	540	ASN
1	A	578	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2,1	14,14,15	0.28	0	17,19,21	0.80	0
2	NAG	B	2	2	14,14,15	0.48	0	17,19,21	1.84	4 (23%)
2	NAG	C	1	2,1	14,14,15	0.74	0	17,19,21	1.47	5 (29%)
2	NAG	C	2	2	14,14,15	0.73	0	17,19,21	1.94	6 (35%)
3	NAG	D	1	1,3	14,14,15	0.88	1 (7%)	17,19,21	1.54	1 (5%)
3	NAG	D	2	3	14,14,15	0.41	0	17,19,21	1.04	1 (5%)
3	BMA	D	3	3	11,11,12	0.63	0	15,15,17	1.10	1 (6%)
3	MAN	D	4	3	11,11,12	0.54	0	15,15,17	1.43	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	4/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	2/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	1	NAG	C1-C2	2.17	1.55	1.52

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	C1-O5-C5	5.46	119.50	112.19
2	B	2	NAG	C2-N2-C7	4.11	128.41	122.90
2	B	2	NAG	C8-C7-N2	3.78	122.38	116.12
2	C	2	NAG	C8-C7-N2	3.71	122.28	116.12
2	C	2	NAG	C3-C4-C5	-3.65	103.62	110.23
2	C	2	NAG	O4-C4-C5	3.09	116.93	109.32
2	B	2	NAG	O7-C7-N2	-3.07	116.56	121.98
3	D	4	MAN	C3-C4-C5	-3.00	104.80	110.23
2	C	1	NAG	O5-C1-C2	-2.90	106.80	111.29
2	C	2	NAG	O7-C7-N2	-2.79	117.06	121.98
2	B	2	NAG	C4-C3-C2	2.58	114.80	111.02
3	D	4	MAN	C1-C2-C3	2.55	113.35	109.64
2	C	2	NAG	O5-C1-C2	-2.49	107.44	111.29
3	D	4	MAN	C1-O5-C5	2.39	115.39	112.19
3	D	2	NAG	O5-C1-C2	-2.23	107.83	111.29
2	C	1	NAG	C2-N2-C7	2.21	125.87	122.90
2	C	1	NAG	C3-C4-C5	-2.14	106.36	110.23
2	C	2	NAG	C1-O5-C5	2.14	115.05	112.19
3	D	3	BMA	C3-C4-C5	2.12	114.07	110.23
2	C	1	NAG	C1-O5-C5	2.03	114.91	112.19
2	C	1	NAG	O4-C4-C5	2.03	114.31	109.32

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
3	D	2	NAG	C8-C7-N2-C2
3	D	2	NAG	O7-C7-N2-C2
3	D	1	NAG	C4-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	D	4	MAN	C4-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
3	D	4	MAN	O5-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 5 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	TAM	A	817	-	10,10,10	0.72	0	12,12,12	0.94	0
9	G8W	A	818	5	30,30,30	1.80	6 (20%)	39,39,39	3.15	12 (30%)
4	NAG	A	805	1	14,14,15	0.81	0	17,19,21	2.37	5 (29%)
4	NAG	A	803	1	14,14,15	0.88	0	17,19,21	2.30	3 (17%)
4	NAG	A	806	1	14,14,15	0.66	0	17,19,21	1.59	3 (17%)
4	NAG	A	804	1	14,14,15	1.05	0	17,19,21	1.25	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	TAM	A	817	-	-	9/12/12/12	-
9	G8W	A	818	5	-	0/31/31/31	0/1/1/1
4	NAG	A	805	1	-	2/6/23/26	0/1/1/1
4	NAG	A	803	1	-	3/6/23/26	0/1/1/1
4	NAG	A	806	1	-	2/6/23/26	0/1/1/1
4	NAG	A	804	1	-	2/6/23/26	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	818	G8W	CAE-CAG	4.50	1.42	1.37
9	A	818	G8W	FAI-CAG	3.86	1.48	1.35
9	A	818	G8W	CAF-NAH	3.67	1.41	1.34
9	A	818	G8W	CAD-NAJ	-2.73	1.36	1.41
9	A	818	G8W	CAX-CAT	2.11	1.58	1.53
9	A	818	G8W	CAG-NAH	2.01	1.36	1.31

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	818	G8W	CAE-CAG-NAH	-12.00	120.56	126.86
9	A	818	G8W	CAF-NAH-CAG	11.21	119.35	115.78
4	A	803	NAG	C1-O5-C5	7.48	122.21	112.19
4	A	805	NAG	C1-O5-C5	7.14	121.75	112.19
9	A	818	G8W	CAC-CAE-CAG	5.86	120.24	116.24
4	A	806	NAG	C1-O5-C5	3.55	116.94	112.19
4	A	805	NAG	C8-C7-N2	3.17	121.37	116.12
9	A	818	G8W	FAI-CAG-CAE	3.11	121.48	118.66
4	A	803	NAG	C8-C7-N2	3.07	121.21	116.12
4	A	806	NAG	O5-C1-C2	-3.06	106.56	111.29
9	A	818	G8W	CAD-CAF-NAH	-3.03	121.57	124.07
9	A	818	G8W	CAD-NAJ-CAB	-2.95	122.30	127.52
4	A	805	NAG	O5-C1-C2	-2.88	106.83	111.29
9	A	818	G8W	CAU-CAT-NAS	-2.77	104.15	110.57
9	A	818	G8W	OBC-CAB-CAK	-2.76	117.02	122.02
9	A	818	G8W	CAL-CB-CA	-2.67	105.59	113.80
4	A	804	NAG	C1-O5-C5	2.57	115.63	112.19
9	A	818	G8W	CB-CA-N	-2.55	105.86	110.91
4	A	805	NAG	C4-C3-C2	-2.40	107.50	111.02
4	A	805	NAG	C6-C5-C4	-2.40	107.13	113.02
9	A	818	G8W	CAK-CAB-NAJ	2.35	118.67	114.59
4	A	806	NAG	O3-C3-C4	-2.32	104.91	110.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	804	NAG	C4-C3-C2	2.28	114.36	111.02
4	A	803	NAG	O6-C6-C5	2.07	118.39	111.33
9	A	818	G8W	FAI-CAG-NAH	2.00	117.95	114.93

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	A	817	TAM	N-C-C1-C4
8	A	817	TAM	N-C-C3-C6
8	A	817	TAM	C-C1-C4-O4
8	A	817	TAM	C-C2-C5-O5
8	A	817	TAM	C-C3-C6-O6
4	A	806	NAG	C4-C5-C6-O6
4	A	803	NAG	C8-C7-N2-C2
4	A	803	NAG	O7-C7-N2-C2
4	A	805	NAG	C8-C7-N2-C2
4	A	805	NAG	O7-C7-N2-C2
4	A	806	NAG	O5-C5-C6-O6
4	A	804	NAG	C4-C5-C6-O6
8	A	817	TAM	C3-C-C1-C4
4	A	803	NAG	C4-C5-C6-O6
4	A	804	NAG	O5-C5-C6-O6
8	A	817	TAM	C2-C-C3-C6
8	A	817	TAM	C2-C-C1-C4
8	A	817	TAM	C1-C-C3-C6


There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	A	817	TAM	1	0
9	A	818	G8W	1	0
4	A	805	NAG	1	0
4	A	803	NAG	1	0
4	A	806	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/707 (98%)	0.29	19 (2%) 56 59	23, 53, 76, 133	47 (6%)

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	546	PHE	5.8
1	A	656[A]	SER	5.4
1	A	653	PHE	3.9
1	A	543	THR	3.5
1	A	279	TYR	3.4
1	A	544	ASN	3.3
1	A	541	TRP	3.3
1	A	548	GLY	3.1
1	A	545	LYS	2.7
1	A	719	VAL	2.4
1	A	152	GLU	2.4
1	A	155	SER	2.3
1	A	547	SER	2.3
1	A	131	ILE	2.2
1	A	657[A]	ASN	2.2
1	A	94[A]	LYS	2.2
1	A	56	HIS	2.2
1	A	506	PHE	2.2
1	A	55	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	C	2	14/15	0.77	0.14	64,75,84,88	0
2	NAG	B	2	14/15	0.80	0.14	70,83,100,104	0
3	MAN	D	4	11/12	0.87	0.13	75,83,87,89	0
3	BMA	D	3	11/12	0.88	0.12	72,76,77,78	0
3	NAG	D	2	14/15	0.89	0.12	67,75,83,89	0
3	NAG	D	1	14/15	0.93	0.10	50,54,70,78	0
2	NAG	B	1	14/15	0.94	0.10	61,73,77,85	0
2	NAG	C	1	14/15	0.95	0.09	58,61,63,72	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NAG	A	805	14/15	0.60	0.19	92,101,104,104	0
4	NAG	A	803	14/15	0.64	0.21	75,90,116,121	0
8	TAM	A	817	11/11	0.79	0.19	91,99,108,111	0
4	NAG	A	804	14/15	0.81	0.14	65,72,77,81	0
4	NAG	A	806	14/15	0.91	0.11	59,73,83,84	0
9	G8W	A	818	30/30	0.96	0.10	43,49,78,89	0
7	CL	A	819	1/1	0.98	0.09	62,62,62,62	0
7	CL	A	816	1/1	0.99	0.04	48,48,48,48	0
5	ZN	A	814	1/1	1.00	0.01	45,45,45,45	0
6	CA	A	815	1/1	1.00	0.02	44,44,44,44	0
5	ZN	A	813	1/1	1.00	0.01	46,46,46,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.