



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 01:22 PM EDT

PDB ID : 6HKZ / pdb_00006hkz
Title : X-ray structure of human glutamate carboxypeptidase II (GCPII) in complex with a inhibitor RNA 2-49-1
Authors : Motlova, L.; Novakova, Z.; Barinka, C.
Deposited on : 2018-09-09
Resolution : 2.09 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	707	3% 89% 8% ..
2	B	2	50% 50%
2	C	2	100%
3	D	4	100%

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 6205 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate carboxypeptidase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	694	Total	C	N	O	S	0	38	0
			5675	3643	951	1060	21			

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	2	Total	C	N	O	0	0	0
			28	16	2	10			
2	C	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 3 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	4	Total	C	N	O	0	0	0
			50	28	2	20			

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Zn	0	0
			2	2		

- Molecule 5 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Cl	0	0
			1	1		

- Molecule 7 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: C₈H₁₅NO₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		
7	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 8 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 9 is (2 {S})-2-[[[(2 {S})-6-[2-[(6-fluoranylpyridin-3-yl)carbonylamino]ethyl-phenyl-amino]-1-oxidanyl-1,6-bis(oxidanylidene)hexan-2-yl]carbamoylamino]pentanedioic acid (CCD ID: GBZ) (formula: C₂₆H₃₀FN₅O₉).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
9	A	1	Total	C	F	N	O	0	0
			41	26	1	5	9		


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	319	Total 319	O 319	0	1

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate carboxypeptidase 2




- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: alpha-D-mannopyranose-(1-3)-beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	102.96Å 132.26Å 161.59Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.59 – 2.09 47.59 – 2.09	Depositor EDS
% Data completeness (in resolution range)	99.2 (47.59-2.09) 99.6 (47.59-2.09)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.30 (at 2.08Å)	Xtriage
Refinement program	REFMAC 5.8.0222	Depositor
R, R_{free}	0.170 , 0.203 0.178 , 0.212	Depositor DCC
R_{free} test set	2100 reflections (3.21%)	wwPDB-VP
Wilson B-factor (Å ²)	33.8	Xtriage
Anisotropy	0.576	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 58.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6205	wwPDB-VP
Average B, all atoms (Å ²)	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, ZN, CL, NAG, MAN, BMA, CA, GBZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.15	16/5988 (0.3%)	1.07	12/8109 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	235	PHE	N-CA	7.31	1.55	1.46
1	A	523	VAL	N-CA	6.37	1.54	1.46
1	A	705	PHE	N-CA	6.19	1.50	1.46
1	A	236	ALA	N-CA	6.12	1.53	1.46
1	A	65	LEU	N-CA	5.78	1.53	1.46
1	A	233	ASP	CA-C	5.52	1.58	1.53
1	A	383	PHE	N-CA	5.49	1.53	1.46
1	A	652	ASP	N-CA	5.48	1.53	1.46
1	A	207[A]	LYS	N-CA	5.26	1.52	1.46
1	A	207[B]	LYS	N-CA	5.26	1.52	1.46
1	A	513	SER	N-CA	5.18	1.52	1.45
1	A	544	ASN	N-CA	5.16	1.52	1.46
1	A	537	TYR	N-CA	5.14	1.52	1.46
1	A	414	ARG	N-CA	5.09	1.51	1.45
1	A	234	TYR	N-CA	5.07	1.52	1.46
1	A	640	THR	N-CA	5.03	1.52	1.46

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	236	ALA	CA-C-N	6.23	127.62	119.84
1	A	236	ALA	C-N-CA	6.23	127.62	119.84
1	A	146	PRO	CA-C-N	-6.20	115.19	119.66
1	A	146	PRO	C-N-CA	-6.20	115.19	119.66
1	A	652	ASP	N-CA-C	5.92	118.47	107.99
1	A	548	GLY	CA-C-O	-5.87	116.51	122.56
1	A	240	LYS	CA-C-N	5.50	128.91	120.82
1	A	240	LYS	C-N-CA	5.50	128.91	120.82
1	A	124	HIS	CA-C-N	5.34	125.28	119.78
1	A	124	HIS	C-N-CA	5.34	125.28	119.78
1	A	295	HIS	CA-C-N	5.14	126.26	119.84
1	A	295	HIS	C-N-CA	5.14	126.26	119.84

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	411	ARG	Sidechain
1	A	414	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5675	0	5546	20	0
2	B	28	0	25	0	0
2	C	28	0	25	0	0
3	D	50	0	43	0	0
4	A	2	0	0	0	0
5	A	1	0	0	0	0
6	A	1	0	0	0	0
7	A	56	0	52	0	0
8	A	4	0	6	0	0
9	A	41	0	0	0	0
10	A	319	0	0	0	0
All	All	6205	0	5697	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:412:PRO:HA	1:A:589:ASN:HD21	1.37	0.90
1:A:235:PHE:HA	1:A:247:ASN:OD1	1.93	0.69
1:A:652:ASP:O	1:A:653:PHE:HB3	1.96	0.64
1:A:412:PRO:HA	1:A:589:ASN:ND2	2.11	0.62
1:A:236:ALA:HB1	1:A:237:PRO:CD	2.31	0.60
1:A:236:ALA:HB1	1:A:237:PRO:HD2	1.85	0.58
1:A:169:MET:HE3	1:A:347:HIS:NE2	2.21	0.54
1:A:526[A]:GLN:OE1	1:A:692:TYR:O	2.27	0.52
1:A:659[B]:ILE:O	1:A:663[B]:MET:HG3	2.10	0.52
1:A:540:ASN:OD1	1:A:542:GLU:HG2	2.14	0.48
1:A:612:TYR:CZ	1:A:616:MET:HG3	2.48	0.47
1:A:652:ASP:N	1:A:652:ASP:OD1	2.47	0.47
1:A:486:GLU:CD	1:A:486:GLU:H	2.24	0.45
1:A:231:PRO:HB3	1:A:235:PHE:HB3	2.00	0.44
1:A:542:GLU:O	1:A:543:THR:OG1	2.34	0.43
1:A:506:PHE:HB2	1:A:509:MET:HG3	2.00	0.42
1:A:325:VAL:HB	1:A:326:PRO:HD2	2.02	0.42
1:A:90:PHE:CE2	1:A:94:LYS:HE2	2.55	0.41
1:A:278:ALA:HB3	1:A:280[B]:ARG:NH1	2.35	0.41
1:A:326:PRO:O	1:A:327:TYR:HB2	2.21	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	729/707 (103%)	708 (97%)	20 (3%)	1 (0%)	48 50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	382	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	631/603 (105%)	612 (97%)	19 (3%)	36 41

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	94	LYS
1	A	102	GLU
1	A	122	LYS
1	A	133	GLU
1	A	187	LYS
1	A	239[A]	VAL
1	A	239[B]	VAL
1	A	303	GLN
1	A	322	SER
1	A	396	HIS
1	A	465	ASP
1	A	486	GLU
1	A	537	TYR
1	A	541	TRP
1	A	557	GLU
1	A	589	ASN
1	A	600	TYR
1	A	652	ASP
1	A	719	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	GLN
1	A	303	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	589	ASN
1	A	618	HIS
1	A	620	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	1,2	14,14,15	0.40	0	17,19,21	0.84	0
2	NAG	B	2	2	14,14,15	0.41	0	17,19,21	1.53	3 (17%)
2	NAG	C	1	1,2	14,14,15	0.52	0	17,19,21	1.26	1 (5%)
2	NAG	C	2	2	14,14,15	0.48	0	17,19,21	1.27	2 (11%)
3	NAG	D	1	3,1	14,14,15	0.56	0	17,19,21	1.61	2 (11%)
3	NAG	D	2	3	14,14,15	0.46	0	17,19,21	1.43	3 (17%)
3	BMA	D	3	3	11,11,12	0.55	0	15,15,17	1.21	1 (6%)
3	MAN	D	4	3	11,11,12	0.43	0	15,15,17	1.11	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1
2	NAG	C	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
3	NAG	D	1	3,1	-	2/6/23/26	0/1/1/1
3	NAG	D	2	3	-	4/6/23/26	0/1/1/1
3	BMA	D	3	3	-	0/2/19/22	0/1/1/1
3	MAN	D	4	3	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	1	NAG	C1-O5-C5	5.26	119.23	112.19
2	B	2	NAG	C2-N2-C7	3.51	127.61	122.90
3	D	2	NAG	C1-O5-C5	3.27	116.58	112.19
3	D	3	BMA	C1-C2-C3	3.26	114.39	109.64
2	B	2	NAG	C4-C3-C2	2.88	115.24	111.02
2	C	2	NAG	C8-C7-N2	2.77	120.72	116.12
2	C	1	NAG	C1-O5-C5	2.70	115.80	112.19
3	D	1	NAG	O5-C1-C2	-2.56	107.33	111.29
2	B	2	NAG	C8-C7-N2	2.38	120.07	116.12
3	D	2	NAG	C8-C7-N2	2.30	119.94	116.12
2	C	2	NAG	O4-C4-C5	2.26	114.88	109.32
3	D	2	NAG	C1-C2-N2	2.18	113.86	110.43
3	D	4	MAN	O5-C5-C6	2.06	111.67	107.66

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	2	NAG	C4-C5-C6-O6
2	B	2	NAG	C8-C7-N2-C2
2	B	2	NAG	O7-C7-N2-C2
2	C	1	NAG	C8-C7-N2-C2
2	C	1	NAG	O7-C7-N2-C2
2	C	2	NAG	C8-C7-N2-C2
2	C	2	NAG	O7-C7-N2-C2
3	D	2	NAG	C8-C7-N2-C2
3	D	2	NAG	O7-C7-N2-C2
3	D	2	NAG	O5-C5-C6-O6

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	D	1	NAG	C4-C5-C6-O6
3	D	1	NAG	O5-C5-C6-O6


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 4 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
7	NAG	A	807	1	14,14,15	0.62	0	17,19,21	1.86	4 (23%)
8	EDO	A	817	-	3,3,3	0.44	0	2,2,2	0.41	0
9	GBZ	A	818	4	42,42,42	1.64	7 (16%)	55,55,55	2.97	12 (21%)
7	NAG	A	808	1	14,14,15	1.01	1 (7%)	17,19,21	1.32	2 (11%)
7	NAG	A	809	1	14,14,15	0.74	0	17,19,21	2.26	6 (35%)
7	NAG	A	810	1	14,14,15	0.47	0	17,19,21	1.41	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	NAG	A	807	1	-	2/6/23/26	0/1/1/1
8	EDO	A	817	-	-	1/1/1/1	-
9	GBZ	A	818	4	-	0/45/45/45	0/2/2/2
7	NAG	A	808	1	-	0/6/23/26	0/1/1/1
7	NAG	A	809	1	-	2/6/23/26	0/1/1/1
7	NAG	A	810	1	-	2/6/23/26	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
9	A	818	GBZ	CAE-CAH	-5.03	1.39	1.50
9	A	818	GBZ	CAF-NAG	4.08	1.42	1.34
9	A	818	GBZ	CBJ-NAM	-3.50	1.36	1.43
9	A	818	GBZ	CAO-CAN	3.10	1.57	1.51
9	A	818	GBZ	CAC-CAA	2.32	1.39	1.37
9	A	818	GBZ	CAA-NAG	2.24	1.36	1.31
7	A	808	NAG	C1-C2	2.23	1.55	1.52
9	A	818	GBZ	CAY-CAW	2.13	1.58	1.53

All (26) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	A	818	GBZ	CAF-NAG-CAA	14.81	120.50	115.78
9	A	818	GBZ	CAC-CAA-NAG	-10.86	121.16	126.86
7	A	809	NAG	C1-O5-C5	5.99	120.21	112.19
7	A	807	NAG	C1-O5-C5	5.24	119.22	112.19
9	A	818	GBZ	CAD-CAC-CAA	4.83	119.53	116.24
9	A	818	GBZ	FAB-CAA-NAG	4.62	121.89	114.93
7	A	810	NAG	C1-O5-C5	3.90	117.41	112.19
9	A	818	GBZ	CAE-CAF-NAG	-3.49	118.46	123.59
7	A	809	NAG	O5-C1-C2	-3.40	106.04	111.29
7	A	809	NAG	C6-C5-C4	-3.16	105.26	113.02
7	A	808	NAG	C1-O5-C5	2.81	115.95	112.19
7	A	807	NAG	C8-C7-N2	2.70	120.60	116.12
9	A	818	GBZ	CAL-NAM-CBJ	2.68	121.94	117.16
9	A	818	GBZ	CA-N-CAU	2.67	126.75	120.64
9	A	818	GBZ	CAX-CAW-NAV	-2.65	104.42	110.57
7	A	810	NAG	C4-C3-C2	-2.55	107.28	111.02
9	A	818	GBZ	CB-CAP-CAO	-2.36	106.97	113.26
7	A	809	NAG	C8-C7-N2	2.36	120.03	116.12
9	A	818	GBZ	OAK-CAH-CAE	-2.33	116.29	120.90
9	A	818	GBZ	CAD-CAE-CAF	2.22	120.09	117.61

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	809	NAG	O5-C5-C6	2.21	111.96	107.66
7	A	808	NAG	C2-N2-C7	2.19	125.83	122.90
7	A	807	NAG	O5-C1-C2	-2.17	107.94	111.29
9	A	818	GBZ	CBL-CBJ-NAM	2.15	123.27	120.18
7	A	807	NAG	O4-C4-C5	2.07	114.42	109.32
7	A	809	NAG	C3-C4-C5	2.06	113.96	110.23

There are no chirality outliers.


All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	A	810	NAG	C4-C5-C6-O6
7	A	810	NAG	O5-C5-C6-O6
7	A	807	NAG	C8-C7-N2-C2
7	A	807	NAG	O7-C7-N2-C2
7	A	809	NAG	C8-C7-N2-C2
7	A	809	NAG	O7-C7-N2-C2
8	A	817	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	694/707 (98%)	0.36	20 (2%) 53 56	25, 55, 82, 110	38 (5%)

All (20) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	234	TYR	7.0
1	A	656[A]	SER	6.3
1	A	235	PHE	5.9
1	A	236	ALA	4.4
1	A	541	TRP	4.3
1	A	653	PHE	3.4
1	A	279	TYR	3.3
1	A	546	PHE	3.0
1	A	543	THR	2.6
1	A	657[A]	ASN	2.5
1	A	131	ILE	2.3
1	A	155	SER	2.3
1	A	55	LYS	2.3
1	A	124	HIS	2.2
1	A	152	GLU	2.2
1	A	545	LYS	2.2
1	A	547	SER	2.1
1	A	733	TYR	2.1
1	A	719	VAL	2.1
1	A	238	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	2	14/15	0.78	0.15	71,87,97,101	0
3	NAG	D	2	14/15	0.82	0.15	72,78,91,95	0
3	MAN	D	4	11/12	0.82	0.14	88,93,98,98	0
2	NAG	C	2	14/15	0.83	0.13	63,71,81,82	0
3	BMA	D	3	11/12	0.89	0.11	76,79,86,87	0
2	NAG	C	1	14/15	0.92	0.12	57,60,63,67	0
3	NAG	D	1	14/15	0.94	0.09	50,57,72,74	0
2	NAG	B	1	14/15	0.94	0.10	60,67,74,84	0


The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
7	NAG	A	809	14/15	0.68	0.17	89,93,100,101	0
7	NAG	A	807	14/15	0.69	0.19	73,84,111,116	0
7	NAG	A	808	14/15	0.78	0.14	67,73,77,79	0
7	NAG	A	810	14/15	0.88	0.13	64,83,98,100	0
8	EDO	A	817	4/4	0.92	0.13	72,74,77,82	0
9	GBZ	A	818	41/41	0.95	0.13	43,54,125,132	0
5	CA	A	803	1/1	1.00	0.02	44,44,44,44	0
6	CL	A	804	1/1	1.00	0.06	49,49,49,49	0
4	ZN	A	801	1/1	1.00	0.01	48,48,48,48	0
4	ZN	A	802	1/1	1.00	0.02	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.