



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:03 PM UTC

PDB ID : 7HKC / pdb_00007hkc
Title : Group deposition for crystallographic fragment screening of the NS5 RNA-dependent RNA polymerase from Dengue virus serotype 2 – Crystal structure of the NS5 RNA-dependent RNA polymerase from Dengue virus serotype 2 in complex with Z4628744292 (DNV2_NS5A-x0160)
Authors : Saini, M.; Chopra, A.; Aschenbrenner, J.C.; Marples, P.G.; Balcomb, B.H.; Fearon, D.; von Delft, F.; Ruiz, F.X.; Arnold, E.
Deposited on : 2024-10-15
Resolution : 1.74 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	: 4-5-2 with Phenix2.0
Mogul	: 2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	: 2.0
EDS	: 3.0
Buster-report	: wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	: 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	: 9.0.010 (Gargrove)
Density-Fitness	: 1.0.12
Ideal geometry (proteins)	: Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.74 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1187 (1.74-1.74)
Clashscore	190562	1207 (1.74-1.74)
Ramachandran outliers	187476	1200 (1.74-1.74)
Sidechain outliers	187428	1200 (1.74-1.74)
RSRZ outliers	180081	1188 (1.74-1.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	637	18% 79% 11% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PO4	A	1008	-	-	X	-

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 5405 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	574	Total	C	N	O	S	0	8	0
			4890	3069	877	910	34			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	264	GLY	-	expression tag	UNP Q91H74
A	265	PRO	-	expression tag	UNP Q91H74

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Zn	0	0
			2	2		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	1
			24	12	2	8	2		

- Molecule 5 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C₂H₆OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		
5	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		
6	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $\text{C}_4\text{H}_{10}\text{O}_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 8 is 4-(4-methyl-1H-pyrazol-5-yl)pyridine (CCD ID: A1AP7) (formula: C₉H₉N₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	N	0	0
			12	9	3		
8	A	1	Total	C	N	0	1
			24	18	6		
8	A	1	Total	C	N	0	0
			12	9	3		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	404	Total	O	0	0
			404	404		

4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	82.57Å 116.51Å 147.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	67.46 – 1.74 67.46 – 1.74	Depositor EDS
% Data completeness (in resolution range)	98.3 (67.46-1.74) 98.4 (67.46-1.74)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 1.74Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC5	Depositor
R, R_{free}	0.189 , 0.227 0.202 , 0.234	Depositor DCC
R_{free} test set	3755 reflections (5.13%)	wwPDB-VP
Wilson B-factor (Å ²)	31.7	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5405	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CL, PO4, PEG, DMS, ZN, MES, A1AP7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	2/4995 (0.0%)	1.31	6/6732 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	650	VAL	N-CA	5.81	1.51	1.46
1	A	768	HIS	CE1-NE2	5.30	1.37	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	702	ASP	CA-CB-CG	5.99	118.59	112.60
1	A	559	GLY	CA-C-N	5.65	128.17	120.54
1	A	559	GLY	C-N-CA	5.65	128.17	120.54
1	A	479	LEU	CA-C-N	5.42	125.96	119.94
1	A	479	LEU	C-N-CA	5.42	125.96	119.94
1	A	650	VAL	CA-C-O	-5.03	115.54	119.46

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	273	ASN	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4890	0	4773	43	0
2	A	1	0	0	0	0
3	A	2	0	0	0	0
4	A	24	0	26	1	0
5	A	12	0	18	4	0
6	A	10	0	0	4	0
7	A	14	0	20	0	0
8	A	48	0	0	0	0
9	A	404	0	0	10	1
All	All	5405	0	4837	47	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:649:ARG:HH11	1:A:649:ARG:HG2	1.01	1.10
1:A:649:ARG:HG2	1:A:649:ARG:NH1	1.74	0.97
1:A:649:ARG:HH11	1:A:649:ARG:CG	1.75	0.97
1:A:664:ASP:OD1	6:A:1008:PO4:O4	1.90	0.90
5:A:1005:DMS:H11	9:A:1304:HOH:O	1.90	0.71
1:A:534:ASP:OD1	6:A:1008:PO4:O4	2.10	0.70
1:A:664:ASP:OD1	6:A:1008:PO4:P	2.50	0.69
1:A:406:ASN:OD1	1:A:423:ARG:NH1	2.36	0.58
5:A:1005:DMS:H12	9:A:1362:HOH:O	2.04	0.58
1:A:638:THR:O	1:A:641:ILE:HG22	2.04	0.57
1:A:886:MET:HE1	9:A:1224:HOH:O	2.07	0.54
1:A:403:VAL:HG21	1:A:426:VAL:HG21	1.90	0.54
1:A:764:LEU:O	1:A:765:MET:HE2	2.09	0.53
1:A:534:ASP:OD1	6:A:1008:PO4:P	2.66	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:649:ARG:NH1	1:A:649:ARG:CG	2.47	0.52
1:A:513:HIS:HB3	1:A:765:MET:HE1	1.92	0.51
1:A:528:GLY:O	1:A:668:LYS:HE3	2.11	0.51
1:A:447:CYS:SG	1:A:450:CYS:HB2	2.51	0.50
1:A:841:LYS:NZ	9:A:1115:HOH:O	2.45	0.48
1:A:475:TRP:HZ3	1:A:576:LYS:HD3	1.79	0.48
1:A:638:THR:HA	1:A:641:ILE:HG22	1.96	0.47
1:A:312:LYS:O	1:A:313:GLN:CG	2.62	0.47
1:A:303:TRP:CE3	1:A:593:ILE:HD12	2.50	0.47
1:A:790:THR:HG22	9:A:1369:HOH:O	2.15	0.46
5:A:1005:DMS:C1	9:A:1304:HOH:O	2.57	0.46
1:A:886:MET:CE	9:A:1224:HOH:O	2.64	0.45
1:A:438:GLU:HG3	1:A:449:THR:OG1	2.17	0.45
1:A:453:ASN:ND2	1:A:579:ARG:HD2	2.32	0.45
1:A:599:ARG:HG2	1:A:606:THR:HG23	1.99	0.44
1:A:305:TYR:OH	1:A:308:SER:OG	2.26	0.44
1:A:303:TRP:CD2	1:A:593:ILE:HD12	2.51	0.44
1:A:562:LYS:NZ	9:A:1124:HOH:O	2.50	0.43
1:A:562:LYS:HE3	1:A:566:GLU:OE2	2.18	0.43
1:A:312:LYS:O	1:A:313:GLN:HG3	2.19	0.42
1:A:452:TYR:O	1:A:578:VAL:HA	2.18	0.42
1:A:337:MET:HE3	9:A:1391:HOH:O	2.19	0.42
1:A:758:TYR:CD1	5:A:1007:DMS:H21	2.55	0.42
1:A:733:GLU:O	1:A:737[C]:ARG:HG3	2.20	0.42
1:A:733:GLU:O	1:A:737[A]:ARG:HG3	2.20	0.42
1:A:400:THR:O	1:A:403:VAL:HG22	2.20	0.41
1:A:834:GLU:OE2	1:A:890:ARG:NE	2.51	0.41
1:A:394:CYS:HB3	1:A:486:PHE:CE2	2.56	0.41
1:A:819:GLN:NE2	9:A:1109:HOH:O	2.42	0.41
4:A:1004[B]:MES:H31	4:A:1004[B]:MES:H81	1.94	0.41
1:A:695:GLU:HA	1:A:696:PRO:HD3	1.91	0.41
1:A:411:ALA:HA	1:A:477:MET:O	2.21	0.40
1:A:765:MET:HE2	1:A:765:MET:HA	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:1139:HOH:O	9:A:1139:HOH:O[2_445]	1.41	0.79

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	584/637 (92%)	560 (96%)	22 (4%)	2 (0%)	36	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	GLN
1	A	418	LYS

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	528/554 (95%)	509 (96%)	19 (4%)	31	9

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	274	LEU
1	A	311	THR
1	A	357	GLU
1	A	358	LYS
1	A	359	VAL
1	A	365	GLU
1	A	375	LYS
1	A	406	ASN
1	A	418	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	420	LYS
1	A	429	SER
1	A	455	MET
1	A	482	ARG
1	A	580	VAL
1	A	601	SER
1	A	633	GLN
1	A	641	ILE
1	A	649	ARG
1	A	830	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	549	ASN
1	A	581	GLN
1	A	603	GLN
1	A	618	GLN
1	A	701	ASN
1	A	786	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 3 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	DMS	A	1007	-	3,3,3	0.13	0	3,3,3	0.25	0
5	DMS	A	1006	-	3,3,3	0.29	0	3,3,3	0.04	0
7	PEG	A	1011	-	6,6,6	0.16	0	5,5,5	0.09	0
6	PO4	A	1008	-	4,4,4	3.93	3 (75%)	6,6,6	0.54	0
8	A1AP7	A	1012	-	13,13,13	0.32	0	16,17,17	0.54	0
8	A1AP7	A	1013[B]	-	13,13,13	0.27	0	16,17,17	0.51	0
4	MES	A	1004[B]	-	12,12,12	0.70	0	15,16,16	0.29	0
5	DMS	A	1005	-	3,3,3	0.52	0	3,3,3	0.45	0
8	A1AP7	A	1014	-	13,13,13	0.23	0	16,17,17	0.51	0
8	A1AP7	A	1013[A]	-	13,13,13	0.26	0	16,17,17	0.47	0
7	PEG	A	1009	-	6,6,6	0.17	0	5,5,5	0.08	0
6	PO4	A	1010	-	4,4,4	1.18	1 (25%)	6,6,6	0.46	0
4	MES	A	1004[A]	-	12,12,12	0.83	0	15,16,16	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	A1AP7	A	1012	-	-	0/4/4/4	0/2/2/2
7	PEG	A	1011	-	-	2/4/4/4	-
8	A1AP7	A	1013[B]	-	-	0/4/4/4	0/2/2/2
4	MES	A	1004[B]	-	-	5/6/14/14	0/1/1/1
8	A1AP7	A	1014	-	-	0/4/4/4	0/2/2/2
8	A1AP7	A	1013[A]	-	-	0/4/4/4	0/2/2/2
7	PEG	A	1009	-	-	1/4/4/4	-
4	MES	A	1004[A]	-	-	0/6/14/14	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1008	PO4	P-O1	6.40	1.65	1.50
6	A	1008	PO4	P-O2	3.55	1.65	1.54
6	A	1008	PO4	P-O3	2.25	1.61	1.54
6	A	1010	PO4	P-O1	2.20	1.55	1.50

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1004[B]	MES	C8-C7-N4-C3
4	A	1004[B]	MES	C7-C8-S-O1S
4	A	1004[B]	MES	C7-C8-S-O3S
7	A	1011	PEG	O2-C3-C4-O4
4	A	1004[B]	MES	C8-C7-N4-C5
4	A	1004[B]	MES	C7-C8-S-O2S
7	A	1011	PEG	C4-C3-O2-C2
7	A	1009	PEG	C4-C3-O2-C2


There are no ring outliers.

4 monomers are involved in 9 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1007	DMS	1	0
6	A	1008	PO4	4	0
4	A	1004[B]	MES	1	0
5	A	1005	DMS	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand A1AP7 A 1012




Ligand A1AP7 A 1013 (B)



Ligand A1AP7 A 1014



Ligand A1AP7 A 1013 (A)



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	574/637 (90%)	1.13	116 (20%) 3 3	3, 38, 103, 152	15 (2%)

All (116) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	512[A]	LEU	14.4
1	A	716	LEU	12.6
1	A	737[A]	ARG	11.8
1	A	763[A]	SER	9.9
1	A	741[A]	SER	9.9
1	A	719	LYS	9.6
1	A	785[A]	SER	9.5
1	A	551[A]	GLU	9.3
1	A	864[A]	GLN	9.1
1	A	801	HIS	8.8
1	A	720	ASP	8.0
1	A	649	ARG	7.2
1	A	809	MET	6.9
1	A	589	VAL	6.7
1	A	413	PHE	5.5
1	A	475	TRP	5.5
1	A	359	VAL	5.3
1	A	426	VAL	5.2
1	A	474	ILE	5.0
1	A	401	ARG	4.9
1	A	363	THR	4.8
1	A	407	ALA	4.7
1	A	419	TRP	4.6
1	A	412	ILE	4.5
1	A	397[A]	GLU	4.5
1	A	317	ALA	4.4
1	A	600	GLY	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	411	ALA	4.4
1	A	476	TYR	4.3
1	A	409	LEU	4.1
1	A	588	THR	4.0
1	A	294	TYR	3.9
1	A	290	THR	3.8
1	A	305	TYR	3.8
1	A	793	THR	3.8
1	A	428	ASP	3.7
1	A	292	TRP	3.7
1	A	584	THR	3.7
1	A	284	ILE	3.6
1	A	431	PHE	3.5
1	A	344	THR	3.5
1	A	880	THR	3.5
1	A	314	THR	3.4
1	A	634	HIS	3.4
1	A	637	VAL	3.4
1	A	311	THR	3.3
1	A	361	THR	3.3
1	A	358	LYS	3.3
1	A	274	LEU	3.2
1	A	449	THR	3.1
1	A	889	PHE	3.1
1	A	281	ILE	3.0
1	A	303	TRP	2.9
1	A	593	ILE	2.9
1	A	410	GLY	2.9
1	A	300	TYR	2.9
1	A	425	ALA	2.9
1	A	452	TYR	2.8
1	A	310	GLU	2.8
1	A	422	ALA	2.8
1	A	745	GLY	2.8
1	A	455	MET	2.8
1	A	746	TRP	2.8
1	A	309	TYR	2.8
1	A	441	LEU	2.8
1	A	635	LEU	2.8
1	A	393	MET	2.7
1	A	641	ILE	2.7
1	A	592	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	717	ILE	2.7
1	A	792	ARG	2.6
1	A	583	PRO	2.6
1	A	273	ASN	2.6
1	A	298	HIS	2.6
1	A	408	ALA	2.6
1	A	478	TRP	2.6
1	A	293	HIS	2.5
1	A	744	ALA	2.5
1	A	577	VAL	2.5
1	A	302	THR	2.5
1	A	528	GLY	2.4
1	A	403	VAL	2.4
1	A	406	ASN	2.4
1	A	443	LEU	2.4
1	A	399	PHE	2.4
1	A	750	GLU	2.3
1	A	288	HIS	2.3
1	A	604	VAL	2.3
1	A	319	SER	2.3
1	A	534	ASP	2.3
1	A	277	ILE	2.3
1	A	434	LEU	2.3
1	A	891	ARG	2.3
1	A	580	VAL	2.3
1	A	650	VAL	2.3
1	A	296	GLN	2.2
1	A	357	GLU	2.2
1	A	648	VAL	2.2
1	A	477	MET	2.2
1	A	571	LEU	2.2
1	A	427	GLU	2.2
1	A	417	ASN	2.2
1	A	304	ALA	2.2
1	A	418	LYS	2.2
1	A	295	ASP	2.1
1	A	564	LEU	2.1
1	A	432	TRP	2.1
1	A	743	GLY	2.1
1	A	748	LEU	2.1
1	A	694	TRP	2.1
1	A	445	GLY	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	435	VAL	2.0
1	A	790	THR	2.0
1	A	424	GLU	2.0
1	A	454	MET	2.0
1	A	275	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	PEG	A	1009	7/7	0.54	0.24	106,116,127,130	0
6	PO4	A	1010	5/5	0.57	0.18	70,70,75,94	0
5	DMS	A	1006	4/4	0.67	0.33	97,112,116,126	0
6	PO4	A	1008	5/5	0.77	0.14	35,38,54,57	0
8	A1AP7	A	1012	12/12	0.80	0.21	28,28,29,29	12
8	A1AP7	A	1013[A]	12/12	0.83	0.21	32,34,37,38	12
8	A1AP7	A	1013[B]	12/12	0.83	0.21	44,47,51,53	12
7	PEG	A	1011	7/7	0.87	0.14	57,69,75,75	0
8	A1AP7	A	1014	12/12	0.90	0.20	46,48,50,50	12
5	DMS	A	1007	4/4	0.92	0.15	57,61,63,66	0
5	DMS	A	1005	4/4	0.93	0.14	46,48,49,50	0
4	MES	A	1004[B]	12/12	0.98	0.31	457,466,488,490	12
4	MES	A	1004[A]	12/12	0.98	0.31	24,27,30,31	12
3	ZN	A	1003	1/1	0.99	0.05	49,49,49,49	0
2	CL	A	1001	1/1	0.99	0.06	41,41,41,41	0
3	ZN	A	1002	1/1	1.00	0.01	22,22,22,22	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A1AP7 A 1012:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1AP7 A 1013 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around A1AP7 A 1013 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around A1AP7 A 1014:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.