



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 10:44 AM UTC

PDB ID : 7HKM / pdb_00007hkm
Title : Group deposition for crystallographic fragment screening of the NS5 RNA-dependent RNA polymerase from Dengue virus serotype 2 – Crystal structure of the NS5 RNA-dependent RNA polymerase from Dengue virus serotype 2 in complex with Z851110644 (DNV2_NS5A-x0288)
Authors : Saini, M.; Chopra, A.; Aschenbrenner, J.C.; Marples, P.G.; Balcomb, B.H.; Fearon, D.; von Delft, F.; Ruiz, F.X.; Arnold, E.
Deposited on : 2024-10-15
Resolution : 1.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1141 (1.64-1.64)
Clashscore	190562	1171 (1.64-1.64)
Ramachandran outliers	187476	1151 (1.64-1.64)
Sidechain outliers	187428	1150 (1.64-1.64)
RSRZ outliers	180081	1141 (1.64-1.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	637	19% 81% 10% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	PEG	A	1009	-	-	-	X
6	PO4	A	1008	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5358 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Genome polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	582	Total	C	N	O	S	0	8	0
			4822	3036	865	887	34			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	264	GLY	-	expression tag	UNP Q91H74
A	265	PRO	-	expression tag	UNP Q91H74

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Zn	0	0
			2	2		

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: C₆H₁₃NO₄S).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	1
			24	12	2	8	2		

- Molecule 4 is DIMETHYL SULFOXIDE (CCD ID: DMS) (formula: C_2H_6OS).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		
4	A	1	Total	C	O	S	0	0
			4	2	1	1		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		
5	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 6 is PHOSPHATE ION (CCD ID: PO4) (formula: O_4P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	O	P	0	0
			5	4	1		

- Molecule 7 is N,2-dimethyl-1,3-thiazole-5-carboxamide (CCD ID: K24) (formula: $C_6H_8N_2OS$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
7	A	1	Total	C	N	O	S	0	0
			10	6	2	1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	457	Total	O	0	0
			457	457		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Genome polypeptide



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	82.44Å 116.79Å 148.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.96 – 1.63 49.96 – 1.63	Depositor EDS
% Data completeness (in resolution range)	96.8 (49.96-1.63) 97.0 (49.96-1.63)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.96 (at 1.63Å)	Xtriage
Refinement program	REFMAC 5.8.0267, REFMAC5	Depositor
R, R_{free}	0.194 , 0.223 0.212 , 0.240	Depositor DCC
R_{free} test set	4552 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	28.6	Xtriage
Anisotropy	0.177	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5358	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DMS, PEG, PO4, K24, MES, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.06	1/4929 (0.0%)	1.32	6/6645 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	768	HIS	CE1-NE2	5.70	1.38	1.32

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	702	ASP	CA-CB-CG	6.28	118.88	112.60
1	A	702	ASP	CB-CA-C	5.45	118.53	110.14
1	A	559	GLY	CA-C-N	5.02	127.32	120.54
1	A	559	GLY	C-N-CA	5.02	127.32	120.54
1	A	322	ASN	CA-C-N	5.02	125.55	119.98
1	A	322	ASN	C-N-CA	5.02	125.55	119.98

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4822	0	4720	38	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	2	0	0	0	0
3	A	24	0	26	0	0
4	A	12	0	18	2	0
5	A	21	0	30	0	0
6	A	10	0	0	4	0
7	A	10	0	0	0	0
8	A	457	0	0	8	1
All	All	5358	0	4794	40	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:1004:DMS:H11	8:A:1321:HOH:O	1.83	0.78
1:A:664:ASP:OD1	6:A:1008:PO4:O4	2.06	0.74
4:A:1004:DMS:H12	8:A:1286:HOH:O	1.92	0.70
1:A:664:ASP:OD1	6:A:1008:PO4:P	2.60	0.59
1:A:733:GLU:O	1:A:737:ARG:HG3	2.06	0.55
1:A:312:LYS:O	1:A:313:GLN:HG2	2.07	0.54
1:A:562:LYS:NZ	8:A:1113:HOH:O	2.40	0.54
1:A:312:LYS:O	1:A:313:GLN:CG	2.56	0.53
1:A:764:LEU:HG	1:A:765:MET:CE	2.39	0.53
1:A:534:ASP:OD1	6:A:1008:PO4:O4	2.29	0.51
1:A:447:CYS:SG	1:A:450:CYS:HB2	2.51	0.50
1:A:834:GLU:OE2	1:A:890:ARG:NE	2.40	0.50
1:A:805:THR:CG2	1:A:809:MET:HE1	2.43	0.49
1:A:886:MET:HE1	8:A:1222:HOH:O	2.14	0.47
1:A:638:THR:HA	1:A:641:ILE:HG22	1.96	0.47
1:A:453:ASN:ND2	1:A:579:ARG:HD2	2.29	0.47
1:A:790:THR:HG22	8:A:1366:HOH:O	2.15	0.47
1:A:801[B]:HIS:H	1:A:801[B]:HIS:CD2	2.31	0.47
1:A:403:VAL:HG21	1:A:426:VAL:HG21	1.97	0.47
1:A:357:GLU:N	1:A:357:GLU:OE2	2.48	0.46
1:A:400:THR:O	1:A:403:VAL:HG22	2.15	0.46
1:A:528:GLY:O	1:A:668:LYS:HE3	2.16	0.46
1:A:638:THR:HA	1:A:641:ILE:CG2	2.46	0.45
1:A:411:ALA:HA	1:A:477:MET:O	2.17	0.45
1:A:579:ARG:HA	1:A:591:ASP:O	2.18	0.44
1:A:474:ILE:HD12	1:A:474:ILE:N	2.32	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:695:GLU:HA	1:A:696:PRO:HD3	1.92	0.44
1:A:305:TYR:OH	1:A:308:SER:OG	2.28	0.43
1:A:886:MET:CE	8:A:1222:HOH:O	2.66	0.43
1:A:321:VAL:HG11	1:A:326:ARG:CZ	2.49	0.42
1:A:764:LEU:C	1:A:765:MET:HE2	2.45	0.42
1:A:394:CYS:HB3	1:A:486:PHE:CE2	2.55	0.42
1:A:303:TRP:CD2	1:A:593:ILE:HD12	2.55	0.42
1:A:475:TRP:CD1	1:A:475:TRP:N	2.88	0.42
1:A:599:ARG:HG2	1:A:606:THR:HG23	2.01	0.41
1:A:356:LYS:HD3	8:A:1523:HOH:O	2.20	0.41
1:A:295:ASP:OD1	1:A:296:GLN:N	2.54	0.40
1:A:384:GLU:OE2	8:A:1102:HOH:O	2.21	0.40
1:A:534:ASP:OD1	6:A:1008:PO4:P	2.79	0.40
1:A:805:THR:HG23	1:A:809:MET:HE1	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
8:A:1167:HOH:O	8:A:1167:HOH:O[2_445]	1.62	0.58

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	576/637 (90%)	554 (96%)	21 (4%)	1 (0%)	43 27

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	473	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	518/554 (94%)	502 (97%)	16 (3%)	35 10

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	274	LEU
1	A	290	THR
1	A	356	LYS
1	A	375	LYS
1	A	426	VAL
1	A	429	SER
1	A	455	MET
1	A	482	ARG
1	A	588	THR
1	A	589	VAL
1	A	631	SER
1	A	635	LEU
1	A	641	ILE
1	A	644	LYS
1	A	693	GLN
1	A	830	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	603	GLN
1	A	618	GLN
1	A	645	ASN
1	A	693	GLN
1	A	701	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 13 ligands modelled in this entry, 2 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PEG	A	1011	-	6,6,6	0.17	0	5,5,5	0.13	0
6	PO4	A	1008	-	4,4,4	4.20	3 (75%)	6,6,6	0.61	0
3	MES	A	1003[B]	-	12,12,12	0.70	0	15,16,16	0.31	0
5	PEG	A	1007	-	6,6,6	0.20	0	5,5,5	0.10	0
6	PO4	A	1010	-	4,4,4	1.19	1 (25%)	6,6,6	0.42	0
4	DMS	A	1004	-	3,3,3	0.65	0	3,3,3	0.45	0
5	PEG	A	1009	-	6,6,6	0.12	0	5,5,5	0.08	0
7	K24	A	1012	-	10,10,10	0.19	0	12,13,13	0.77	1 (8%)
4	DMS	A	1005	-	3,3,3	0.34	0	3,3,3	0.07	0
4	DMS	A	1006	-	3,3,3	0.12	0	3,3,3	0.29	0
3	MES	A	1003[A]	-	12,12,12	0.74	0	15,16,16	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	A	1011	-	-	2/4/4/4	-
3	MES	A	1003[A]	-	-	3/6/14/14	0/1/1/1
3	MES	A	1003[B]	-	-	5/6/14/14	0/1/1/1
5	PEG	A	1009	-	-	3/4/4/4	-
7	K24	A	1012	-	-	2/6/6/6	0/1/1/1
5	PEG	A	1007	-	-	2/4/4/4	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	A	1008	PO4	P-O1	6.97	1.66	1.50
6	A	1008	PO4	P-O2	3.61	1.65	1.54
6	A	1008	PO4	P-O3	2.35	1.61	1.54
6	A	1010	PO4	P-O1	2.26	1.55	1.50

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1012	K24	C3-C2-C1	2.40	134.38	130.64

There are no chirality outliers.

All (17) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1003[A]	MES	C7-C8-S-O2S
3	A	1003[A]	MES	C7-C8-S-O3S
3	A	1003[B]	MES	C8-C7-N4-C3
3	A	1003[B]	MES	C7-C8-S-O2S
3	A	1003[B]	MES	C7-C8-S-O3S
7	A	1012	K24	O-C1-N-C
7	A	1012	K24	C2-C1-N-C
5	A	1011	PEG	O2-C3-C4-O4
5	A	1007	PEG	O2-C3-C4-O4
3	A	1003[B]	MES	C8-C7-N4-C5
3	A	1003[A]	MES	C7-C8-S-O1S
3	A	1003[B]	MES	C7-C8-S-O1S
5	A	1009	PEG	O1-C1-C2-O2
5	A	1009	PEG	O2-C3-C4-O4
5	A	1007	PEG	O1-C1-C2-O2
5	A	1011	PEG	C4-C3-O2-C2
5	A	1009	PEG	C4-C3-O2-C2

There are no ring outliers.

2 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1008	PO4	4	0
4	A	1004	DMS	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	582/637 (91%)	1.29	123 (21%) 2 3	5, 36, 102, 144	25 (4%)

All (123) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	512[A]	LEU	16.2
1	A	515	LEU	13.7
1	A	801[A]	HIS	13.1
1	A	719[A]	LYS	12.8
1	A	713	PHE	12.6
1	A	725	VAL	12.3
1	A	511	GLY	12.2
1	A	800	THR	12.1
1	A	513	HIS	10.7
1	A	646	TRP	10.6
1	A	514	LYS	9.6
1	A	741[A]	SER	9.4
1	A	650	VAL	9.2
1	A	396	ARG	9.2
1	A	401	ARG	9.1
1	A	509	GLY	8.9
1	A	551[A]	GLU	8.6
1	A	785[A]	SER	8.6
1	A	864[A]	GLN	8.5
1	A	763[A]	SER	8.4
1	A	510	GLU	8.0
1	A	705	GLN	8.0
1	A	397	GLU	7.9
1	A	413	PHE	6.3
1	A	799	ALA	6.1
1	A	589	VAL	5.9
1	A	355	PHE	5.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	649	ARG	5.0
1	A	645	ASN	4.9
1	A	475	TRP	4.8
1	A	359	VAL	4.8
1	A	294	TYR	4.6
1	A	411	ALA	4.4
1	A	284	ILE	4.3
1	A	637	VAL	4.3
1	A	600	GLY	4.2
1	A	409	LEU	4.1
1	A	431	PHE	4.1
1	A	412	ILE	4.0
1	A	584	THR	4.0
1	A	309	TYR	3.9
1	A	363	THR	3.9
1	A	889	PHE	3.9
1	A	314	THR	3.9
1	A	419	TRP	3.9
1	A	426	VAL	3.8
1	A	298	HIS	3.7
1	A	290	THR	3.7
1	A	281	ILE	3.6
1	A	274	LEU	3.6
1	A	344	THR	3.6
1	A	794	THR	3.6
1	A	300	TYR	3.5
1	A	311	THR	3.5
1	A	361	THR	3.5
1	A	299	PRO	3.5
1	A	317	ALA	3.4
1	A	292	TRP	3.4
1	A	293	HIS	3.4
1	A	428	ASP	3.3
1	A	407	ALA	3.3
1	A	305	TYR	3.3
1	A	583	PRO	3.3
1	A	473	ALA	3.3
1	A	356	LYS	3.2
1	A	454	MET	3.1
1	A	793	THR	3.1
1	A	577	VAL	3.1
1	A	310	GLU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	588	THR	3.0
1	A	634	HIS	3.0
1	A	403	VAL	3.0
1	A	410	GLY	2.9
1	A	592	ILE	2.9
1	A	430	GLY	2.9
1	A	443	LEU	2.8
1	A	528	GLY	2.8
1	A	635	LEU	2.8
1	A	582	ARG	2.8
1	A	358	LYS	2.8
1	A	474	ILE	2.7
1	A	295	ASP	2.7
1	A	288	HIS	2.6
1	A	417	ASN	2.6
1	A	745	GLY	2.6
1	A	792	ARG	2.6
1	A	304	ALA	2.6
1	A	308	SER	2.6
1	A	744	ALA	2.6
1	A	273	ASN	2.6
1	A	880	THR	2.6
1	A	303	TRP	2.6
1	A	422	ALA	2.5
1	A	590	MET	2.5
1	A	597	ASP	2.5
1	A	890	ARG	2.5
1	A	791	SER	2.5
1	A	591	ASP	2.4
1	A	399	PHE	2.4
1	A	434	LEU	2.4
1	A	441	LEU	2.4
1	A	478	TRP	2.4
1	A	408	ALA	2.4
1	A	275	ASP	2.4
1	A	891	ARG	2.4
1	A	648	VAL	2.4
1	A	406	ASN	2.3
1	A	452	TYR	2.3
1	A	571	LEU	2.2
1	A	593	ILE	2.2
1	A	427	GLU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	277	ILE	2.2
1	A	472	ARG	2.2
1	A	455	MET	2.2
1	A	289	GLU	2.2
1	A	884	PRO	2.1
1	A	476	TYR	2.1
1	A	276	ILE	2.1
1	A	435	VAL	2.0
1	A	432	TRP	2.0
1	A	694	TRP	2.0
1	A	746	TRP	2.0
1	A	596	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	PEG	A	1009	7/7	0.53	0.48	220,224,227,228	7
7	K24	A	1012	10/10	0.59	0.37	64,64,66,67	10
6	PO4	A	1010	5/5	0.60	0.19	77,77,89,103	0
5	PEG	A	1007	7/7	0.62	0.20	79,86,92,94	0
4	DMS	A	1005	4/4	0.69	0.30	70,93,94,108	0
6	PO4	A	1008	5/5	0.74	0.14	36,38,55,57	0
5	PEG	A	1011	7/7	0.81	0.17	60,69,72,74	0
4	DMS	A	1004	4/4	0.92	0.15	40,45,47,47	0
4	DMS	A	1006	4/4	0.95	0.12	53,56,58,59	0
3	MES	A	1003[B]	12/12	0.97	0.35	698,716,740,742	12
3	MES	A	1003[A]	12/12	0.97	0.35	21,24,27,28	12

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	ZN	A	1002	1/1	0.99	0.05	48,48,48,48	0
2	ZN	A	1001	1/1	1.00	0.01	20,20,20,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.