



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 08:04 AM UTC

PDB ID : 8HKN / pdb_00008hkn
Title : Mutated human ADP-ribosyltransferase 2 (PARP2) catalytic domain bound to approved drug Fluzoparib
Authors : Wang, X.Y.; Zhou, J.; Xu, B.L.
Deposited on : 2022-11-27
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	5829 (2.50-2.50)
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)
Sidechain outliers	187428	6380 (2.50-2.50)
RSRZ outliers	180081	5833 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	% 85% 13% ..
1	B	353	% 78% 20% .

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5913 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Poly [ADP-ribose] polymerase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	353	Total	C	N	O	S	0	0	0
			2802	1777	486	520	19			
1	A	353	Total	C	N	O	S	0	0	0
			2802	1777	486	520	19			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	229	GLY	-	expression tag	UNP Q9UGN5
B	349	SER	THR	engineered mutation	UNP Q9UGN5
B	351	ARG	LEU	engineered mutation	UNP Q9UGN5
B	353	GLY	SER	engineered mutation	UNP Q9UGN5
B	354	LEU	PRO	engineered mutation	UNP Q9UGN5
A	229	GLY	-	expression tag	UNP Q9UGN5
A	349	SER	THR	engineered mutation	UNP Q9UGN5
A	351	ARG	LEU	engineered mutation	UNP Q9UGN5
A	353	GLY	SER	engineered mutation	UNP Q9UGN5
A	354	LEU	PRO	engineered mutation	UNP Q9UGN5

- Molecule 2 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is Fluzoparib (CCD ID: 25I) (formula: $C_{22}H_{16}F_4N_6O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	F	N	O	0	0
			34	22	4	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	F	N	O	0	0
			34	22	4	6	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	110	Total	O	0	0
			110	110		
4	A	119	Total	O	0	0
			119	119		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Poly [ADP-ribose] polymerase 2



- Molecule 1: Poly [ADP-ribose] polymerase 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	41.01Å 85.58Å 95.41Å 90.00° 90.03° 90.00°	Depositor
Resolution (Å)	36.98 – 2.50 36.98 – 2.50	Depositor EDS
% Data completeness (in resolution range)	99.3 (36.98-2.50) 99.3 (36.98-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.43 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.181 , 0.261 0.181 , 0.261	Depositor DCC
R_{free} test set	1102 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	40.2	Xtriage
Anisotropy	0.026	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 43.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.024 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5913	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 25I

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.59	0/2861	1.05	1/3860 (0.0%)
1	B	0.62	0/2861	1.09	2/3860 (0.1%)
All	All	0.60	0/5722	1.07	3/7720 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	3

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	396	ASP	CA-CB-CG	7.22	119.82	112.60
1	B	422	ASN	CB-CA-C	6.42	120.04	111.82
1	A	568	ARG	CB-CA-C	-5.07	104.03	111.23

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	308	ARG	Sidechain
1	B	431	ARG	Sidechain
1	B	478	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2802	0	2808	34	0
1	B	2802	0	2808	45	0
2	A	6	0	8	1	0
2	B	6	0	8	0	0
3	A	34	0	0	1	0
3	B	34	0	0	0	0
4	A	119	0	0	2	0
4	B	110	0	0	3	0
All	All	5913	0	5632	80	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (80) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:384:GLN:O	1:B:388:SER:HB3	1.62	0.99
1:A:502:ASN:ND2	1:A:508:LEU:HD11	2.00	0.76
1:A:370:ARG:NH1	1:A:406:GLU:OE2	2.20	0.74
1:B:374:HIS:O	4:B:701:HOH:O	2.06	0.73
1:B:377:TYR:O	1:B:381:VAL:HG23	1.90	0.72
1:B:362:TYR:O	4:B:702:HOH:O	2.10	0.69
1:B:439:LEU:HD23	1:B:569:MET:HE1	1.74	0.69
1:B:568:ARG:NH2	4:B:705:HOH:O	2.26	0.67
1:A:509:LEU:O	1:A:510:GLN:HG2	2.02	0.59
1:B:391:ALA:HB1	1:B:556:TYR:CZ	2.39	0.57
1:A:481:ASN:OD1	4:A:701:HOH:O	2.17	0.57
1:A:351:ARG:O	1:A:352:GLN:HB2	2.05	0.56
1:A:253:GLU:HA	1:A:256:MET:HE3	1.88	0.55
1:B:430:SER:O	1:B:474:CYS:HA	2.07	0.55
1:A:331:ILE:CD1	1:A:454:GLY:HA3	2.37	0.55
1:B:497:GLU:HB3	1:B:518:LEU:HD11	1.90	0.54
1:B:261:ASN:ND2	1:B:312:ASP:HB2	2.22	0.53
1:B:500:GLU:HG2	1:B:548:ASN:ND2	2.24	0.53
1:A:553:THR:O	1:A:554:LEU:HB2	2.09	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:479:LEU:HD22	1:A:479:LEU:H	1.74	0.52
1:A:502:ASN:HD22	1:A:508:LEU:HD11	1.73	0.52
1:B:486:LEU:HD21	1:B:569:MET:HG2	1.91	0.51
1:B:373:ASP:O	1:B:376:SER:OG	2.25	0.51
1:A:361:HIS:HD2	4:A:705:HOH:O	1.93	0.50
1:B:489:GLU:HG3	1:B:570:ARG:CG	2.42	0.50
1:B:439:LEU:HD23	1:B:569:MET:CE	2.42	0.49
1:A:237:ARG:CZ	1:A:371:PRO:HB2	2.43	0.49
1:B:331:ILE:CD1	1:B:454:GLY:HA3	2.42	0.49
1:B:524:SER:O	1:B:525:SER:HB3	2.13	0.49
1:A:456:MET:HE1	1:A:554:LEU:HD21	1.95	0.49
1:B:237:ARG:CZ	1:B:371:PRO:HB2	2.44	0.47
1:B:363:ARG:HG2	1:B:363:ARG:HH11	1.78	0.47
1:B:388:SER:HG	1:B:539:GLY:H	1.60	0.47
1:B:279:TYR:CD1	1:B:450:ALA:HB2	2.50	0.47
1:B:261:ASN:HD22	1:B:312:ASP:HB2	1.81	0.46
1:B:304:GLU:HG2	1:B:308:ARG:HD2	1.97	0.46
1:B:265:ALA:HB2	1:B:310:PRO:HG2	1.97	0.46
1:B:350:GLU:O	1:B:351:ARG:HB2	2.16	0.46
2:A:601:GOL:H11	3:A:602:25I:C12	2.46	0.46
1:B:346:LEU:C	1:B:346:LEU:HD23	2.42	0.45
1:A:331:ILE:HD13	1:A:454:GLY:HA3	1.98	0.45
1:B:238:VAL:CG2	1:B:371:PRO:HD3	2.46	0.45
1:A:333:LEU:C	1:A:333:LEU:HD23	2.41	0.45
1:A:456:MET:HE3	1:A:501:ALA:HB2	1.99	0.45
1:A:551:GLY:O	1:A:552:TYR:C	2.59	0.45
1:A:253:GLU:O	1:A:257:GLU:HG3	2.15	0.45
1:B:308:ARG:O	1:B:310:PRO:HD3	2.16	0.45
1:B:481:ASN:HB2	1:B:577:PHE:O	2.17	0.45
1:B:419:ASP:OD1	1:B:419:ASP:N	2.48	0.45
1:B:262:THR:O	1:B:266:PRO:HA	2.16	0.45
1:A:345:LYS:HD2	1:A:345:LYS:C	2.42	0.44
1:B:330:LYS:HD3	1:B:330:LYS:HA	1.70	0.44
1:B:524:SER:O	1:B:525:SER:CB	2.65	0.44
1:B:573:LEU:HD12	1:B:573:LEU:N	2.33	0.44
1:A:546:ILE:O	1:A:546:ILE:HG23	2.17	0.44
1:B:368:ALA:HB3	1:B:408:GLU:HB3	1.99	0.44
1:B:388:SER:OG	1:B:539:GLY:N	2.44	0.44
1:A:497:GLU:HA	1:A:516:LYS:O	2.18	0.44
1:B:385:TYR:HA	1:B:538:LEU:HD23	2.00	0.43
1:A:560:ILE:N	1:A:560:ILE:HD12	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:536:VAL:HG12	1:B:538:LEU:HG	2.01	0.43
1:A:230:PRO:HA	1:A:354:LEU:HD12	1.99	0.43
1:A:265:ALA:HB2	1:A:310:PRO:HG2	2.01	0.43
1:B:268:GLY:HA2	1:B:344:ILE:HG21	2.00	0.43
1:B:249:GLN:O	1:B:253:GLU:HG3	2.19	0.43
1:A:286:GLU:OE2	1:A:453:THR:OG1	2.29	0.43
1:B:349:SER:O	1:B:351:ARG:N	2.52	0.42
1:B:489:GLU:HG3	1:B:570:ARG:HG3	2.01	0.42
1:A:432:MET:HA	1:A:432:MET:HE2	2.01	0.42
1:A:259:LYS:O	1:A:311:HIS:HA	2.18	0.42
1:A:553:THR:O	1:A:554:LEU:CB	2.68	0.42
1:A:550:ASP:O	1:A:551:GLY:C	2.63	0.42
1:B:489:GLU:HG3	1:B:570:ARG:HG2	2.03	0.41
1:A:505:ALA:HA	1:A:508:LEU:HD12	2.03	0.41
1:A:385:TYR:HA	1:A:538:LEU:HD23	2.03	0.41
1:A:487:LEU:HB2	1:A:571:TYR:HB2	2.02	0.41
1:B:340:ILE:O	1:B:344:ILE:HG12	2.21	0.41
1:B:484:LEU:HD23	1:B:574:LYS:HA	2.02	0.41
1:A:509:LEU:HD23	1:A:509:LEU:HA	1.90	0.40
1:A:285:ILE:HG13	1:A:301:ALA:HB1	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	351/353 (99%)	332 (95%)	14 (4%)	5 (1%)	9	17
1	B	351/353 (99%)	328 (93%)	18 (5%)	5 (1%)	9	17
All	All	702/706 (99%)	660 (94%)	32 (5%)	10 (1%)	9	17

All (10) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	350	GLU
1	B	376	SER
1	B	525	SER
1	A	352	GLN
1	A	552	TYR
1	B	351	ARG
1	A	554	LEU
1	A	553	THR
1	A	545	GLY
1	B	529	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	305/305 (100%)	296 (97%)	9 (3%)	36	64
1	B	305/305 (100%)	290 (95%)	15 (5%)	22	45
All	All	610/610 (100%)	586 (96%)	24 (4%)	28	55

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	317	THR
1	B	345	LYS
1	B	354	LEU
1	B	370	ARG
1	B	374	HIS
1	B	388	SER
1	B	403	ASP
1	B	409	LYS
1	B	413	LYS
1	B	432	MET
1	B	456	MET
1	B	479	LEU
1	B	480	LYS
1	B	524	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	570	ARG
1	A	271	THR
1	A	303	ASN
1	A	344	ILE
1	A	345	LYS
1	A	479	LEU
1	A	510	GLN
1	A	552	TYR
1	A	553	THR
1	A	570	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	352	GLN
1	B	366	HIS
1	B	422	ASN
1	B	548	ASN
1	B	557	ASN
1	B	576	GLN
1	B	578	ASN
1	A	361	HIS
1	A	555	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	25I	A	602	-	38,38,38	1.34	3 (7%)	43,57,57	1.19	4 (9%)
3	25I	B	602	-	38,38,38	1.25	4 (10%)	43,57,57	1.39	8 (18%)
2	GOL	B	601	-	5,5,5	1.35	1 (20%)	5,5,5	0.74	0
2	GOL	A	601	-	5,5,5	1.23	0	5,5,5	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	25I	A	602	-	-	0/18/27/27	0/5/5/5
3	25I	B	602	-	-	0/18/27/27	0/5/5/5
2	GOL	B	601	-	-	0/4/4/4	-
2	GOL	A	601	-	-	3/4/4/4	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	25I	C13-N02	3.98	1.38	1.34
3	A	602	25I	C04-N05	3.95	1.32	1.29
3	A	602	25I	N02-N04	-3.45	1.30	1.36
3	B	602	25I	C21-C20	2.70	1.44	1.38
3	B	602	25I	C05-C06	-2.35	1.47	1.51
2	B	601	GOL	O2-C2	2.17	1.49	1.43
3	B	602	25I	C01-N06	2.14	1.38	1.35
3	B	602	25I	C22-C02	2.07	1.43	1.39

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	25I	C17-C16-C08	-4.15	118.59	123.07
3	B	602	25I	C18-C17-C16	3.02	122.77	119.04

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	602	25I	C08-C09-N01	2.94	122.81	118.24
3	B	602	25I	C08-C09-N01	2.67	122.38	118.24
3	A	602	25I	F02-C15-C14	-2.66	106.79	112.25
3	B	602	25I	O02-C09-N01	-2.34	118.67	122.35
3	B	602	25I	C22-C02-C03	2.22	121.75	119.26
3	B	602	25I	C21-C22-C02	-2.20	115.82	119.80
3	B	602	25I	F01-C15-C14	-2.17	107.80	112.25
3	B	602	25I	C20-C21-C22	2.05	122.77	120.24
3	A	602	25I	O02-C09-C08	-2.03	116.07	120.06
3	A	602	25I	C22-C02-C03	2.01	121.51	119.26

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	601	GOL	C1-C2-C3-O3
2	A	601	GOL	O2-C2-C3-O3
2	A	601	GOL	O1-C1-C2-O2


There are no ring outliers.

2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	25I	1	0
2	A	601	GOL	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand 25I A 602



Ligand 25I B 602





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	353/353 (100%)	-0.20	5 (1%) 73 70	24, 41, 72, 126	0
1	B	353/353 (100%)	-0.08	3 (0%) 82 80	21, 42, 75, 106	0
All	All	706/706 (100%)	-0.14	8 (1%) 78 75	21, 42, 75, 126	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	546	ILE	3.8
1	A	551	GLY	3.2
1	A	553	THR	2.9
1	B	580	LEU	2.3
1	A	547	LEU	2.3
1	B	353	GLY	2.3
1	B	552	TYR	2.1
1	A	313	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	GOL	A	601	6/6	0.85	0.14	57,80,84,93	0
2	GOL	B	601	6/6	0.91	0.10	53,58,60,62	0
3	25I	A	602	34/34	0.95	0.06	24,28,39,44	0
3	25I	B	602	34/34	0.96	0.06	21,25,43,45	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around GOL B 601:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 25I A 602:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 25I B 602:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.