



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 06:03 AM UTC

PDB ID : 5HLH / pdb_00005hlh
Title : Crystal structure of the overoxidized AbfR bound to DNA
Authors : Liu, G.; Liu, X.; Gan, J.; Yang, C.-G.
Deposited on : 2016-01-15
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	I	24	96% .
1	J	24	88% 12%
1	K	24	92% 8%
1	L	24	96% .
1	M	24	96% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	N	24	 92% 8%
1	O	24	 96% •
1	P	24	 92% 8%
2	A	147	 88% • 8%
2	B	147	 90% • 6%
2	C	147	 86% • 10%
2	D	147	 86% 6% 7%
2	E	147	 84% 9% • 7%
2	F	147	 86% 6% • 7%
2	G	147	 86% • 11%
2	H	147	 84% 8% • 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	OCS	A	13	-	X	-	-
2	OCS	F	13	-	X	-	-

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 12949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	I	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			
1	J	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			
1	K	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			
1	L	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			
1	M	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			
1	N	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			
1	O	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			
1	P	24	Total	C	N	O	P	0	0	0
			489	234	90	142	23			

- Molecule 2 is a protein called MarR family transcriptional regulator.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	A	135	Total	C	N	O	S	0	0	0
			1129	713	193	219	4			
2	B	138	Total	C	N	O	S	0	0	0
			1153	729	196	223	5			
2	C	133	Total	C	N	O	S	0	0	0
			1110	703	189	214	4			
2	D	136	Total	C	N	O	S	0	0	0
			1131	715	194	217	5			
2	E	137	Total	C	N	O	S	0	0	0
			1136	718	195	218	5			
2	F	137	Total	C	N	O	S	0	0	0
			1145	724	195	221	5			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	G	131	Total	C	N	O	S	0	0	0
			1089	690	188	207	4			
2	H	137	Total	C	N	O	S	0	0	0
			1144	724	195	220	5			


There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ALA	-	expression tag	UNP A0A0N1EJ89
A	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89
B	0	ALA	-	expression tag	UNP A0A0N1EJ89
B	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89
C	0	ALA	-	expression tag	UNP A0A0N1EJ89
C	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89
D	0	ALA	-	expression tag	UNP A0A0N1EJ89
D	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89
E	0	ALA	-	expression tag	UNP A0A0N1EJ89
E	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89
F	0	ALA	-	expression tag	UNP A0A0N1EJ89
F	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89
G	0	ALA	-	expression tag	UNP A0A0N1EJ89
G	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89
H	0	ALA	-	expression tag	UNP A0A0N1EJ89
H	72	MET	LEU	engineered mutation	UNP A0A0N1EJ89

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain I:  96%




- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain J:  88%




- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain K:  92%



- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain L:  96%




- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain M:  96%




- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain N: 92% 8%



- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain O: 96% .




- Molecule 1: DNA (5'-D(*TP*AP*AP*CP*TP*CP*AP*AP*TP*CP*GP*CP*GP*CP*GP*CP*GP*AP*TP*TP*GP*AP*GP*T)-3')

Chain P: 92% 8%



- Molecule 2: MarR family transcriptional regulator

Chain A: 88% . 8%



- Molecule 2: MarR family transcriptional regulator

Chain B: 90% . 6%




- Molecule 2: MarR family transcriptional regulator


Chain C: 86% . 10%




- Molecule 2: MarR family transcriptional regulator




- Molecule 2: MarR family transcriptional regulator




- Molecule 2: MarR family transcriptional regulator



- Molecule 2: MarR family transcriptional regulator



- Molecule 2: MarR family transcriptional regulator



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	150.31Å 290.91Å 52.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 3.00 30.00 – 3.00	Depositor EDS
% Data completeness (in resolution range)	93.8 (30.00-3.00) 93.8 (30.00-3.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.64 (at 2.96Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.224 , 0.272 0.230 , 0.281	Depositor DCC
R_{free} test set	2336 reflections (4.75%)	wwPDB-VP
Wilson B-factor (Å ²)	55.7	Xtriage
Anisotropy	0.980	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 39.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	12949	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: OCS

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	I	0.51	0/548	0.88	0/844
1	J	0.51	0/548	0.86	0/844
1	K	0.48	0/548	0.83	0/844
1	L	0.53	0/548	0.89	0/844
1	M	0.50	0/548	0.85	0/844
1	N	0.52	0/548	0.82	0/844
1	O	0.49	0/548	0.86	0/844
1	P	0.55	0/548	0.91	0/844
2	A	0.82	0/1137	0.98	0/1528
2	B	0.84	0/1162	1.02	1/1561 (0.1%)
2	C	0.83	0/1118	1.00	0/1504
2	D	0.87	0/1139	1.04	0/1531
2	E	0.89	0/1144	0.97	0/1538
2	F	0.93	0/1153	1.04	0/1549
2	G	0.83	0/1097	0.98	0/1475
2	H	0.88	0/1153	1.02	0/1549
All	All	0.76	0/13487	0.96	1/18987 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	101	ASN	N-CA-C	5.38	117.84	111.33

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	I	489	0	272	4	0
1	J	489	0	272	2	0
1	K	489	0	272	2	0
1	L	489	0	272	2	0
1	M	489	0	272	1	0
1	N	489	0	272	2	0
1	O	489	0	272	1	0
1	P	489	0	272	3	0
2	A	1129	0	1117	3	0
2	B	1153	0	1146	2	0
2	C	1110	0	1101	1	0
2	D	1131	0	1133	3	0
2	E	1136	0	1137	8	0
2	F	1145	0	1141	6	0
2	G	1089	0	1090	2	0
2	H	1144	0	1140	6	0
All	All	12949	0	11181	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:1:DT:H71	1:L:24:DT:H2''	1.78	0.64
2:F:43:VAL:HG21	2:F:72:MET:HE2	1.83	0.61
2:E:129:THR:HB	2:F:132:VAL:CG1	2.35	0.57
1:M:8:DA:N7	2:E:66:SER:OG	2.33	0.55
1:I:1:DT:C7	1:L:24:DT:H2''	2.38	0.53
1:I:1:DT:O2	1:K:3:DA:C2	2.64	0.51
1:P:14:DC:OP2	2:G:75:ARG:NH1	2.44	0.51
2:A:47:LEU:HD11	2:A:58:ILE:HD11	1.92	0.51
2:E:88:GLN:HA	2:E:88:GLN:OE1	2.11	0.50
1:P:14:DC:OP1	2:G:37:THR:OG1	2.27	0.50
1:N:8:DA:N7	2:F:66:SER:OG	2.38	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:94:LEU:HD22	2:E:96:ILE:HD11	1.96	0.46
2:E:20:SER:HB3	2:F:16:ALA:O	2.15	0.46
2:D:54:THR:O	2:D:55:LEU:C	2.60	0.44
1:J:14:DC:OP2	2:A:75:ARG:NH1	2.51	0.44
2:F:55:LEU:C	2:F:55:LEU:HD23	2.43	0.43
1:P:15:DG:OP2	2:H:17:TYR:OH	2.29	0.43
2:E:140:LEU:HD21	2:F:23:PHE:CE1	2.53	0.43
1:N:14:DC:OP2	2:E:75:ARG:NH1	2.50	0.43
2:E:55:LEU:C	2:E:55:LEU:HD23	2.44	0.42
2:H:55:LEU:C	2:H:55:LEU:HD23	2.44	0.42
2:H:87:GLN:HE21	2:H:90:ASP:HB3	1.85	0.42
1:O:14:DC:OP2	2:H:75:ARG:NH1	2.53	0.41
2:B:87:GLN:HE21	2:B:95:ILE:HD11	1.86	0.41
2:C:128:GLU:O	2:C:129:THR:C	2.62	0.41
1:I:1:DT:H5'	1:I:1:DT:H6	1.85	0.41
2:D:55:LEU:C	2:D:55:LEU:HD23	2.45	0.41
2:A:143:LEU:HD11	2:B:116:CYS:O	2.21	0.41
2:H:54:THR:O	2:H:55:LEU:C	2.64	0.41
2:H:12:LEU:O	2:H:13:OCS:C	2.69	0.40
1:J:6:DC:H2'	1:J:7:DA:C8	2.56	0.40
1:K:14:DC:OP2	2:D:75:ARG:NH1	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	A	130/147 (88%)	129 (99%)	1 (1%)	0	100	100
2	B	133/147 (90%)	133 (100%)	0	0	100	100
2	C	128/147 (87%)	125 (98%)	3 (2%)	0	100	100
2	D	131/147 (89%)	129 (98%)	2 (2%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	E	132/147 (90%)	131 (99%)	1 (1%)	0	100	100
2	F	132/147 (90%)	131 (99%)	1 (1%)	0	100	100
2	G	126/147 (86%)	126 (100%)	0	0	100	100
2	H	132/147 (90%)	130 (98%)	2 (2%)	0	100	100
All	All	1044/1176 (89%)	1034 (99%)	10 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	A	125/136 (92%)	124 (99%)	1 (1%)	73	86
2	B	128/136 (94%)	127 (99%)	1 (1%)	73	86
2	C	123/136 (90%)	120 (98%)	3 (2%)	43	73
2	D	126/136 (93%)	121 (96%)	5 (4%)	28	62
2	E	126/136 (93%)	121 (96%)	5 (4%)	28	62
2	F	127/136 (93%)	124 (98%)	3 (2%)	43	73
2	G	121/136 (89%)	119 (98%)	2 (2%)	53	78
2	H	127/136 (93%)	122 (96%)	5 (4%)	28	62
All	All	1003/1088 (92%)	978 (98%)	25 (2%)	42	72

All (25) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	A	8	LEU
2	B	105	GLN
2	C	37	THR
2	C	105	GLN
2	C	113	ILE
2	D	1	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	D	11	GLN
2	D	12	LEU
2	D	23	PHE
2	D	39	SER
2	E	1	MET
2	E	23	PHE
2	E	96	ILE
2	E	97	THR
2	E	135	GLU
2	F	39	SER
2	F	72	MET
2	F	143	LEU
2	G	39	SER
2	G	143	LEU
2	H	39	SER
2	H	105	GLN
2	H	111	GLU
2	H	113	ILE
2	H	130	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
2	A	142	HIS
2	B	25	GLN
2	B	56	ASN
2	C	40	GLN
2	C	142	HIS
2	D	11	GLN
2	D	40	GLN
2	D	87	GLN
2	D	101	ASN
2	D	142	HIS
2	E	56	ASN
2	E	87	GLN
2	F	56	ASN
2	F	93	GLN
2	F	103	GLN
2	G	103	GLN
2	H	103	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	OCS	F	13	2	6,8,9	2.40	3 (50%)	7,11,13	5.84	6 (85%)
2	OCS	A	13	2	6,8,9	3.65	2 (33%)	7,11,13	3.94	6 (85%)
2	OCS	D	13	2	6,8,9	3.34	2 (33%)	7,11,13	4.79	4 (57%)
2	OCS	B	13	2	2,7,9	0.83	0	2,8,13	5.80	2 (100%)
2	OCS	C	13	2	6,8,9	3.28	2 (33%)	7,11,13	4.44	5 (71%)
2	OCS	H	13	2	2,7,9	0.78	0	2,8,13	5.51	2 (100%)
2	OCS	E	13	2	6,8,9	3.35	2 (33%)	7,11,13	6.08	6 (85%)
2	OCS	G	13	2	2,7,9	0.72	0	2,8,13	6.17	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	OCS	F	13	2	-	4/4/7/9	-
2	OCS	A	13	2	-	4/4/7/9	-
2	OCS	D	13	2	-	1/4/7/9	-
2	OCS	B	13	2	-	1/3/6/9	-
2	OCS	C	13	2	-	1/4/7/9	-
2	OCS	H	13	2	-	3/3/6/9	-
2	OCS	E	13	2	-	1/4/7/9	-
2	OCS	G	13	2	-	0/3/6/9	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	13	OCS	CB-CA	6.65	1.58	1.53
2	A	13	OCS	CB-CA	6.38	1.58	1.53
2	A	13	OCS	OD3-SG	6.06	1.62	1.45
2	E	13	OCS	OD1-SG	5.97	1.62	1.45
2	C	13	OCS	CB-CA	5.65	1.57	1.53
2	C	13	OCS	OD3-SG	5.40	1.60	1.45
2	E	13	OCS	CB-CA	5.37	1.57	1.53
2	D	13	OCS	OD2-SG	4.46	1.64	1.47
2	F	13	OCS	OD2-SG	4.16	1.62	1.47
2	F	13	OCS	CB-CA	3.31	1.56	1.53
2	F	13	OCS	OD3-SG	-2.24	1.38	1.45

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	13	OCS	OD3-SG-CB	9.48	120.91	106.76
2	F	13	OCS	OD3-SG-CB	9.14	120.41	106.76
2	G	13	OCS	OD1-SG-CB	8.70	117.86	105.35
2	E	13	OCS	OD3-SG-OD1	-8.55	86.02	113.82
2	D	13	OCS	OD3-SG-CB	8.21	119.01	106.76
2	E	13	OCS	CA-CB-SG	7.26	124.39	113.61
2	F	13	OCS	OD1-SG-CB	7.01	117.22	106.76
2	C	13	OCS	CA-CB-SG	6.68	123.53	113.61
2	F	13	OCS	OD2-SG-OD3	-6.68	94.69	111.40
2	D	13	OCS	OD3-SG-OD1	-6.49	92.71	113.82
2	A	13	OCS	CA-CB-SG	6.06	122.61	113.61
2	H	13	OCS	OD1-SG-CB	6.02	114.01	105.35
2	C	13	OCS	OD2-SG-CB	5.92	117.41	105.97
2	B	13	OCS	OD1-SG-CB	5.92	113.86	105.35
2	C	13	OCS	OD2-SG-OD3	-5.86	96.75	111.40
2	D	13	OCS	OD1-SG-CB	5.80	115.42	106.76
2	B	13	OCS	OD3-SG-CB	5.68	113.52	105.35
2	F	13	OCS	CA-CB-SG	5.52	121.82	113.61
2	A	13	OCS	OD1-SG-CB	5.02	114.25	106.76
2	H	13	OCS	OD3-SG-CB	4.95	112.47	105.35
2	E	13	OCS	OD1-SG-CB	4.93	114.11	106.76
2	D	13	OCS	CA-CB-SG	4.14	119.76	113.61
2	F	13	OCS	OD3-SG-OD1	-4.13	100.38	113.82
2	C	13	OCS	OD1-SG-CB	3.86	112.52	106.76
2	A	13	OCS	OD2-SG-OD3	-3.70	102.15	111.40
2	A	13	OCS	OD3-SG-CB	3.62	112.16	106.76
2	F	13	OCS	OD2-SG-CB	3.49	112.71	105.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	13	OCS	OD2-SG-CB	3.47	112.68	105.97
2	E	13	OCS	OD2-SG-CB	3.23	112.20	105.97
2	A	13	OCS	OD3-SG-OD1	-2.74	104.90	113.82
2	E	13	OCS	OD2-SG-OD3	-2.37	105.47	111.40
2	C	13	OCS	OD2-SG-OD1	-2.33	105.58	111.40

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	13	OCS	N-CA-CB-SG
2	A	13	OCS	CA-CB-SG-OD2
2	C	13	OCS	N-CA-CB-SG
2	D	13	OCS	N-CA-CB-SG
2	E	13	OCS	N-CA-CB-SG
2	F	13	OCS	N-CA-CB-SG
2	F	13	OCS	CA-CB-SG-OD1
2	F	13	OCS	CA-CB-SG-OD2
2	F	13	OCS	CA-CB-SG-OD3
2	H	13	OCS	N-CA-CB-SG
2	H	13	OCS	CA-CB-SG-OD3
2	A	13	OCS	CA-CB-SG-OD1
2	A	13	OCS	CA-CB-SG-OD3
2	B	13	OCS	CA-CB-SG-OD3
2	H	13	OCS	CA-CB-SG-OD1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	H	13	OCS	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	I	24/24 (100%)	0.20	0 100 100	47, 66, 80, 84	0
1	J	24/24 (100%)	0.28	0 100 100	51, 63, 79, 89	0
1	K	24/24 (100%)	0.01	0 100 100	42, 61, 87, 92	0
1	L	24/24 (100%)	0.08	0 100 100	48, 62, 69, 77	0
1	M	24/24 (100%)	0.19	0 100 100	40, 62, 76, 81	0
1	N	24/24 (100%)	-0.05	0 100 100	43, 61, 67, 67	0
1	O	24/24 (100%)	0.16	0 100 100	46, 66, 78, 81	0
1	P	24/24 (100%)	0.33	0 100 100	47, 65, 77, 78	0
2	A	134/147 (91%)	0.36	2 (1%) 72 49	57, 92, 122, 134	0
2	B	137/147 (93%)	0.16	4 (2%) 53 31	48, 74, 123, 136	0
2	C	132/147 (89%)	0.24	1 (0%) 82 64	52, 82, 110, 118	0
2	D	135/147 (91%)	0.13	3 (2%) 62 39	48, 69, 122, 134	0
2	E	136/147 (92%)	0.04	0 100 100	48, 73, 108, 124	0
2	F	136/147 (92%)	0.01	2 (1%) 72 49	45, 66, 110, 129	0
2	G	130/147 (88%)	0.24	3 (2%) 61 38	57, 88, 110, 124	0
2	H	136/147 (92%)	0.03	1 (0%) 84 66	49, 71, 123, 141	0
All	All	1268/1368 (92%)	0.15	16 (1%) 75 53	40, 74, 116, 141	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	A	108	ALA	4.6
2	G	33	GLN	3.4
2	D	114	SER	3.3
2	B	131	TYR	3.1
2	C	135	GLU	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	142	HIS	2.8
2	B	15	SER	2.8
2	D	1	MET	2.8
2	A	49	GLU	2.6
2	D	115	SER	2.5
2	G	102	GLY	2.4
2	F	1	MET	2.3
2	B	85	GLU	2.2
2	B	1	MET	2.2
2	G	24	ALA	2.1
2	H	8	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	OCS	H	13	8/10	0.89	0.11	78,80,84,85	0
2	OCS	F	13	9/10	0.91	0.13	52,59,65,68	0
2	OCS	E	13	9/10	0.91	0.14	53,56,59,60	0
2	OCS	D	13	9/10	0.92	0.14	65,72,79,83	0
2	OCS	A	13	9/10	0.92	0.11	55,66,76,81	0
2	OCS	B	13	8/10	0.93	0.11	76,86,93,95	0
2	OCS	G	13	8/10	0.95	0.10	51,63,69,76	0
2	OCS	C	13	9/10	0.95	0.10	62,68,77,78	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.