



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 10:23 PM UTC

PDB ID : 6HL9 / pdb_00006hl9
Title : Crystal Structure of the CsiD Glutarate Hydroxylase in complex with Succinate
Authors : Williams, R.M.; Mayans, O.; Hartig, J.S.
Deposited on : 2018-09-10
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	353	6% 79% 5% 15%
1	B	353	7% 78% 5% 17%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 5408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutarate 2-hydroxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	299	Total	C	N	O	S	0	2	0
			2438	1560	428	440	10			
1	B	294	Total	C	N	O	S	0	3	0
			2425	1550	427	438	10			

There are 56 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-19	MET	-	initiating methionine	UNP P76621
A	-18	GLY	-	expression tag	UNP P76621
A	-17	SER	-	expression tag	UNP P76621
A	-16	SER	-	expression tag	UNP P76621
A	-15	HIS	-	expression tag	UNP P76621
A	-14	HIS	-	expression tag	UNP P76621
A	-13	HIS	-	expression tag	UNP P76621
A	-12	HIS	-	expression tag	UNP P76621
A	-11	HIS	-	expression tag	UNP P76621
A	-10	HIS	-	expression tag	UNP P76621
A	-9	SER	-	expression tag	UNP P76621
A	-8	SER	-	expression tag	UNP P76621
A	-7	GLY	-	expression tag	UNP P76621
A	-6	LEU	-	expression tag	UNP P76621
A	-5	VAL	-	expression tag	UNP P76621
A	-4	PRO	-	expression tag	UNP P76621
A	-3	ARG	-	expression tag	UNP P76621
A	-2	GLY	-	expression tag	UNP P76621
A	-1	SER	-	expression tag	UNP P76621
A	0	HIS	-	expression tag	UNP P76621
A	326	LEU	-	expression tag	UNP P76621
A	327	GLU	-	expression tag	UNP P76621
A	328	HIS	-	expression tag	UNP P76621
A	329	HIS	-	expression tag	UNP P76621
A	330	HIS	-	expression tag	UNP P76621

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	331	HIS	-	expression tag	UNP P76621
A	332	HIS	-	expression tag	UNP P76621
A	333	HIS	-	expression tag	UNP P76621
B	-19	MET	-	initiating methionine	UNP P76621
B	-18	GLY	-	expression tag	UNP P76621
B	-17	SER	-	expression tag	UNP P76621
B	-16	SER	-	expression tag	UNP P76621
B	-15	HIS	-	expression tag	UNP P76621
B	-14	HIS	-	expression tag	UNP P76621
B	-13	HIS	-	expression tag	UNP P76621
B	-12	HIS	-	expression tag	UNP P76621
B	-11	HIS	-	expression tag	UNP P76621
B	-10	HIS	-	expression tag	UNP P76621
B	-9	SER	-	expression tag	UNP P76621
B	-8	SER	-	expression tag	UNP P76621
B	-7	GLY	-	expression tag	UNP P76621
B	-6	LEU	-	expression tag	UNP P76621
B	-5	VAL	-	expression tag	UNP P76621
B	-4	PRO	-	expression tag	UNP P76621
B	-3	ARG	-	expression tag	UNP P76621
B	-2	GLY	-	expression tag	UNP P76621
B	-1	SER	-	expression tag	UNP P76621
B	0	HIS	-	expression tag	UNP P76621
B	326	LEU	-	expression tag	UNP P76621
B	327	GLU	-	expression tag	UNP P76621
B	328	HIS	-	expression tag	UNP P76621
B	329	HIS	-	expression tag	UNP P76621
B	330	HIS	-	expression tag	UNP P76621
B	331	HIS	-	expression tag	UNP P76621
B	332	HIS	-	expression tag	UNP P76621
B	333	HIS	-	expression tag	UNP P76621

- Molecule 2 is FE (II) ION (CCD ID: FE2) (formula: Fe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe 1 1	0	0
2	B	1	Total Fe 1 1	0	0

- Molecule 3 is SUCCINIC ACID (CCD ID: SIN) (formula: C₄H₆O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			8	4	4		
3	B	1	Total	C	O	0	0
			8	4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	309	Total	O	0	0
			309	309		
4	B	218	Total	O	0	0
			218	218		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glutarate 2-hydroxylase



• Molecule 1: Glutarate 2-hydroxylase



4 Data and refinement statistics

Property	Value	Source
Space group	P 4 21 2	Depositor
Cell constants a, b, c, α , β , γ	123.03Å 123.03Å 138.33Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.11 – 2.30 46.11 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (46.11-2.30) 100.0 (46.11-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.14	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.186 , 0.210 0.189 , 0.190	Depositor DCC
R_{free} test set	2592 reflections (3.46%)	wwPDB-VP
Wilson B-factor (Å ²)	37.4	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5408	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE2, SIN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.78	0/2504	0.78	0/3395
1	B	0.70	0/2486	0.79	1/3366 (0.0%)
All	All	0.74	0/4990	0.79	1/6761 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	155	ARG	CB-CG-CD	7.54	128.64	111.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2438	0	2376	11	0
1	B	2425	0	2381	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	8	0	4	0	0
3	B	8	0	4	0	0
4	A	309	0	0	1	1
4	B	218	0	0	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	5408	0	4765	29	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (29) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:242[B]:ARG:NH2	4:B:501:HOH:O	2.13	0.80
1:B:242[B]:ARG:HH21	1:B:242[B]:ARG:HG2	1.58	0.68
1:B:183:MET:HE2	1:B:305[A]:ARG:NH2	2.11	0.66
1:A:305[B]:ARG:CZ	1:A:307:LEU:HD21	2.30	0.61
1:B:40:GLN:NE2	4:B:503:HOH:O	2.33	0.61
1:B:305[B]:ARG:CZ	1:B:307:LEU:HD21	2.33	0.58
1:B:242[B]:ARG:HH21	1:B:242[B]:ARG:CG	2.19	0.55
1:A:203:PHE:O	1:A:209:ARG:NH2	2.41	0.53
1:A:281:LYS:NZ	4:A:512:HOH:O	2.45	0.49
1:B:186:GLY:O	1:B:305[A]:ARG:NH1	2.46	0.48
1:A:200:ASP:O	1:A:204:ARG:HD3	2.14	0.46
1:B:180:GLU:OE1	1:B:305[A]:ARG:NH2	2.47	0.46
1:B:168:GLU:CD	4:B:515:HOH:O	2.59	0.46
1:B:155:ARG:HH22	1:B:158:GLU:CD	2.24	0.46
1:A:136:ALA:O	1:A:308:MET:HA	2.16	0.46
1:B:183:MET:HE2	1:B:305[A]:ARG:HH21	1.81	0.45
1:B:139:VAL:HG22	1:B:306:GLU:HG3	1.99	0.45
1:B:306:GLU:C	1:B:307:LEU:HD22	2.42	0.45
1:B:35:LEU:O	1:B:97:ILE:HA	2.18	0.44
1:A:153:PRO:HB2	1:A:154:HIS:CE1	2.52	0.44
1:B:123:GLY:HA3	1:B:313:TYR:O	2.18	0.43
1:A:62:PHE:CD1	1:A:62:PHE:C	2.97	0.42
1:A:123:GLY:HA3	1:A:313:TYR:O	2.20	0.42
1:B:54:VAL:O	1:B:58:GLU:HG3	2.20	0.42
1:A:183:MET:HE2	1:A:305[A]:ARG:CZ	2.50	0.42
1:A:45:PHE:O	1:A:49:VAL:HG22	2.20	0.41
1:A:35:LEU:O	1:A:97:ILE:HA	2.21	0.41
1:B:155:ARG:NH2	1:B:158:GLU:OE1	2.54	0.40
1:B:294:ARG:HG2	1:B:295:ASP:N	2.37	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:711:HOH:O	4:A:711:HOH:O[8_556]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/353 (84%)	290 (98%)	5 (2%)	0	100	100
1	B	291/353 (82%)	284 (98%)	7 (2%)	0	100	100
All	All	586/706 (83%)	574 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/312 (83%)	260 (100%)	0	100	100
1	B	261/312 (84%)	261 (100%)	0	100	100
All	All	521/624 (84%)	521 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	40	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	182	ASN
1	B	98	ASN
1	B	198	HIS
1	B	310	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SIN	A	402	2	7,7,7	1.05	0	8,8,8	1.58	2 (25%)
3	SIN	B	402	-	7,7,7	1.10	0	8,8,8	1.54	1 (12%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SIN	A	402	2	-	3/5/5/5	-
3	SIN	B	402	-	-	2/5/5/5	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	SIN	O2-C1-C2	2.24	121.09	114.00
3	A	402	SIN	C3-C2-C1	-2.20	107.83	113.67
3	A	402	SIN	O4-C4-C3	2.16	120.83	114.00

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	402	SIN	C2-C3-C4-O3
3	A	402	SIN	C2-C3-C4-O3
3	B	402	SIN	C2-C3-C4-O4
3	A	402	SIN	C2-C3-C4-O4
3	A	402	SIN	O2-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	299/353 (84%)	0.04	22 (7%)	20 22	17, 37, 73, 105	2 (0%)
1	B	294/353 (83%)	0.24	26 (8%)	15 17	20, 42, 85, 124	3 (1%)
All	All	593/706 (83%)	0.14	48 (8%)	18 19	17, 39, 82, 124	5 (0%)

All (48) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	154	HIS	6.4
1	B	218	PRO	5.2
1	B	140	VAL	5.1
1	A	150	LEU	5.1
1	B	142	ASN	5.0
1	A	218	PRO	4.9
1	A	323	THR	4.6
1	A	149	TYR	4.4
1	B	130	MET	4.3
1	B	324	HIS	4.3
1	B	223	SER	4.0
1	A	17	ASP	3.6
1	A	324	HIS	3.5
1	A	297	PHE	3.5
1	A	223	SER	3.4
1	A	141	LYS	3.4
1	A	16	GLN	3.4
1	A	154	HIS	3.3
1	A	142	ASN	3.2
1	B	141	LYS	3.1
1	B	217	PRO	3.0
1	A	217	PRO	2.8
1	B	184	GLN	2.8
1	B	157	MET	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	168	GLU	2.8
1	B	323	THR	2.8
1	B	139	VAL	2.8
1	B	169	ILE	2.7
1	A	157	MET	2.7
1	A	152	GLN	2.7
1	B	155	ARG	2.6
1	A	130	MET	2.5
1	A	153	PRO	2.5
1	B	138	PHE	2.5
1	B	43	LYS	2.5
1	B	156	VAL	2.5
1	A	307	LEU	2.4
1	A	151	ARG	2.4
1	B	183	MET	2.3
1	B	307	LEU	2.3
1	B	59	TYR	2.3
1	A	183	MET	2.2
1	A	139	VAL	2.2
1	B	305[A]	ARG	2.2
1	B	224	LYS	2.1
1	B	301	PRO	2.1
1	B	296	ARG	2.0
1	A	204	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SIN	B	402	8/8	0.84	0.16	85,85,87,88	0
3	SIN	A	402	8/8	0.87	0.14	64,67,73,74	0
2	FE2	A	401	1/1	0.99	0.02	47,47,47,47	0
2	FE2	B	401	1/1	0.99	0.05	55,55,55,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.