



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 03:53 AM UTC

PDB ID : 8HLN / pdb_00008hln
Title : Crystal structure of p53/BCL2 fusion complex(complex3)
Authors : Guo, M.; Wei, H.; Wang, H.; Chen, Y.
Deposited on : 2022-11-30
Resolution : 2.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	6% 87% 12% .
2	B	175	7% 69% 12% .. 18%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2663 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cellular tumor antigen p53.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	0	0
			1502	927	273	286	16			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	91	GLY	-	expression tag	UNP P04637
A	92	SER	-	expression tag	UNP P04637
A	93	GLY	-	expression tag	UNP P04637
A	94	GLY	-	expression tag	UNP P04637

- Molecule 2 is a protein called Apoptosis regulator Bcl-2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	143	Total	C	N	O	S	0	0	0
			1119	712	191	210	6			

There are 49 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	6	GLY	-	expression tag	UNP P10415
B	7	PRO	-	expression tag	UNP P10415
B	8	SER	-	expression tag	UNP P10415
B	9	GLN	-	expression tag	UNP P10415
B	20	SER	HIS	engineered mutation	UNP P10415
B	78	GLU	-	linker	UNP P10415
B	79	GLU	-	linker	UNP P10415
B	80	ASN	-	linker	UNP P10415
B	81	ARG	-	linker	UNP P10415
B	82	THR	-	linker	UNP P10415
B	83	GLU	-	linker	UNP P10415
B	84	ALA	-	linker	UNP P10415

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	85	PRO	-	linker	UNP P10415
B	86	GLU	-	linker	UNP P10415
B	87	GLY	-	linker	UNP P10415
B	88	THR	-	linker	UNP P10415
B	89	GLU	-	linker	UNP P10415
B	90	SER	-	linker	UNP P10415
B	91	GLU	-	linker	UNP P10415
B	95	GLN	LEU	engineered mutation	UNP P10415
B	106	LEU	ARG	engineered mutation	UNP P10415
B	124	GLY	PHE	engineered mutation	UNP P10415
B	127	TYR	ARG	engineered mutation	UNP P10415
B	128	ALA	GLY	engineered mutation	UNP P10415
B	129	SER	ARG	engineered mutation	UNP P10415
B	168	VAL	PRO	engineered mutation	UNP P10415
B	175	ALA	LEU	engineered mutation	UNP P10415
B	178	ALA	THR	engineered mutation	UNP P10415
B	179	THR	GLU	engineered mutation	UNP P10415
B	183	ASP	ARG	engineered mutation	UNP P10415
B	204	ASN	-	expression tag	UNP P10415
B	205	ASN	-	expression tag	UNP P10415
B	206	GLY	-	expression tag	UNP P10415
B	207	GLY	-	expression tag	UNP P10415
B	208	SER	-	expression tag	UNP P10415
B	209	GLY	-	expression tag	UNP P10415
B	210	GLY	-	expression tag	UNP P10415
B	211	GLY	-	expression tag	UNP P10415
B	212	GLY	-	expression tag	UNP P10415
B	213	SER	-	expression tag	UNP P10415
B	214	LEU	-	expression tag	UNP P10415
B	215	VAL	-	expression tag	UNP P10415
B	216	PRO	-	expression tag	UNP P10415
B	217	ARG	-	expression tag	UNP P10415
B	218	GLY	-	expression tag	UNP P10415
B	219	SER	-	expression tag	UNP P10415
B	220	GLY	-	expression tag	UNP P10415
B	221	GLY	-	expression tag	UNP P10415
B	222	GLY	-	expression tag	UNP P10415

- Molecule 3 is ZINC ION (CCD ID: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	30	Total 30	O 30	0	0
4	B	11	Total 11	O 11	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cellular tumor antigen p53



- Molecule 2: Apoptosis regulator Bcl-2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	44.70Å 70.53Å 127.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.83 – 2.35 30.83 – 2.35	Depositor EDS
% Data completeness (in resolution range)	96.8 (30.83-2.35) 96.7 (30.83-2.35)	Depositor EDS
R_{merge}	0.17	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.85 (at 2.36Å)	Xtriage
Refinement program	PHENIX 1.15.2_3472	Depositor
R, R_{free}	0.202 , 0.253 0.213 , 0.259	Depositor DCC
R_{free} test set	1687 reflections (9.71%)	wwPDB-VP
Wilson B-factor (Å ²)	39.4	Xtriage
Anisotropy	0.866	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 35.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2663	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.32	0/1536	0.42	0/2086
2	B	0.51	0/1147	0.49	2/1557 (0.1%)
All	All	0.41	0/2683	0.45	2/3643 (0.1%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	107	ARG	N-CA-C	5.53	118.24	111.82
2	B	200	GLU	CB-CG-CD	5.13	121.32	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1502	0	1436	18	0
2	B	1119	0	1021	17	0
3	A	1	0	0	0	0
4	A	30	0	0	0	0
4	B	11	0	0	1	0
All	All	2663	0	2457	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 32 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:133:MET:HE1	1:A:135:CYS:SG	2.24	0.77
1:A:147:VAL:HG11	1:A:151:PRO:HD3	1.68	0.76
2:B:143:ASN:ND2	2:B:146:ARG:H	1.83	0.75
1:A:133:MET:CE	1:A:135:CYS:SG	2.77	0.73
2:B:123:PRO:HA	2:B:169:LEU:HD21	1.73	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/200 (98%)	187 (95%)	9 (5%)	0	100	100
2	B	139/175 (79%)	136 (98%)	3 (2%)	0	100	100
All	All	335/375 (89%)	323 (96%)	12 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/177 (94%)	167 (100%)	0	100	100
2	B	112/140 (80%)	107 (96%)	5 (4%)	24	32
All	All	279/317 (88%)	274 (98%)	5 (2%)	51	66

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	107	ARG
2	B	114	GLU
2	B	118	GLN
2	B	143	ASN
2	B	201	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	235	ASN
1	A	268	ASN
2	B	143	ASN
2	B	184	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/200 (99%)	0.45	13 (6%) 24 27	27, 51, 84, 103	0
2	B	143/175 (81%)	0.53	13 (9%) 15 17	33, 53, 82, 110	0
All	All	341/375 (90%)	0.48	26 (7%) 20 22	27, 52, 84, 110	0

The worst 5 of 26 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	288	ASN	3.7
1	A	187	GLY	3.4
2	B	90	SER	3.3
2	B	7	PRO	3.2
2	B	86	GLU	3.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	ZN	A	401	1/1	0.66	0.17	30,30,30,30	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers ⓘ

There are no such residues in this entry.