



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 17, 2026 – 10:22 PM UTC

PDB ID : 4HMZ / pdb_00004hmz
Title : Crystal Structure of ChmJ, a 3'-monoepimerase from *Streptomyces bikiniensis*
in complex with dTDP-quinovose
Authors : Holden, H.M.; Kubiak, R.L.
Deposited on : 2012-10-18
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	FAILED
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	FAILED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	200	
1	B	200	
1	C	200	
1	D	200	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6861 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative 3-epimerase in D-allose pathway.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	198	Total	C	N	O	S	0	1	0
			1560	980	283	293	4			
1	B	199	Total	C	N	O	S	0	0	0
			1559	980	282	293	4			
1	C	200	Total	C	N	O	S	0	4	0
			1593	1002	290	297	4			
1	D	200	Total	C	N	O	S	0	3	0
			1596	1004	296	292	4			

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	197	LEU	-	expression tag	UNP Q5SFD1
A	198	GLU	-	expression tag	UNP Q5SFD1
A	199	HIS	-	expression tag	UNP Q5SFD1
A	200	HIS	-	expression tag	UNP Q5SFD1
B	197	LEU	-	expression tag	UNP Q5SFD1
B	198	GLU	-	expression tag	UNP Q5SFD1
B	199	HIS	-	expression tag	UNP Q5SFD1
B	200	HIS	-	expression tag	UNP Q5SFD1
C	197	LEU	-	expression tag	UNP Q5SFD1
C	198	GLU	-	expression tag	UNP Q5SFD1
C	199	HIS	-	expression tag	UNP Q5SFD1
C	200	HIS	-	expression tag	UNP Q5SFD1
D	197	LEU	-	expression tag	UNP Q5SFD1
D	198	GLU	-	expression tag	UNP Q5SFD1
D	199	HIS	-	expression tag	UNP Q5SFD1
D	200	HIS	-	expression tag	UNP Q5SFD1

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	B	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	C	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		
2	D	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is [(2R,3S,5R)-3-hydroxy-5-(5-methyl-2,4-dioxo-3,4-dihydropyrimidin-1(2H)-yl)tetrahydrofuran-2-yl]methyl (2R,3R,4S,5S,6R)-3,4,5-trihydroxy-6-methyltetrahydro-2H-pyran-2-yl dihydrogen diphosphate (CCD ID: 18T) (formula: C₁₆H₂₆N₂O₁₅P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 35	C 16	N 2	O 15	P 2	0	0
3	B	1	Total 35	C 16	N 2	O 15	P 2	0	0
3	C	1	Total 35	C 16	N 2	O 15	P 2	0	0
3	D	1	Total 35	C 16	N 2	O 15	P 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	123	Total O 123 123	0	0
4	B	99	Total O 99 99	0	0
4	C	79	Total O 79 79	0	0
4	D	84	Total O 84 84	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative 3-epimerase in D-allose pathway

Chain A:  86% 12% .




- Molecule 1: Putative 3-epimerase in D-allose pathway

Chain B:  88% 10% .



- Molecule 1: Putative 3-epimerase in D-allose pathway

Chain C:  80% 18% .



- Molecule 1: Putative 3-epimerase in D-allose pathway

Chain D:  86% 12% .



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	140.72Å 140.72Å 117.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.00 30.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	94.9 (30.00-2.00) 94.9 (30.00-2.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.71 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.199 , 0.249 0.201 , 0.250	Depositor DCC
R_{free} test set	3880 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtriage
Anisotropy	0.109	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 19.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.488 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6861	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 18T, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/1603	1.33	8/2188 (0.4%)
1	B	0.55	0/1603	1.33	8/2189 (0.4%)
1	C	0.53	0/1651	1.28	9/2254 (0.4%)
1	D	0.54	0/1654	1.28	3/2257 (0.1%)
All	All	0.55	0/6511	1.30	28/8888 (0.3%)

There are no bond length outliers.

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	157	ASP	N-CA-C	9.18	121.29	111.28
1	A	157	ASP	N-CA-C	8.20	122.24	111.75
1	B	157	ASP	N-CA-C	6.80	119.53	111.71
1	B	192	GLN	N-CA-C	-6.09	104.72	111.36
1	B	88	ILE	CB-CA-C	-6.05	102.15	111.71
1	C	159	ILE	N-CA-C	6.04	116.62	108.17
1	C	182	LEU	CA-C-N	-6.02	114.56	120.52
1	C	182	LEU	C-N-CA	-6.02	114.56	120.52
1	B	24	SER	CB-CA-C	-5.97	97.62	110.45
1	D	144	VAL	N-CA-C	-5.96	99.83	108.89
1	C	157	ASP	N-CA-C	5.89	118.66	111.82
1	C	99	VAL	N-CA-C	5.88	113.89	107.60
1	A	117	ARG	CA-CB-CG	-5.71	102.68	114.10
1	A	192	GLN	N-CA-C	-5.66	105.19	111.36
1	C	120	LEU	CB-CA-C	-5.53	101.13	110.19
1	A	87	ARG	N-CA-C	5.41	117.12	108.41
1	B	87	ARG	N-CA-C	5.40	117.10	108.41
1	A	12	GLN	CB-CG-CD	5.36	121.72	112.60
1	A	75	VAL	N-CA-C	5.33	117.23	112.17
1	B	70	LYS	CB-CG-CD	-5.29	99.14	111.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	179	LEU	N-CA-C	5.28	119.47	113.19
1	B	157	ASP	CB-CA-C	-5.28	100.87	110.63
1	D	120	LEU	CB-CA-C	-5.22	101.73	110.19
1	A	157	ASP	CB-CA-C	-5.17	100.42	110.46
1	C	50	VAL	CB-CA-C	-5.09	104.38	110.99
1	C	87	ARG	NE-CZ-NH1	-5.05	116.45	121.50
1	C	144	VAL	N-CA-C	-5.03	101.10	108.85
1	A	125	ASP	N-CA-C	5.00	118.97	112.86

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1560	0	1493	11	0
1	B	1559	0	1488	11	0
1	C	1593	0	1526	18	0
1	D	1596	0	1537	16	0
2	A	8	0	12	0	0
2	B	4	0	6	0	0
2	C	8	0	12	1	0
2	D	8	0	12	0	0
3	A	35	0	24	0	0
3	B	35	0	24	0	0
3	C	35	0	24	0	0
3	D	35	0	24	0	0
4	A	123	0	0	2	0
4	B	99	0	0	2	0
4	C	79	0	0	2	0
4	D	84	0	0	1	0
All	All	6861	0	6182	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:23[C]:ARG:NH2	1:D:106:ASN:OD1	2.14	0.80
1:B:70:LYS:HE3	4:B:404:HOH:O	1.87	0.73
1:A:117:ARG:HG2	1:A:117:ARG:O	1.90	0.63
1:B:111:LEU:CD2	1:B:117:ARG:HB3	2.29	0.61
1:C:76:ARG:HD2	1:D:76:ARG:HD2	1.84	0.58
1:C:66:PRO:HA	1:C:191:GLN:NE2	2.19	0.58
1:B:51:SER:HB2	1:B:121:SER:OG	2.05	0.56
1:C:69:ALA:H	1:C:133:SER:HG	1.57	0.52
1:D:12:GLN:NE2	4:D:403:HOH:O	2.42	0.52
1:C:23[A]:ARG:NH2	4:C:413:HOH:O	2.24	0.52
1:D:90:SER:HB2	1:D:185:TYR:HD1	1.75	0.51
1:C:66:PRO:HA	1:C:191:GLN:HE22	1.76	0.51
1:D:13:GLU:HG2	1:D:105:ARG:NH1	2.25	0.50
1:C:143:SER:HB2	1:C:170:ALA:HB2	1.93	0.49
1:B:88:ILE:CD1	1:B:177:GLU:HG3	2.43	0.49
1:C:59:ILE:O	1:C:143:SER:HA	2.13	0.49
1:A:88:ILE:HD11	1:A:177:GLU:HG3	1.95	0.49
4:C:471:HOH:O	1:D:26:GLU:HG2	2.12	0.49
1:D:90:SER:HB2	1:D:185:TYR:CD1	2.47	0.49
1:A:2:HIS:NE2	4:A:403:HOH:O	2.35	0.48
1:B:165:ASP:OD1	1:C:2[B]:HIS:NE2	2.44	0.48
1:C:117:ARG:C	1:C:117:ARG:HD3	2.38	0.48
1:D:1:MET:HB3	1:D:10:TRP:HB3	1.95	0.47
1:C:90:SER:HB2	1:C:185:TYR:CD1	2.49	0.47
1:D:59:ILE:O	1:D:143:SER:HA	2.15	0.47
1:C:1:MET:HE3	1:C:12:GLN:HG2	1.97	0.47
1:B:111:LEU:HD21	1:B:117:ARG:HB3	1.96	0.46
1:C:5:SER:HB2	1:C:95:ARG:HH11	1.80	0.46
1:D:117:ARG:C	1:D:117:ARG:HD3	2.41	0.46
1:A:59:ILE:O	1:A:143:SER:HA	2.17	0.45
1:C:60:HIS:O	1:C:116:GLY:HA2	2.17	0.45
1:D:51:SER:HB2	1:D:121:SER:OG	2.16	0.45
1:C:51:SER:HB3	1:C:55:ALA:HB3	1.98	0.45
1:C:143:SER:HB2	1:C:170:ALA:CB	2.47	0.44
1:D:17:HIS:O	1:D:23[A]:ARG:HA	2.17	0.44
1:C:16:ILE:CG2	1:C:23[A]:ARG:HD2	2.47	0.44
1:A:158:ASP:OD1	1:A:158:ASP:N	2.47	0.43
1:B:136:TYR:CZ	1:B:138:PRO:HG3	2.54	0.43
1:C:33:PHE:HB2	2:C:303:EDO:H11	2.00	0.43
1:B:51:SER:HB3	1:B:55:ALA:HB3	2.01	0.43
1:B:196:ARG:NH1	4:B:427:HOH:O	2.41	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:140[B]:ARG:NE	4:A:462:HOH:O	2.22	0.42
1:A:160:GLU:HA	1:A:161:PRO:HD3	1.89	0.42
1:A:60:HIS:O	1:A:116:GLY:HA2	2.20	0.42
1:D:17:HIS:O	1:D:23[C]:ARG:HA	2.19	0.42
1:A:54:GLY:O	1:A:55:ALA:C	2.62	0.42
1:A:151:LEU:HD23	1:A:151:LEU:HA	1.92	0.42
1:D:60:HIS:O	1:D:116:GLY:HA2	2.19	0.42
1:D:148:ASP:C	1:D:148:ASP:OD2	2.63	0.42
1:A:1:MET:HA	1:A:12:GLN:HE21	1.85	0.41
1:C:8:GLY:HA2	1:C:10:TRP:CZ3	2.56	0.41
1:B:1:MET:HA	1:B:12:GLN:HE21	1.85	0.41
1:B:82:VAL:O	1:B:117:ARG:HA	2.21	0.40
1:D:51:SER:HB3	1:D:55:ALA:HB3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/200 (98%)	194 (98%)	3 (2%)	0	100	100
1	B	197/200 (98%)	195 (99%)	2 (1%)	0	100	100
1	C	202/200 (101%)	198 (98%)	4 (2%)	0	100	100
1	D	202/200 (101%)	197 (98%)	5 (2%)	0	100	100
All	All	798/800 (100%)	784 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	163/164 (99%)	157 (96%)	6 (4%)	30	30
1	B	163/164 (99%)	158 (97%)	5 (3%)	35	37
1	C	168/164 (102%)	156 (93%)	12 (7%)	13	10
1	D	167/164 (102%)	158 (95%)	9 (5%)	20	17
All	All	661/656 (101%)	629 (95%)	32 (5%)	25	21

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LEU
1	A	117	ARG
1	A	120	LEU
1	A	159	ILE
1	A	166	ARG
1	A	198	GLU
1	B	104	GLU
1	B	117	ARG
1	B	120	LEU
1	B	151	LEU
1	B	178	ARG
1	C	13	GLU
1	C	23[A]	ARG
1	C	23[B]	ARG
1	C	90	SER
1	C	120	LEU
1	C	140	ARG
1	C	143	SER
1	C	151	LEU
1	C	166	ARG
1	C	193	GLN
1	C	197	LEU
1	C	198	GLU
1	D	13	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	23[A]	ARG
1	D	23[B]	ARG
1	D	23[C]	ARG
1	D	120	LEU
1	D	151	LEU
1	D	190	GLU
1	D	197	LEU
1	D	200	HIS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	12	GLN
1	A	186	GLN
1	A	192	GLN
1	B	12	GLN
1	B	192	GLN
1	B	199	HIS
1	C	25	HIS
1	C	191	GLN
1	D	12	GLN
1	D	192	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

Mogul failed to run properly - this section is therefore empty.

5.5 Carbohydrates ⓘ

Mogul failed to run properly - this section is therefore empty.

5.6 Ligand geometry ⓘ

Mogul failed to run properly - this section is therefore empty.

5.7 Other polymers [i](#)

Mogul failed to run properly - this section is therefore empty.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	198/200 (99%)	-1.51	0 100 100	7, 19, 36, 52	1 (0%)
1	B	199/200 (99%)	-1.52	0 100 100	9, 19, 35, 56	0
1	C	200/200 (100%)	-1.41	0 100 100	10, 23, 43, 65	4 (2%)
1	D	200/200 (100%)	-1.42	0 100 100	5, 24, 42, 66	3 (1%)
All	All	797/800 (99%)	-1.47	0 100 100	5, 21, 40, 66	8 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	301	4/4	0.99	0.02	16,18,21,23	0
2	EDO	A	302	4/4	0.99	0.02	18,20,21,21	0
2	EDO	B	302	4/4	0.99	0.03	16,17,20,21	0
2	EDO	C	303	4/4	0.99	0.03	20,21,22,23	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	D	301	4/4	0.99	0.04	20,22,22,23	0
2	EDO	C	301	4/4	1.00	0.02	12,16,20,25	0
2	EDO	D	302	4/4	1.00	0.03	11,15,17,23	0
3	18T	A	303	35/35	1.00	0.02	17,21,25,27	0
3	18T	B	301	35/35	1.00	0.02	12,16,21,22	0
3	18T	C	302	35/35	1.00	0.03	18,24,29,29	0
3	18T	D	303	35/35	1.00	0.02	18,23,26,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.