



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:12 AM UTC

PDB ID : 9HML / pdb_00009hml
Title : KIR2DL1 bound to RIFIN PfKE01_040007400
Authors : Chamberlain, S.G.; Higgins, M.K.
Deposited on : 2024-12-09
Resolution : 2.17 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.17 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8975 (2.20-2.16)
Clashscore	190562	9786 (2.20-2.16)
Ramachandran outliers	187476	9664 (2.20-2.16)
Sidechain outliers	187428	9664 (2.20-2.16)
RSRZ outliers	180081	8979 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	140	14% 75% 5% 19%
2	B	204	14% 78% 9% 11%
3	C	126	2% 84% 7% 6%
4	D	2	50% 50%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3380 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIFIN PfKE01_040007400.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	114	Total	C	N	O	S	0	0	0
			851	535	145	167	4			

- Molecule 2 is a protein called KIR2DL protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	181	Total	C	N	O	S	0	0	0
			1416	892	245	271	8			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	25	THR	-	expression tag	UNP A0A191URJ7
B	26	GLY	-	expression tag	UNP A0A191URJ7
B	220	GLY	-	expression tag	UNP A0A191URJ7
B	221	THR	-	expression tag	UNP A0A191URJ7
B	222	LYS	-	expression tag	UNP A0A191URJ7
B	223	HIS	-	expression tag	UNP A0A191URJ7
B	224	HIS	-	expression tag	UNP A0A191URJ7
B	225	HIS	-	expression tag	UNP A0A191URJ7
B	226	HIS	-	expression tag	UNP A0A191URJ7
B	227	HIS	-	expression tag	UNP A0A191URJ7
B	228	HIS	-	expression tag	UNP A0A191URJ7

- Molecule 3 is a protein called nanobody Nb1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	119	Total	C	N	O	S	0	0	0
			917	573	159	181	4			

- Molecule 4 is an oligosaccharide called alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	D	2	Total	C	N	O	0	0	0
			24	14	1	9			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	B	1	Total	C	N	O	0	0
			14	8	1	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	36	Total	O	0	0
			36	36		
6	B	70	Total	O	0	0
			70	70		
6	C	52	Total	O	0	0
			52	52		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: RIFIN PfKE01_040007400



- Molecule 2: KIR2DL protein



- Molecule 3: nanobody Nb1



- Molecule 4: alpha-L-fucopyranose-(1-6)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	97.17Å 97.17Å 109.20Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.05 – 2.17 28.05 – 2.17	Depositor EDS
% Data completeness (in resolution range)	100.0 (28.05-2.17) 99.9 (28.05-2.17)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.98 (at 2.18Å)	Xtriage
Refinement program	BUSTER 2.10.4	Depositor
R, R_{free}	0.221 , 0.232 0.214 , 0.234	Depositor DCC
R_{free} test set	1592 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	57.6	Xtriage
Anisotropy	0.078	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 52.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.026 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3380	wwPDB-VP
Average B, all atoms (Å ²)	72.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, FUC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.75	1/864 (0.1%)	1.05	0/1172
2	B	0.69	0/1454	1.01	3/1969 (0.2%)
3	C	0.78	1/939 (0.1%)	1.09	3/1271 (0.2%)
All	All	0.73	2/3257 (0.1%)	1.04	6/4412 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	34	MET	SD-CE	-7.16	1.61	1.79
1	A	101	MET	SD-CE	-6.50	1.63	1.79

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	63	VAL	N-CA-CB	-8.24	103.14	112.21
3	C	48	VAL	N-CA-CB	7.83	118.06	111.64
3	C	67	PHE	CA-CB-CG	5.61	119.41	113.80
2	B	122	ILE	N-CA-CB	5.31	116.51	110.72
2	B	165	GLU	CA-C-N	5.20	127.96	120.38
2	B	165	GLU	C-N-CA	5.20	127.96	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	851	0	857	4	0
2	B	1416	0	1345	10	0
3	C	917	0	867	8	0
4	D	24	0	22	0	0
5	B	14	0	13	0	0
6	A	36	0	0	0	0
6	B	70	0	0	2	0
6	C	52	0	0	0	0
All	All	3380	0	3104	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:32:TYR:CB	3:C:34:MET:HE2	2.21	0.70
2:B:121:VAL:HG11	2:B:209:TRP:CD2	2.31	0.65
3:C:32:TYR:HB3	3:C:34:MET:HE2	1.79	0.64
2:B:92:GLN:HA	2:B:122:ILE:HD11	1.82	0.60
1:A:80:ALA:HB1	1:A:101:MET:HE1	1.83	0.60
3:C:32:TYR:HB2	3:C:34:MET:HE2	1.83	0.59
3:C:90:THR:HG23	3:C:116:THR:HA	1.85	0.58
2:B:150:SER:HB3	2:B:182:GLN:HG3	1.87	0.56
2:B:107:SER:HB2	2:B:108:PRO:HD3	1.89	0.55
2:B:89:ARG:NH1	6:B:502:HOH:O	2.43	0.51
2:B:124:GLY:HA2	2:B:211:LYS:HA	1.95	0.48
3:C:63:VAL:HG13	3:C:67:PHE:HB2	1.94	0.48
2:B:133:ALA:HB2	2:B:216:LEU:HD21	1.96	0.47
3:C:32:TYR:HB2	3:C:34:MET:CE	2.46	0.45
1:A:89:LYS:HB2	1:A:94:ILE:HG22	1.99	0.45
1:A:18:ALA:HB2	1:A:120:ALA:HB2	1.98	0.44
2:B:128:LYS:NZ	6:B:509:HOH:O	2.53	0.42
1:A:63:SER:HB3	2:B:61:HIS:CD2	2.55	0.42
3:C:87:PRO:HA	3:C:117:VAL:HB	2.01	0.42
3:C:18:LEU:HB3	3:C:82:MET:HE3	2.03	0.41
2:B:126:TYR:H	2:B:210:SER:HA	1.87	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	112/140 (80%)	110 (98%)	2 (2%)	0	100	100
2	B	175/204 (86%)	172 (98%)	3 (2%)	0	100	100
3	C	117/126 (93%)	115 (98%)	2 (2%)	0	100	100
All	All	404/470 (86%)	397 (98%)	7 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	92/106 (87%)	89 (97%)	3 (3%)	33	42
2	B	158/176 (90%)	153 (97%)	5 (3%)	34	43
3	C	96/102 (94%)	92 (96%)	4 (4%)	26	32
All	All	346/384 (90%)	334 (96%)	12 (4%)	32	40

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	GLN
1	A	66	LEU
1	A	94	ILE
2	B	32	LEU
2	B	56	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	80	VAL
2	B	122	ILE
2	B	216	LEU
3	C	4	LEU
3	C	48	VAL
3	C	63	VAL
3	C	78	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	61	HIS
3	C	39	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NAG	D	1	2,4	14,14,15	0.34	0	17,19,21	0.78	1 (5%)
4	FUC	D	2	4	10,10,11	0.51	0	14,14,16	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	2,4	-	2/6/23/26	0/1/1/1
4	FUC	D	2	4	-	-	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	D	1	NAG	C1-O5-C5	2.18	115.11	112.19

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	D	1	NAG	C4-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	NAG	B	400	2	14,14,15	0.29	0	17,19,21	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	B	400	2	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	114/140 (81%)	0.97	20 (17%) 4 3	51, 76, 122, 136	0
2	B	181/204 (88%)	0.87	29 (16%) 5 4	47, 71, 120, 135	0
3	C	119/126 (94%)	0.31	2 (1%) 69 68	46, 59, 84, 104	0
All	All	414/470 (88%)	0.73	51 (12%) 8 7	46, 68, 117, 136	0

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	177	VAL	6.4
1	A	9	ALA	5.5
2	B	188	GLY	5.3
2	B	141	ALA	5.0
1	A	41	PHE	4.0
2	B	179	GLY	3.9
2	B	133	ALA	3.8
2	B	27	ARG	3.8
3	C	2	ARG	3.8
2	B	167	HIS	3.6
1	A	120	ALA	3.6
2	B	147	LEU	3.5
1	A	10	ALA	3.4
2	B	149	CYS	3.4
2	B	219	SER	3.3
3	C	1	GLN	3.2
2	B	108	PRO	3.2
2	B	158	TYR	3.0
1	A	43	ALA	3.0
2	B	32	LEU	2.9
2	B	146	THR	2.9
2	B	106	HIS	2.9
1	A	94	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	178	ASN	2.8
2	B	105	THR	2.7
1	A	122	THR	2.7
1	A	91	PRO	2.7
2	B	50	TRP	2.6
1	A	118	LYS	2.6
1	A	93	PHE	2.5
2	B	142	GLY	2.5
2	B	104	VAL	2.5
1	A	70	SER	2.5
2	B	176	LYS	2.5
1	A	88	ARG	2.4
2	B	107	SER	2.4
1	A	38	LYS	2.4
1	A	119	THR	2.3
2	B	175	PRO	2.3
1	A	67	THR	2.3
2	B	183	ALA	2.2
2	B	185	PHE	2.2
2	B	122	ILE	2.1
1	A	37	LEU	2.1
2	B	180	THR	2.1
1	A	90	ILE	2.1
2	B	181	PHE	2.1
1	A	34	VAL	2.1
1	A	46	TYR	2.0
2	B	109	TYR	2.0
1	A	12	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	FUC	D	2	10/11	0.87	0.19	76,78,79,79	0
4	NAG	D	1	14/15	0.92	0.10	62,65,70,73	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	NAG	B	400	14/15	0.89	0.10	70,71,72,72	0

6.5 Other polymers [i](#)

There are no such residues in this entry.