



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 04:46 PM UTC

PDB ID : 4HN4 / pdb_00004hn4
Title : Tryptophan synthase in complex with alpha aminoacrylate E(A-A) form and the F9 inhibitor in the alpha site
Authors : Hilario, E.; Niks, D.; Dunn, M.F.; Mueller, L.J.; Fan, L.
Deposited on : 2012-10-18
Resolution : 1.64 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.


The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49

ene]amino}prop-2-enoic acid (CCD ID: 0JO) (formula: $C_{11}H_{13}N_2O_7P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	B	1	Total	C	N	O	P	0	0
			21	11	2	7	1		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			7	4	3		
5	B	1	Total	C	O	0	0
			7	4	3		

Continued on next page...

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.97Å 59.72Å 67.40Å 90.00° 94.69° 90.00°	Depositor
Resolution (Å)	28.59 – 1.64 28.59 – 1.64	Depositor EDS
% Data completeness (in resolution range)	95.7 (28.59-1.64) 95.7 (28.59-1.64)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.04 (at 1.64Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.148 , 0.200 0.144 , 0.197	Depositor DCC
R_{free} test set	4281 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	17.4	Xtriage
Anisotropy	0.651	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5908	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.33% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	22	0	9	1	0
4	B	21	0	9	0	0
5	B	21	0	30	2	0
6	B	33	0	36	0	0
7	B	3	0	0	1	0
8	A	248	0	0	1	0
8	B	428	0	0	6	0
All	All	5908	0	5271	33	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (33) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:408:CS:CS	8:B:901:HOH:O	2.04	1.03
1:A:192:PRO:HG2	1:A:223:ALA:HA	1.59	0.82
5:B:403:PEG:H32	8:B:884:HOH:O	1.90	0.71
1:A:112[A]:ASP:OD1	1:A:146:HIS:HE1	1.79	0.65
1:A:221:SER:HB3	1:A:225[A]:ARG:HH12	1.64	0.63
1:A:192:PRO:CG	1:A:223:ALA:HA	2.30	0.60
2:B:301[A]:SER:OG	2:B:350:GLU:HG3	2.04	0.57
1:A:221:SER:HB3	1:A:225[A]:ARG:NH1	2.20	0.56
1:A:211:GLY:O	1:A:212:PHE:HB2	2.08	0.53
1:A:58[B]:LEU:HD12	2:B:174:LEU:HD23	1.91	0.51
2:B:37[A]:LYS:HE3	8:B:655:HOH:O	2.13	0.49
1:A:108:ASN:HA	8:A:596:HOH:O	2.12	0.49
1:A:178:SER:HB3	1:A:189:GLY:HA3	1.95	0.49
2:B:16:TYR:O	2:B:281:GLY:HA2	2.14	0.47
1:A:22:PHE:HA	1:A:49:GLU:O	2.15	0.46
2:B:222:ARG:NH1	8:B:904:HOH:O	2.26	0.46
5:B:403:PEG:H41	8:B:798:HOH:O	2.16	0.46
2:B:216:ILE:HG21	2:B:224:PRO:HD3	1.98	0.45
1:A:22:PHE:CD1	1:A:22:PHE:C	2.95	0.45
2:B:44[B]:GLN:HG3	8:B:601:HOH:O	2.17	0.44
2:B:11:GLU:O	2:B:11:GLU:HG2	2.17	0.44
2:B:392:LYS:HB3	2:B:392:LYS:HE3	1.68	0.43
1:A:215[B]:SER:HB3	1:A:236:ALA:HB2	2.01	0.43
1:A:20:VAL:HG22	1:A:47:ALA:HB3	2.01	0.43
2:B:286[A]:MET:HE2	2:B:286[A]:MET:HB3	1.94	0.43
2:B:59:LEU:O	2:B:219[C]:LYS:NZ	2.48	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:337:LYS:O	2:B:341[B]:ARG:HG3	2.20	0.42
2:B:300:ILE:HD11	2:B:390:ILE:CD1	2.49	0.42
2:B:295[B]:GLU:OE2	2:B:295[B]:GLU:HA	2.19	0.41
2:B:146:VAL:HA	2:B:149:MET:HE3	2.03	0.41
1:A:175:TYR:CE2	3:A:301:F9F:H3	2.55	0.41
2:B:99[A]:LYS:HD2	2:B:126:LEU:HB3	2.03	0.41
1:A:178:SER:HB3	1:A:189:GLY:CA	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/268 (102%)	266 (98%)	5 (2%)	1 (0%)	30	14
2	B	409/397 (103%)	401 (98%)	8 (2%)	0	100	100
All	All	681/665 (102%)	667 (98%)	13 (2%)	1 (0%)	43	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	212	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	214/208 (103%)	214 (100%)	0	100	100
2	B	325/311 (104%)	323 (99%)	2 (1%)	78	66
All	All	539/519 (104%)	537 (100%)	2 (0%)	84	76

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	30	GLU
2	B	392	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	HIS
1	A	157	ASN
1	A	195	HIS
2	B	26	ASN
2	B	51	ASN
2	B	317	ASN
2	B	342	HIS
2	B	365	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 3 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	PEG	B	404	-	6,6,6	0.62	0	5,5,5	0.29	0
5	PEG	B	402	-	6,6,6	0.48	0	5,5,5	0.24	0
5	PEG	B	403	-	6,6,6	0.48	0	5,5,5	0.50	0
3	F9F	A	301	-	22,22,22	1.90	2 (9%)	32,33,33	1.88	3 (9%)
6	BCN	B	407	-	10,10,10	1.03	1 (10%)	11,11,11	0.78	1 (9%)
4	OJO	B	401	-	20,21,21	2.44	2 (10%)	23,30,30	1.91	7 (30%)
6	BCN	B	406	-	10,10,10	0.87	0	11,11,11	0.73	0
6	BCN	B	405	-	10,10,10	0.80	0	11,11,11	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	PEG	B	404	-	-	3/4/4/4	-
5	PEG	B	402	-	-	3/4/4/4	-
5	PEG	B	403	-	-	2/4/4/4	-
3	F9F	A	301	-	-	6/20/20/20	0/1/1/1
6	BCN	B	407	-	-	2/10/10/10	-
4	OJO	B	401	-	-	0/11/15/15	0/1/1/1
6	BCN	B	406	-	-	2/10/10/10	-
6	BCN	B	405	-	-	5/10/10/10	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	401	OJO	C4A-N	9.93	1.41	1.28
3	A	301	F9F	C4-S12	-7.74	1.64	1.76
6	B	407	BCN	O21-C2	2.43	1.30	1.22
4	B	401	OJO	OXT-C	-2.06	1.25	1.30
3	A	301	F9F	O7-C8	2.00	1.42	1.31

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	301	F9F	O22-S12-O21	-8.01	109.79	119.52
4	B	401	OJO	C4-C4A-N	-4.17	113.38	123.01
4	B	401	OJO	OP4-C5A-C5	3.95	116.76	109.36
4	B	401	OJO	C5-C4-C4A	-3.18	116.56	121.47
3	A	301	F9F	C6-C5-C4	2.92	122.28	119.44
3	A	301	F9F	O21-S12-N13	2.90	111.55	107.03
4	B	401	OJO	C6-N1-C2	2.48	123.70	119.20
4	B	401	OJO	C3-C4-C4A	2.33	124.60	120.40
4	B	401	OJO	C3-C2-N1	-2.24	118.14	120.96
6	B	407	BCN	C1-N1-C3	-2.13	106.77	111.91
4	B	401	OJO	C2A-C2-N1	2.01	121.42	117.64

There are no chirality outliers.

All (23) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
5	B	403	PEG	O2-C3-C4-O4
6	B	405	BCN	N1-C3-C4-O4
3	A	301	F9F	C5-C4-S12-O21
3	A	301	F9F	C3-C4-S12-O21
5	B	402	PEG	O2-C3-C4-O4
5	B	403	PEG	O1-C1-C2-O2
6	B	405	BCN	C2-C1-N1-C3
5	B	404	PEG	O2-C3-C4-O4
3	A	301	F9F	N13-C14-C15-O16
5	B	402	PEG	C4-C3-O2-C2
5	B	404	PEG	O1-C1-C2-O2
6	B	405	BCN	C2-C1-N1-C5
3	A	301	F9F	C2-C1-O7-C8
6	B	407	BCN	C2-C1-N1-C5
5	B	404	PEG	C1-C2-O2-C3
5	B	402	PEG	C1-C2-O2-C3
3	A	301	F9F	C6-C1-O7-C8
6	B	405	BCN	C4-C3-N1-C5
3	A	301	F9F	C5-C4-S12-N13
6	B	406	BCN	C2-C1-N1-C3
6	B	406	BCN	N1-C5-C6-O6
6	B	405	BCN	C4-C3-N1-C1
6	B	407	BCN	C2-C1-N1-C3


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	403	PEG	2	0
3	A	301	F9F	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/268 (100%)	-0.26	6 (2%) 62 68	13, 21, 42, 65	6 (2%)
2	B	395/397 (99%)	-0.64	1 (0%) 90 91	6, 15, 25, 57	14 (3%)
All	All	663/665 (99%)	-0.49	7 (1%) 78 83	6, 18, 36, 65	20 (3%)

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	191	LEU	4.3
1	A	268	ALA	3.3
1	A	190	ALA	3.2
1	A	192	PRO	3.1
1	A	193	LEU	3.1
1	A	189	GLY	2.9
2	B	63	GLN	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	PEG	B	403	7/7	0.79	0.15	29,34,44,57	0
5	PEG	B	402	7/7	0.87	0.10	32,40,46,55	0
5	PEG	B	404	7/7	0.88	0.10	30,31,40,40	0
6	BCN	B	405	11/11	0.93	0.09	21,31,35,37	0
6	BCN	B	406	11/11	0.95	0.07	16,24,29,32	0
6	BCN	B	407	11/11	0.95	0.08	15,25,32,37	0
3	F9F	A	301	22/22	0.99	0.04	15,19,22,23	0
4	OJO	B	401	21/21	0.99	0.04	10,12,15,17	0
7	CS	B	408	1/1	0.99	0.03	25,25,25,25	1
7	CS	B	409[A]	1/1	1.00	0.02	16,16,16,16	1
7	CS	B	409[B]	1/1	1.00	0.02	17,17,17,17	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around F9F A 301:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 0JO B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.