



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 20, 2026 – 07:49 PM UTC

PDB ID : 4HNX / pdb_00004hnx
Title : The NatA Acetyltransferase Complex Bound To ppGpp
Authors : Neubauer, J.L.; Immormino, R.M.; Dollins, D.E.; Endo-Streeter, S.T.; Pemble IV, C.W.; York, J.D.
Deposited on : 2012-10-21
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	863	13% 74% 8% 17%
2	B	248	15% 74% 12% 14%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N-terminal acetyltransferase A complex subunit NAT1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	713	Total	C	N	O	S	1	1	0
			5541	3568	927	1024	22			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	855	TYR	-	expression tag	UNP P12945
A	856	PRO	-	expression tag	UNP P12945
A	857	TYR	-	expression tag	UNP P12945
A	858	ASP	-	expression tag	UNP P12945
A	859	VAL	-	expression tag	UNP P12945
A	860	PRO	-	expression tag	UNP P12945
A	861	ASP	-	expression tag	UNP P12945
A	862	TYR	-	expression tag	UNP P12945
A	863	ALA	-	expression tag	UNP P12945

- Molecule 2 is a protein called N-terminal acetyltransferase A complex catalytic subunit ARD1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	213	Total	C	N	O	S	0	0	0
			1646	1044	276	315	11			

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	239	GLU	-	expression tag	UNP P07347
B	240	GLN	-	expression tag	UNP P07347
B	241	LYS	-	expression tag	UNP P07347
B	242	LEU	-	expression tag	UNP P07347
B	243	ILE	-	expression tag	UNP P07347
B	244	SER	-	expression tag	UNP P07347

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	245	GLU	-	expression tag	UNP P07347
B	246	GLU	-	expression tag	UNP P07347
B	247	ASP	-	expression tag	UNP P07347
B	248	LEU	-	expression tag	UNP P07347

- Molecule 3 is GUANOSINE-5',3'-TETRAPHOSPHATE (CCD ID: G4P) (formula: $C_{10}H_{17}N_5O_{17}P_4$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			36	10	5	17	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	139	Total	O	0	0
			139	139		
4	B	46	Total	O	0	0
			46	46		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: N-terminal acetyltransferase A complex subunit NAT1



- Molecule 2: N-terminal acetyltransferase A complex catalytic subunit ARD1



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	137.10Å 137.10Å 179.60Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.14 – 2.34 40.14 – 2.34	Depositor EDS
% Data completeness (in resolution range)	99.2 (40.14-2.34) 93.4 (40.14-2.34)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.34Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.6.4_486)	Depositor
R, R_{free}	0.213 , 0.240 (Not available) , 0.238	Depositor DCC
R_{free} test set	4110 reflections (4.96%)	wwPDB-VP
Wilson B-factor (Å ²)	48.5	Xtriage
Anisotropy	0.150	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 63.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.021 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	7408	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: G4P

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.49	0/5662	0.75	4/7674 (0.1%)
2	B	0.47	0/1676	0.78	2/2279 (0.1%)
All	All	0.48	0/7338	0.76	6/9953 (0.1%)

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	82	ASP	CA-C-N	5.69	125.40	119.82
2	B	82	ASP	C-N-CA	5.69	125.40	119.82
1	A	601	LYS	CA-C-N	5.47	124.93	119.24
1	A	601	LYS	C-N-CA	5.47	124.93	119.24
1	A	810	ILE	N-CA-C	5.46	115.66	110.42
1	A	534	GLU	N-CA-C	-5.20	107.31	113.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5541	0	5207	47	0
2	B	1646	0	1553	19	0
3	A	36	0	10	3	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	139	0	0	0	0
4	B	46	0	0	1	0
All	All	7408	0	6770	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:111:ILE:O	1:A:115:THR:HG23	1.86	0.75
1:A:314:ILE:O	1:A:317:THR:HB	1.88	0.73
1:A:164:SER:HB3	1:A:580:MET:HG3	1.73	0.69
1:A:86:GLY:C	1:A:88:SER:H	2.06	0.62
2:B:150:VAL:HG23	2:B:194:LEU:HD11	1.83	0.60
2:B:225:LEU:O	2:B:228:ILE:HG13	2.01	0.59
1:A:80:ALA:O	1:A:84:ILE:HG13	2.01	0.59
1:A:532:LYS:C	1:A:534:GLU:H	2.09	0.58
1:A:54:VAL:O	1:A:57:LEU:HB2	2.04	0.58
1:A:441:LEU:HD21	1:A:457:LYS:HG2	1.87	0.56
1:A:55:ASP:HB3	1:A:93:CYS:HB2	1.88	0.56
1:A:476:SER:O	1:A:479:THR:HB	2.07	0.55
1:A:84:ILE:O	1:A:84:ILE:HG22	2.07	0.55
1:A:543:ASN:O	1:A:547:VAL:HG23	2.10	0.52
2:B:152:LEU:C	2:B:152:LEU:HD12	2.35	0.51
1:A:559:ASN:O	1:A:562:PRO:HD2	2.10	0.51
2:B:178:LYS:HD2	2:B:186:ASP:OD1	2.11	0.51
1:A:19:LYS:O	1:A:23:GLN:HG3	2.11	0.50
1:A:159:ARG:HD2	1:A:185:PHE:CE2	2.46	0.50
1:A:40:LEU:HG	1:A:63:ASP:HB3	1.92	0.50
1:A:381:ILE:HB	1:A:382:PRO:HD3	1.94	0.49
2:B:227:ASP:HB3	2:B:239:GLU:OE2	2.13	0.49
1:A:831:LYS:O	1:A:835:ILE:HG13	2.13	0.49
2:B:181:TYR:CE2	2:B:187:ALA:HB2	2.48	0.48
2:B:20:ASN:HB2	4:B:339:HOH:O	2.13	0.48
2:B:87:ALA:HB1	2:B:88:PRO:HD2	1.95	0.48
1:A:854:LEU:O	1:A:855:TYR:C	2.55	0.48
1:A:86:GLY:C	1:A:88:SER:N	2.71	0.48
2:B:12:ASP:HA	2:B:86:LEU:HD21	1.95	0.48
1:A:696:ALA:HA	1:A:700:TYR:HB3	1.96	0.48
2:B:22:HIS:ND1	2:B:29:MET:HE2	2.29	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:GLU:HB3	1:A:235:PRO:HD3	1.97	0.47
1:A:309:GLU:HB2	1:A:310:PRO:HD3	1.97	0.47
1:A:829:ASN:O	1:A:833:MET:HG3	2.15	0.46
1:A:429:LYS:NZ	3:A:901:G4P:O3D	2.47	0.46
1:A:82:ARG:C	1:A:84:ILE:H	2.24	0.46
1:A:42:LEU:O	1:A:46:ILE:HG13	2.16	0.46
1:A:241:PHE:HB2	2:B:7:ARG:NH2	2.31	0.46
1:A:532:LYS:C	1:A:534:GLU:N	2.73	0.45
1:A:548:ARG:HH22	1:A:687:GLU:CD	2.24	0.45
1:A:430:HIS:NE2	3:A:901:G4P:O1B	2.46	0.44
2:B:82:ASP:OD1	2:B:82:ASP:C	2.60	0.44
1:A:92:ILE:O	1:A:96:VAL:HG23	2.18	0.44
2:B:12:ASP:O	2:B:16:MET:HG3	2.19	0.43
1:A:606:ARG:O	1:A:610:GLU:HG2	2.19	0.43
1:A:462:PHE:CE1	1:A:470:LYS:HG2	2.54	0.43
2:B:228:ILE:HG13	2:B:228:ILE:H	1.60	0.42
2:B:156:GLN:HA	2:B:188:TYR:CE2	2.54	0.42
1:A:389:TYR:CD1	1:A:389:TYR:C	2.97	0.42
1:A:459:VAL:HG21	1:A:475:ALA:HB2	2.02	0.42
1:A:839:SER:N	1:A:840:PRO:HD2	2.35	0.41
2:B:229:ILE:HD12	2:B:229:ILE:HA	1.91	0.41
1:A:407:ILE:HD12	1:A:407:ILE:HA	1.83	0.41
1:A:65:TYR:CD1	1:A:65:TYR:C	2.98	0.41
1:A:571:ASP:OD2	2:B:27:ASN:ND2	2.46	0.41
1:A:514:ARG:HA	1:A:514:ARG:HD3	1.90	0.41
1:A:558:PHE:CE1	1:A:610:GLU:HB2	2.56	0.41
1:A:697:THR:HG22	1:A:698:GLU:OE2	2.21	0.41
3:A:901:G4P:N2	2:B:81:LEU:HB3	2.36	0.41
1:A:159:ARG:HB2	1:A:185:PHE:CZ	2.55	0.41
2:B:237:ILE:O	2:B:240:GLN:HB2	2.21	0.40
1:A:159:ARG:CD	1:A:185:PHE:CE2	3.04	0.40
1:A:546:LEU:HD23	1:A:546:LEU:HA	1.91	0.40
1:A:601:LYS:HA	1:A:602:PRO:HD3	1.91	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	702/863 (81%)	678 (97%)	24 (3%)	0	100	100
2	B	207/248 (84%)	200 (97%)	7 (3%)	0	100	100
All	All	909/1111 (82%)	878 (97%)	31 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	538/767 (70%)	532 (99%)	6 (1%)	65	77
2	B	170/226 (75%)	168 (99%)	2 (1%)	63	75
All	All	708/993 (71%)	700 (99%)	8 (1%)	65	77

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	164	SER
1	A	317	THR
1	A	407	ILE
1	A	521	LEU
1	A	577	SER
1	A	841	LEU
2	B	11	ASN
2	B	157	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	GLN
1	A	184	GLN
1	A	256	GLN
1	A	392	GLN
1	A	559	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	G4P	A	901	-	36,38,38	2.65	14 (38%)	56,61,61	2.05	13 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	G4P	A	901	-	-	8/27/43/43	0/3/3/3

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	901	G4P	O6-C6	7.74	1.38	1.23
3	A	901	G4P	PA-O3A	-6.06	1.53	1.59
3	A	901	G4P	C2-N2	4.86	1.45	1.34
3	A	901	G4P	PC-O3C	4.58	1.64	1.59
3	A	901	G4P	C8-N9	-3.67	1.29	1.37
3	A	901	G4P	O2'-C2'	-3.44	1.34	1.43
3	A	901	G4P	C5-N7	-3.33	1.32	1.39
3	A	901	G4P	C8-N7	-3.22	1.23	1.32
3	A	901	G4P	C2'-C3'	-3.21	1.46	1.53
3	A	901	G4P	C5-C4	2.85	1.46	1.38
3	A	901	G4P	PD-O2D	-2.32	1.46	1.54
3	A	901	G4P	C1'-N9	-2.31	1.41	1.47
3	A	901	G4P	C4-N9	2.13	1.43	1.38
3	A	901	G4P	C5'-C4'	-2.06	1.45	1.51

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	901	G4P	N9-C4-N3	5.40	136.75	125.95
3	A	901	G4P	C5-C4-N3	-5.30	119.96	128.39
3	A	901	G4P	C2-N3-C4	5.27	121.38	112.30
3	A	901	G4P	C6-C5-N7	5.20	139.75	130.29
3	A	901	G4P	C8-N7-C5	5.09	113.33	104.26
3	A	901	G4P	C4-C5-N7	-4.34	103.79	110.67
3	A	901	G4P	C5-C6-N1	2.91	120.67	113.25
3	A	901	G4P	O6-C6-C5	-2.70	119.42	126.53
3	A	901	G4P	O2C-PC-O3C	2.42	113.81	107.27
3	A	901	G4P	C2-N1-C6	-2.19	121.13	125.11
3	A	901	G4P	O2A-PA-O3A	2.14	113.05	107.27
3	A	901	G4P	O4'-C1'-N9	2.08	113.07	108.36
3	A	901	G4P	O2B-PB-O3A	2.05	111.52	104.64

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	901	G4P	PA-O3A-PB-O2B

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	A	901	G4P	C3'-O3'-PC-O3C
3	A	901	G4P	C3'-C4'-C5'-O5'
3	A	901	G4P	PB-O3A-PA-O5'
3	A	901	G4P	O4'-C4'-C5'-O5'
3	A	901	G4P	C3'-O3'-PC-O1C
3	A	901	G4P	PA-O3A-PB-O3B
3	A	901	G4P	C3'-O3'-PC-O2C

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	901	G4P	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	713/863 (82%)	0.68	111 (15%) 5 5	27, 58, 122, 168	1 (0%)
2	B	213/248 (85%)	0.74	37 (17%) 4 4	33, 61, 109, 126	0
All	All	926/1111 (83%)	0.69	148 (15%) 5 5	27, 59, 118, 168	1 (0%)

All (148) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	623	LEU	9.4
2	B	108	ASN	7.5
1	A	808	ASN	7.5
1	A	854	LEU	7.4
2	B	245	GLU	6.9
1	A	87	ALA	6.5
1	A	855	TYR	6.3
1	A	530	SER	6.0
2	B	79	ILE	5.9
2	B	207	ARG	5.9
1	A	85	GLU	5.8
1	A	51	GLY	5.6
2	B	106	GLN	5.5
1	A	88	SER	5.5
1	A	821	ASP	5.3
1	A	853	TYR	5.3
1	A	825	ILE	5.2
1	A	531	ASP	5.1
2	B	182	GLN	5.0
1	A	494	VAL	5.0
2	B	176	ILE	4.9
1	A	621	ASP	4.9
1	A	747	LEU	4.9
1	A	529	GLU	4.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	756	LEU	4.8
1	A	824	PRO	4.8
1	A	743	THR	4.5
1	A	536	ILE	4.5
2	B	243	ILE	4.5
1	A	538	ASN	4.5
1	A	532	LYS	4.5
1	A	21	ASN	4.4
1	A	848	GLU	4.3
1	A	533	SER	4.3
1	A	758	ALA	4.3
1	A	479	THR	4.3
1	A	18	LYS	4.2
1	A	839	SER	4.2
2	B	103	ASP	4.1
1	A	748	PHE	4.0
2	B	107	GLN	4.0
2	B	54	CYS	3.8
1	A	823	VAL	3.8
1	A	744	THR	3.8
1	A	809	ASP	3.7
2	B	23	ASN	3.7
1	A	852	TYR	3.7
1	A	829	ASN	3.7
1	A	19	LYS	3.7
1	A	850	LEU	3.6
1	A	757	HIS	3.6
2	B	216	LEU	3.5
2	B	238	VAL	3.5
1	A	735	ASN	3.5
1	A	55	ASP	3.5
1	A	41	LYS	3.4
1	A	750	SER	3.4
1	A	540	ILE	3.3
1	A	827	SER	3.3
1	A	826	GLY	3.3
1	A	617	GLN	3.2
1	A	838	LEU	3.2
2	B	229	ILE	3.2
1	A	828	SER	3.2
1	A	813	LEU	3.2
1	A	49	LYS	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	522	ALA	3.1
1	A	622	ARG	3.1
1	A	810	ILE	3.1
2	B	179	SER	3.1
2	B	241	LYS	3.1
1	A	52	SER	3.0
1	A	44	ASP	3.0
1	A	836	SER	3.0
1	A	745	SER	3.0
1	A	38	LYS	3.0
1	A	837	SER	3.0
1	A	86	GLY	2.9
1	A	534	GLU	2.9
1	A	705	MET	2.9
2	B	178	LYS	2.9
2	B	173	VAL	2.9
1	A	196	SER	2.8
1	A	289	GLY	2.8
1	A	48	LYS	2.8
1	A	724	ILE	2.8
2	B	226	GLU	2.8
1	A	619	HIS	2.7
2	B	244	SER	2.7
2	B	80	LYS	2.7
1	A	141	PHE	2.7
2	B	215	LYS	2.7
1	A	82	ARG	2.6
1	A	686	ILE	2.6
1	A	696	ALA	2.6
1	A	728	ALA	2.6
1	A	476	SER	2.6
1	A	478	PHE	2.5
1	A	537	ALA	2.5
1	A	851	GLN	2.5
2	B	188	TYR	2.5
1	A	689	SER	2.5
1	A	741	PHE	2.5
1	A	30	LEU	2.5
2	B	184	GLY	2.5
1	A	679	ASP	2.4
1	A	693	GLU	2.4
1	A	832	GLU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	242	LEU	2.4
2	B	197	GLU	2.4
1	A	73	ALA	2.4
1	A	740	ARG	2.4
1	A	819	TYR	2.4
1	A	754	VAL	2.4
1	A	521	LEU	2.4
2	B	110	PRO	2.3
2	B	187	ALA	2.3
1	A	78	ALA	2.3
2	B	109	GLU	2.3
1	A	708	ARG	2.3
2	B	157	SER	2.3
1	A	535	GLN	2.3
1	A	731	PHE	2.2
1	A	218	ASP	2.2
1	A	725	GLY	2.2
1	A	844	HIS	2.2
1	A	515	LYS	2.2
1	A	845	SER	2.2
2	B	181	TYR	2.2
1	A	746	GLY	2.2
1	A	53	HIS	2.2
1	A	517	LYS	2.2
1	A	23	GLN	2.2
1	A	722	TYR	2.1
1	A	232	ASP	2.1
2	B	185	GLU	2.1
2	B	206	HIS	2.1
1	A	742	GLY	2.1
2	B	21	LEU	2.1
2	B	20	ASN	2.1
1	A	812	GLY	2.1
1	A	752	ALA	2.1
1	A	477	LEU	2.1
1	A	45	ALA	2.0
1	A	35	GLN	2.0
1	A	755	LEU	2.0
2	B	186	ASP	2.0
1	A	849	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	G4P	A	901	36/36	0.90	0.14	61,87,98,100	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around G4P A 901:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.