



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 12:29 PM UTC

PDB ID : 3HOX / pdb_00003hox
Title : Complete RNA polymerase II elongation complex V
Authors : Sydow, J.F.; Brueckner, F.; Cheung, A.C.M.; Damsma, G.E.; Dengl, S.;
Lehmann, E.; Vassylyev, D.; Cramer, P.
Deposited on : 2009-06-03
Resolution : 3.65 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1062 (3.78-3.54)
Clashscore	190562	1009 (3.76-3.56)
Ramachandran outliers	187476	1054 (3.78-3.54)
Sidechain outliers	187428	1052 (3.78-3.54)
RSRZ outliers	180081	1061 (3.78-3.54)
RNA backbone	3983	1027 (4.26-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1733	2% 18% 46% 15% • 18%
2	B	1224	4% 17% 53% 18% • 9%
3	C	347	12% 46% 16% • 23%
4	D	221	5% 17% 46% 17% • 19%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	E	215	
6	F	155	
7	G	171	
8	H	146	
9	I	122	
10	J	70	
11	K	120	
12	L	70	
13	N	12	
14	T	26	
15	P	18	

2 Entry composition [i](#)

There are 17 unique types of molecules in this entry. The entry contains 31918 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1417	Total	C	N	O	S	0	0	0
			11151	7027	1950	2112	62			

- Molecule 2 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	1113	Total	C	N	O	S	0	0	0
			8847	5600	1552	1640	55			

- Molecule 3 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	266	Total	C	N	O	S	0	0	0
			2095	1317	348	417	13			

There are 30 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-28	MET	-	expression tag	UNP P16370
C	-27	GLY	-	expression tag	UNP P16370
C	-26	SER	-	expression tag	UNP P16370
C	-25	HIS	-	expression tag	UNP P16370
C	-24	HIS	-	expression tag	UNP P16370
C	-23	HIS	-	expression tag	UNP P16370
C	-22	HIS	-	expression tag	UNP P16370
C	-21	HIS	-	expression tag	UNP P16370
C	-20	HIS	-	expression tag	UNP P16370
C	-19	SER	-	expression tag	UNP P16370
C	-18	ASN	-	expression tag	UNP P16370
C	-17	SER	-	expression tag	UNP P16370
C	-16	GLY	-	expression tag	UNP P16370
C	-15	LEU	-	expression tag	UNP P16370

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-14	ASN	-	expression tag	UNP P16370
C	-13	ASP	-	expression tag	UNP P16370
C	-12	ILE	-	expression tag	UNP P16370
C	-11	PHE	-	expression tag	UNP P16370
C	-10	GLU	-	expression tag	UNP P16370
C	-9	ALA	-	expression tag	UNP P16370
C	-8	GLN	-	expression tag	UNP P16370
C	-7	LYS	-	expression tag	UNP P16370
C	-6	ILE	-	expression tag	UNP P16370
C	-5	GLU	-	expression tag	UNP P16370
C	-4	TRP	-	expression tag	UNP P16370
C	-3	HIS	-	expression tag	UNP P16370
C	-2	GLU	-	expression tag	UNP P16370
C	-1	ASP	-	expression tag	UNP P16370
C	0	THR	-	expression tag	UNP P16370
C	1	GLY	-	expression tag	UNP P16370

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	178	Total	C	N	O	S	0	0	0
			1434	887	257	288	2			

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	214	Total	C	N	O	S	0	0	0
			1752	1111	309	321	11			

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	87	Total	C	N	O	S	0	0	0
			705	451	119	132	3			

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	G	171	Total	C	N	O	S	0	0	0
			1340	861	222	249	8			

- Molecule 8 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	H	135	Total	C	N	O	S	0	0	0
			1081	680	183	214	4			

- Molecule 9 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	I	119	Total	C	N	O	S	0	0	0
			971	596	179	186	10			

- Molecule 10 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	J	65	Total	C	N	O	S	0	0	0
			532	339	93	94	6			

- Molecule 11 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
11	K	114	Total	C	N	O	S	0	0	0
			919	590	156	171	2			

- Molecule 12 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
12	L	46	Total	C	N	O	S	0	0	0
			363	224	72	63	4			

- Molecule 13 is a DNA chain called 5'-D(*AP*CP*TP*AP*CP*TP*TP*GP*AP*GP*CP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
13	N	7	Total	C	N	O	P	0	0	0
			137	68	22	41	6			

- Molecule 14 is a DNA chain called 5'-D(*AP*GP*CP*TP*C*AP*AP*GP*TP*AP*GP*TP*TP*AP*AP*GP*CP*CP*(BRU)P*GP*GP*TP*CP*AP*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
14	T	18	Total	Br	C	N	O	P	0	0
			369	1	176	69	106	17		

- Molecule 15 is a RNA chain called 5'-R(*UP*GP*CP*AP*UP*UP*U*CP*AP*AP*CP*CP

*AP*GP*GP*CP*UP*U)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
15	P	11	Total	C	N	O	P	0	0	0
			213	95	39	69	10			

- Molecule 16 is ZINC ION (CCD ID: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
16	A	2	Total	Zn	0	0
			2	2		
16	B	1	Total	Zn	0	0
			1	1		
16	C	1	Total	Zn	0	0
			1	1		
16	I	2	Total	Zn	0	0
			2	2		
16	J	1	Total	Zn	0	0
			1	1		
16	L	1	Total	Zn	0	0
			1	1		

- Molecule 17 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	1	Total	Mg	0	0
			1	1		









• Molecule 3: DNA-directed RNA polymerase II subunit RPB3

Chain C: 12% 46% 16% 23%




SER
ASN
MET
LEU
GLY
SER
ASN
GLU
ASP
VAL
MET
THR
GLY
ALA
GLU
GLN
ASP
PRO
TYR
SER
ASN
ALA
SER
GLN
MET
GLY
ASN
THR
GLY
TYR
ASP
ASP
ALA
TRP


● Molecule 4: DNA-directed RNA polymerase II subunit RPB4




● Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1




● Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2




• Molecule 7: DNA-directed RNA polymerase II subunit RPB7




• Molecule 8: DNA-directed RNA polymerases I, II, and III subunit RPABC3



• Molecule 9: DNA-directed RNA polymerase II subunit RPB9



• Molecule 10: DNA-directed RNA polymerases I, II, and III subunit RPABC5



• Molecule 11: DNA-directed RNA polymerase II subunit RPB11





- Molecule 12: DNA-directed RNA polymerases I, II, and III subunit RPABC4





- Molecule 13: 5'-D(*AP*CP*TP*AP*CP*TP*TP*GP*AP*GP*CP*T)-3'



- Molecule 14: 5'-D(*AP*GP*CP*TP*C*AP*AP*GP*TP*AP*GP*TP*TP*AP*AP*GP*CP*C P*(BRU)P*GP*GP*TP*CP*AP*TP*T)-3'



- Molecule 15: 5'-R(*UP*GP*CP*AP*UP*UP*U*CP*AP*AP*CP*CP*AP*GP*GP*CP*UP*U)-3'



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	221.62Å 393.72Å 282.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 3.65 50.00 – 3.65	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.00-3.65) 99.9 (50.00-3.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.64 (at 3.67Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.212 , 0.250 0.214 , 0.255	Depositor DCC
R_{free} test set	2686 reflections (1.97%)	wwPDB-VP
Wilson B-factor (Å ²)	86.1	Xtriage
Anisotropy	0.065	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 100.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.015 for 1/2*h-1/2*k,-3/2*h-1/2*k,-l 0.021 for 1/2*h+1/2*k,3/2*h-1/2*k,-l	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	31918	wwPDB-VP
Average B, all atoms (Å ²)	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BRU, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	2/11351 (0.0%)	1.19	117/15350 (0.8%)
2	B	0.64	0/9019	1.12	63/12160 (0.5%)
3	C	0.69	0/2133	1.20	32/2891 (1.1%)
4	D	0.59	0/1444	1.19	13/1935 (0.7%)
5	E	0.60	0/1788	1.19	24/2406 (1.0%)
6	F	0.80	0/717	1.34	10/967 (1.0%)
7	G	0.67	0/1368	1.22	11/1844 (0.6%)
8	H	0.57	0/1099	1.12	9/1488 (0.6%)
9	I	0.59	0/989	1.18	11/1331 (0.8%)
10	J	0.64	0/541	1.13	3/727 (0.4%)
11	K	0.60	0/937	1.08	3/1265 (0.2%)
12	L	0.72	0/365	1.23	4/485 (0.8%)
13	N	0.54	0/152	0.98	0/232
14	T	0.49	0/391	0.82	0/600
15	P	0.53	0/237	0.90	0/368
All	All	0.65	2/32531 (0.0%)	1.16	300/44049 (0.7%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	2
14	T	0	3
All	All	0	5

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	55	ASP	CA-C	-5.32	1.46	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1424	VAL	CA-CB	-5.27	1.48	1.54

The worst 5 of 300 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	56	PRO	N-CA-C	-17.16	94.50	114.92
4	D	26	THR	N-CA-C	-14.68	92.33	112.30
6	F	71	GLU	N-CA-C	-12.72	97.14	113.17
2	B	1185	CYS	N-CA-C	-12.22	98.55	114.31
5	E	5	ASN	N-CA-C	-11.78	98.89	113.97

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	1192	TYR	Sidechain
2	B	431	TYR	Sidechain
14	T	14	DA	Sidechain
14	T	16	DT	Sidechain
14	T	21	DC	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	11151	0	11226	1688	0
2	B	8847	0	8884	1444	0
3	C	2095	0	2051	369	0
4	D	1434	0	1460	227	0
5	E	1752	0	1776	283	0
6	F	705	0	731	118	0
7	G	1340	0	1357	188	0
8	H	1081	0	1051	213	0
9	I	971	0	929	158	0
10	J	532	0	542	120	0
11	K	919	0	929	127	0
12	L	363	0	388	98	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
13	N	137	0	82	10	0
14	T	369	0	202	27	0
15	P	213	0	110	19	0
16	A	2	0	0	0	0
16	B	1	0	0	0	0
16	C	1	0	0	0	0
16	I	2	0	0	0	0
16	J	1	0	0	0	0
16	L	1	0	0	0	0
17	A	1	0	0	0	0
All	All	31918	0	31718	4698	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 74.

The worst 5 of 4698 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:559:SER:HA	2:B:563:MET:HB3	1.20	1.16
2:B:1072:MET:HE3	2:B:1085:ILE:HB	1.28	1.13
12:L:55:ILE:HG12	12:L:56:LEU:H	0.97	1.12
12:L:55:ILE:CG1	12:L:56:LEU:H	1.59	1.12
3:C:43:THR:HG22	3:C:44:LEU:H	1.07	1.12

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	1407/1733 (81%)	977 (69%)	291 (21%)	139 (10%)	0 5
2	B	1097/1224 (90%)	736 (67%)	235 (21%)	126 (12%)	0 3

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	264/347 (76%)	182 (69%)	50 (19%)	32 (12%)	0	3
4	D	174/221 (79%)	116 (67%)	41 (24%)	17 (10%)	0	5
5	E	212/215 (99%)	141 (66%)	51 (24%)	20 (9%)	0	5
6	F	85/155 (55%)	66 (78%)	15 (18%)	4 (5%)	2	15
7	G	169/171 (99%)	138 (82%)	23 (14%)	8 (5%)	2	15
8	H	131/146 (90%)	84 (64%)	21 (16%)	26 (20%)	0	1
9	I	117/122 (96%)	80 (68%)	26 (22%)	11 (9%)	0	5
10	J	63/70 (90%)	39 (62%)	12 (19%)	12 (19%)	0	1
11	K	112/120 (93%)	82 (73%)	25 (22%)	5 (4%)	2	15
12	L	44/70 (63%)	25 (57%)	7 (16%)	12 (27%)	0	0
All	All	3875/4594 (84%)	2666 (69%)	797 (21%)	412 (11%)	0	4

5 of 412 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	4	GLN
1	A	5	GLN
1	A	54	ASN
1	A	58	LEU
1	A	62	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	1240/1520 (82%)	1096 (88%)	144 (12%)	5	23
2	B	965/1061 (91%)	826 (86%)	139 (14%)	3	16
3	C	234/299 (78%)	202 (86%)	32 (14%)	3	18
4	D	160/200 (80%)	138 (86%)	22 (14%)	3	18
5	E	196/197 (100%)	174 (89%)	22 (11%)	6	24
6	F	77/137 (56%)	68 (88%)	9 (12%)	5	23

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
7	G	152/152 (100%)	133 (88%)	19 (12%)	4	20
8	H	118/128 (92%)	98 (83%)	20 (17%)	2	12
9	I	113/116 (97%)	97 (86%)	16 (14%)	3	17
10	J	60/65 (92%)	55 (92%)	5 (8%)	10	33
11	K	99/102 (97%)	90 (91%)	9 (9%)	9	31
12	L	40/57 (70%)	36 (90%)	4 (10%)	7	28
All	All	3454/4034 (86%)	3013 (87%)	441 (13%)	4	20

5 of 441 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	806	THR
3	C	53	THR
12	L	54	ARG
8	H	89	LEU
2	B	878	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 141 such sidechains are listed below:

Mol	Chain	Res	Type
5	E	104	ASN
5	E	147	HIS
8	H	128	ASN
1	A	1218	GLN
1	A	1203	ASN

5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
15	P	10/18 (55%)	1 (10%)	1 (10%)

All (1) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
15	P	2	A

All (1) RNA pucker outliers are listed below:

Mol	Chain	Res	Type
15	P	1	C

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
14	BRU	T	23	15,14	18,21,22	3.90	1 (5%)	25,30,33	1.03	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
14	BRU	T	23	15,14	-	0/7/21/22	0/2/2/2

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
14	T	23	BRU	BR-C5	-16.43	1.50	1.88

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
14	T	23	BRU	C6-C5-C4	-2.86	117.77	120.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
14	T	23	BRU	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å ²)	Q<0.9
1	A	1417/1733 (81%)	0.08	35 (2%)	58	35	18, 67, 100, 123	0
2	B	1113/1224 (90%)	0.37	55 (4%)	35	21	21, 77, 109, 121	0
3	C	266/347 (76%)	0.00	1 (0%)	88	71	34, 66, 93, 107	0
4	D	178/221 (80%)	0.39	12 (6%)	24	16	46, 76, 106, 111	0
5	E	214/215 (99%)	0.44	7 (3%)	49	29	43, 88, 108, 117	0
6	F	87/155 (56%)	-0.28	2 (2%)	61	36	16, 44, 74, 84	0
7	G	171/171 (100%)	0.07	0	100	100	47, 63, 94, 101	0
8	H	135/146 (92%)	0.68	11 (8%)	18	13	71, 94, 110, 116	0
9	I	119/122 (97%)	0.62	7 (5%)	28	18	59, 92, 110, 119	0
10	J	65/70 (92%)	0.03	0	100	100	45, 62, 87, 98	0
11	K	114/120 (95%)	-0.23	1 (0%)	81	57	28, 67, 85, 96	0
12	L	46/70 (65%)	0.92	8 (17%)	4	4	51, 94, 110, 115	0
13	N	7/12 (58%)	2.24	4 (57%)	0	0	120, 125, 135, 137	0
14	T	17/26 (65%)	1.87	9 (52%)	0	0	99, 119, 135, 136	0
15	P	11/18 (61%)	1.75	3 (27%)	1	2	115, 119, 129, 139	0
All	All	3960/4650 (85%)	0.23	155 (3%)	43	26	16, 73, 107, 139	0

The worst 5 of 155 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	471	LYS	5.8
2	B	883	LEU	5.6
2	B	506	GLY	4.8
4	D	18	VAL	4.7
2	B	472	ALA	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
14	BRU	T	23	20/21	0.73	0.20	105,113,118,121	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
17	MG	A	2458	1/1	0.86	0.13	151,151,151,151	0
16	ZN	I	1122	1/1	0.95	0.08	131,131,131,131	0
16	ZN	L	1071	1/1	0.98	0.04	106,106,106,106	0
16	ZN	C	1269	1/1	0.99	0.11	40,40,40,40	0
16	ZN	A	2456	1/1	0.99	0.03	87,87,87,87	0
16	ZN	B	2225	1/1	1.00	0.02	42,42,42,42	0
16	ZN	J	1066	1/1	1.00	0.03	47,47,47,47	0
16	ZN	A	2457	1/1	1.00	0.08	42,42,42,42	0
16	ZN	I	1121	1/1	1.00	0.02	79,79,79,79	0

6.5 Other polymers [i](#)

There are no such residues in this entry.